Is living in a gas-flaring host community associated with being hypertensive? Evidence from the Niger Delta region of Nigeria

Omosivie Maduka, Charles Tobin-West

To cite: Maduka O, Tobin-West C. Is living in a gas-flaring host community associated with being hypertensive? Evidence from the Niger Delta region of Nigeria. *BMJ Glob Health* 2017;2:e000413. doi:10.1136/bmjgh-2017-000413

Handling editor Seye Abimbola

Received 18 May 2017
Revised 3 September 2017
Accepted 6 September 2017

ABSTRACT

Background Researchers have linked gas flaring to climate change, the hastening of the epidemiological transition and an upsurge in the prevalence of non-communicable diseases. We sought to determine if a relationship exists between residing in a gas-flaring host community and hypertension.

Methods We conducted an analytical cross-sectional household survey among residents of 600 households in three gas-flaring and three non-gas-flaring host communities in the Niger Delta region of Nigeria. We took geo-coordinates, administered a modified WHO-STEP questionnaire and built on Android mobile phones using Open-Data-Kit (ODK) software. We also took biological measurements and carried out descriptive and inferential statistical analysis using SPSS and STATA.

Results We interviewed a total of 912 adults: 437 (47.9%) from non-gas-flaring and 475 (52.1%) from gas-flaring host communities. There were differences in level of education ($\chi^2=42.99$; $p=0.00$), occupation category ($\chi^2=25.42$; $p=0.00$) and BMI category ($\chi^2=15.37$; $p=0.003$) among the two groups. The overall prevalence of hypertension was 23.7%: 20.7% among persons living in non-gas-flaring host communities compared with 25.3% among persons living in gas-flaring host communities ($\chi^2=2.89$; $p=0.89$). Residence in a gas-flaring host community, (AdjOR=1.75; 95% CI=1.11 to 2.74) and mean age (AdjOR=1.05; 95% CI=1.03 to 1.07) were identified as the predictors of hypertension. There was a significant association between hypertension and age, 1.05 (1.04–1.06) while the probability of being hypertensive was higher among residents of gas-flaring host communities between 20 to 40 years and 60 to 80 years.

Conclusion There is a need for the relevant agencies to scale up environmental and biological monitoring of air pollutants. The implication of a possible relationship between gas-flaring and hypertension brings to the fore the need for interventions to regulate gas-flaring activities.

INTRODUCTION

Nigeria is Africa's largest producer of crude oil and the sixth largest producer in the world,^{1–3} with a capacity to produce approximately 2.5 million barrels per day.⁴ The country has been producing crude oil since the 1950s when

Key questions

What is already known about this topic?

- ▶ Air pollution has been associated with an increase in the prevalence of hypertension and other non-communicable diseases.
- ▶ Communities in the Niger Delta which are host to gas-flaring activities are exposed to gas-flaring-induced air pollution.
- ▶ A previous study in the Niger Delta has linked oil pollution to increased prevalence of hypertension.

What are the new findings?

- ▶ Persons living in gas-flaring host communities are 1.75 times more likely to be hypertensive than persons resident in communities without oil exploration activities.
- ▶ The prevalence of hypertension in the study population was moderated by the age of study participants with those at both extremes of adult age residing in gas-flaring host communities being more hypertensive compared with similar persons in non-gas-flaring communities.

Recommendations for policy

- ▶ Study findings provide the impetus for stricter prevention and control measures for gas-flaring activities in the Niger Delta region of Nigeria.
- ▶ More attention need also be given to interventions for prevention and control of socio-economic lifestyle determinants of hypertension, especially among residents of oil-bearing communities.

deposits were first discovered in Oloibiri in what is now known as Bayelsa state. Oil and natural gas extraction currently account for up to 97% of the country's revenue from foreign exchange, 20% of the country's GDP and 65% of budgetary revenue. Nigeria is also blessed with huge deposits of natural gas located mainly in the Niger Delta region of the country.^{5,6} Despite this obvious wealth, Nigeria is ranked among the poorest countries in the world with over half of its population living on less than two dollars a day.⁶

CrossMark

Department of Preventive and Social Medicine, College of Health Sciences, University of Port Harcourt, Port Harcourt, Nigeria

Correspondence to

Dr Omosivie Maduka;
omosivie.maduka@uniport.edu.ng

Gas-flaring, which is routinely carried out by oil exploration companies in Nigeria, also poses a significant hazard to the health of populations exposed to it. It pollutes the air, heats up the atmosphere and releases greenhouse gases.^{7,8} Although Nigeria pledged to stop gas-flaring and has imposed fines on oil exploration companies still flaring gas in the country, the practice of gas-flaring has not stopped.⁹ Many of these companies admit that flaring of associated and non-associated gas is wrong and wasteful, but they also contend that the oil and gas sector in Nigeria faces infrastructure and technology obstacles in adequately harnessing natural gas. Consequently, Nigeria ranks as the 10th highest contributor to flaring of natural gas worldwide with flares of up to 426 billion cubic feet (bcf) of gas in 2013, representing 10% of gas-flaring worldwide.^{9,10} Many companies in the Niger Delta have only reduced flaring by a mere 10% since 2007 preferring to pay fines rather than to abate gas-flaring activities.^{8,9} The country had only succeeded in achieving a minimal reduction in gas-flaring from 540 bcf in 2010 to 428 bcf in 2013.¹⁰

The process of gas-flaring results in the dissemination of greenhouse gases (GHG) and other air pollutants such as carbon dioxide (CO₂), methane (CH₄), ethane, propane, butane, hydrogen sulphide (H₂S) and nitrous oxide (NO₂). Greenhouse gases have long been implicated in global warming and climate change.⁷ There is evidence of the adverse effects of heat-trapping gases on surface temperature.^{7,11} Apart from the effect on atmospheric temperature, the gases flared serve as pollutants to air and water. Greenhouse gases also precipitate the formation of acid rain. All this leads to negative outcomes in the health of the people who live and work in communities that are exposed to gas-flaring activities. These health outcomes range from chronic and recurrent respiratory diseases, abnormal haematological indices and increased susceptibility to blood dyscrasias, dermatological diseases and malignancies among others. The association between global warming and disease transmission has also been noted.^{7,8}

Several researchers have hinted at the relationship between air pollution and non-communicable diseases such as hypertension.^{12–21} However, this has not been demonstrated in the Niger Delta region of the country except for a recent publication which showed an association between residing in an oil-polluted environment and the prevalence of hypertension.²²

We determined the relationship between residence in a gas-flaring host community and the prevalence of hypertension in six Niger Delta communities in Nigeria. This research was part of a larger survey titled 'Epidemiological assessment of the health and well-being of residents of communities exposed to gas-flaring activities in the Niger Delta region of Nigeria' which explored morbidity and mortality patterns of persons living in gas-flaring host communities.

MATERIALS AND METHODS

The Niger Delta region of Nigeria as defined by the Nigerian government occupies about 70 000 km² and makes up 7.5% of Nigeria's land mass. It comprises the nine oil-producing states in the country namely: Bayelsa, Rivers, Delta, Akwa Ibom, Cross River, Edo, Abia, Imo and Ondo. These states are home to about 31 million people spanning more than 40 ethnic groups and 185 Local Government Areas (LGAs). The inhabitants speak about 250 different dialects. This research was part of a larger study conducted to compare morbidity and mortality of persons residing in gas-flaring host communities with those residing in communities where there are no oil exploration activities, using an analytical cross-sectional household survey. Rivers, Bayelsa and Delta states arguably have the highest number of oil exploration and gas-flaring activities in the Niger Delta of Nigeria and so were purposively sampled for the study. We chose two communities from each of the three states: one 'exposed' community and one 'un-exposed' community.

A total of six communities, Sampou and Nedugo in Bayelsa state, Ibada-Elume and Oton-Yasere in Delta state, and Omerelu and Mbodo-Aluu in Rivers state were selected. Nedugo, Oton-Yasere and Mbodo-Aluu were designated as gas-flaring host communities (host to gas-flaring activities for a minimum of 10 years with gas-flaring activities occurring within a five-kilometre radius from community settlements). Sampou, Ibada-Elume and Omerelu were designated as non-gas-flaring communities (no history of exposure to any oil exploration activities and not contiguous with exposed communities). These communities were, however, in the same state as exposed communities with similar population size, structure and socio-demographic characteristics as gas-flaring host communities. Individuals included in the household survey were all current residents of the selected communities, who were born into that community or had lived there for a minimum of 10 years.

A minimum sample size of 964 adults (482 resident in gas-flaring host communities and 482 resident in non-gas-flaring, and 880 not exposed) was considered adequate to determine a difference in prevalence of hypertension among the two groups of communities using the formula for difference between two proportions. This calculation was done using Winpepi version 9.7,²³ with power set at 80% ($u=0.84$) and a 5% significance level ($v=1.96$). From a previous study, the proportion of study participants in the oil-polluted community in Rivers state who were hypertensive (p_1) was 43.3%, while the proportion of hypertensive persons in the non-polluted community (p_2) was 31.2%.²² A 10% mark-up in sample size was done to allow for non-response and multiplication by a factor of two used to correct for design effect.

Based on an estimate of an average of two adults per household, a total of 600 households (100 households per community) were targeted for data collection. We visited each household once and studied all eligible and

available adults in 600 households. We did not conduct repeat visits. The geographical zones/villages in each community were identified based on an oral history provided by community gatekeepers such that the total number of households per community was further divided into these zones/villages. The first households studied in each zone/village was selected by tossing a coin between the two households closest to the village/zone centre. Subsequently, every household adjacent to the last sampled household was included if the residents met the eligibility criteria.

An interviewer-administered questionnaire adapted from the WHO STEPS questionnaire was used to obtain information on morbidity and mortality patterns in the study communities. It had the following sections: consent form, socio-demographic and household information, and physical measurements such as weight, height and blood pressure.

Every respondent had their blood pressure measured twice (at the start and end of the interview) using the Omron 3 Series Upper Arm Blood Pressure Monitor with wide-range cuff (BP710N). Appropriate cuffs sizes were used at all times, and blood pressure was taken

while the study participant was comfortably seated. The mean of the two readings (rounded off to the nearest whole number) was used. Hypertension was defined as systolic blood pressure ≥ 140 mmHg and/or diastolic BP ≥ 90 mmHg, or persons on anti-hypertensive medication irrespective of their observed blood pressure. Persons identified as hypertensive for the first time were referred to the nearest Primary Health Centre.

The weight of study participants was measured in kilograms using the Seca Clara 803 Digital Personal Scale. Study participants stood barefoot on the scale, and the weight was recorded by the research assistant based on readings from the scale. The Doran Scales Mechanical Height Rod Stadiometer was used to take all height measurements. Height was measured in metres and measurements taken barefoot with the study participants standing up straight against the vertical ruler.

The data collection tool was built and uploaded on Android mobile phones using the Open Data Kit software. A server domiciled in the WHO state office was set up for real-time upload and storage of completed questionnaires. Six Android mobile phones which are GPS-enabled were used for data collection including the

Table 1 Socio-demographic characteristics of persons residing in gas-flaring host communities compared with non-gas-flaring communities

Characteristics	Non gas-flaring	Gas-flaring	Chi-square (p Value)
Sex (n=912)			
Male	148 (33.9)	178 (37.5)	1.29 (0.26)
Female	289 (66.1)	297 (62.5)	
Level of education (n=836)			
Primary	84 (21.6)	49 (10.9)	*42.99 (0.00)
Secondary	257 (66.2)	27 (60.5)	
Tertiary	47 (12.1)	128 (28.6)	
Marital status (n=912)			
Married/living with partner	306 (70.0)	326 (68.6)	6.44 (0.09)
Widowed	24 (5.5)	16 (3.4)	
Divorced/separated	11 (2.5)	6 (1.3)	
Single	96 (22.0)	127 (26.7)	
Occupation category (n=912)			
Unemployed/student/pensioner	168 (51.7)	142 (42.5)	25.42 (0.00)
Unskilled	95 (29.2)	70 (21.0)	
Skilled	56 (17.2)	110 (32.9)	
Professional	6 (1.8)	12 (3.6)	
BMI category			
Underweight	63 (14.4)	51 (10.7)	*15.37 (0.003)
Normal	245 (56.1)	226 (47.6)	
Overweight	74 (16.9)	107 (22.5)	
Obese	55 (12.6)	91 (19.2)	
Mean age (SD)†	36.6 (16.1)	35.6 (14.0)	0.97 (0.33)

*significant Chi-square values at $p < 0.05$, †t-test (p-value).

capture of the GPS coordinates for each household. We recruited and trained six field data collectors who were university graduates to administer the study tool. These data collectors were blinded to the objectives of the study and the purpose of the physical measurements as a means of controlling measurement bias. In each community, we further recruited another five persons who were indigenes to work with each data collector (making up a two-person team of one data collector and one indigene). The local recruits served as interpreters/guides/assistants to each data collector. Data collection was carried out over 14 working days in May 2016 with an average of 2 days spent in each community. Each data collection team collected data from a minimum of 20 households per day.

DATA ANALYSIS

Data was downloaded from the WHO server in Microsoft Excel 2013 version for data cleaning and analysed using Statistical Package for the Social Sciences (SPSS) version

21 and STATA version 14 with descriptive and inferential statistics carried out, Chi-square test, t-tests, analysis of variance, simple logistic and multiple logistic regression. We determined the fitness of the regression model using Hosmer–Lemeshow goodness-of-fit test regression diagnostics. We also performed the tolerance test for multicollinearity, link test to check for model specification error and plotted the receiver operating characteristic curves (c-statistics) analyses to determine the predictive powers of the final multivariate model. Results were presented as odds ratios (ORs) with 95% confidence intervals (CIs). The significance tests were 2-tailed, and statistical significance was defined at the alpha level of 0.05.

ETHICAL CONSIDERATIONS

Ethical approval for the research was obtained from the research ethics committee of the University of Port Harcourt. A consent form explaining the study objectives, procedures, benefits and risks was prepared for all potential study participants to review and give written informed consent

Table 2 Prevalence of hypertension by socio-demographic characteristics in the study population

	Study category		Chi-square (p Value)
	Non gas-flaring (n=93)	Gas-flaring (n=123)	
Hypertension status			
Hypertensive	93 (20.7)	123 (25.3)	2.89 (0.89)
Sex			
Male	39 (26.4)	45 (25.6)	0.03 (0.87)
Female	54 (18.8)	18 (26.4)	*4.82 (0.03)
Level of education			
Primary	23 (27.4)	18 (37.5)	1.46 (0.23)
Secondary	39 (15.2)	56 (20.7)	2.69 (0.10)
Tertiary	10 (21.3)	38 (29.9)	1.28 (0.28)
Non-response	21 (42.9)	11 (40.7)	0.32 (0.86)
Occupation			
Unemployed/student	39 (23.3)	42 (29.8)	1.71 (0.19)
Unskilled	15 (15.8)	20 (29.0)	*4.15 (0.04)
Skilled	12 (21.8)	22 (19.3)	0.11 (0.7)
Professional	2 (33.3)	4 (33.3)	0.00 (1.00)
Age group			
18 to 34 years	24 (9.5)	45 (16.1)	*5.14 (0.02)
35 to 54 years	38 (30.9)	46 (33.8)	0.25 (0.62)
55 to 74 years	23 (50.0)	27 (57.4)	0.52 (0.47)
75 years and above	8 (57.1)	5 (55.6)	0.006 (0.94)
Body mass index			
Underweight	13 (20.6)	5 (9.8)	1.49 (0.12)
Normal	40 (16.3)	46 (20.5)	1.38 (0.24)
Overweight	21 (28.8)	34 (31.8)	0.19 (0.67)
Obese	19 (34.5)	38 (42.2)	0.84 (0.36)

*significant Chi-square values at $p < 0.05$.

(by signature or thumb print). Permission to carry out the study was obtained from the local government headquarters of each community and the community gatekeepers. Study participants were assured of confidentiality. Data collection tools did not bear participants names. All records generated remained in the custody of the researchers and were kept in a secure location. Data entry files were passworded to further guarantee confidentiality. Study participants were also informed of their rights to withdraw consent at any time during the study without concern for negative consequences.

RESULTS

Socio-demographic characteristics of study participants

A total of 912 adults (aged 18 years and above) were interviewed during the household survey giving a response rate of 94.6%. A total of 437 (47.9%) of these were residents of communities not exposed to gas-flaring, while 475 (52.1%) persons were residents of communities exposed to gas-flaring activities. There was no difference in the mean age (SD) of respondents in communities not exposed to gas-flaring (36.6; 16.1 years) compared with respondents residing in communities exposed to gas-flaring activities (35.6; 14.0 years). A comparison of the distribution of socio-demographic characteristics across study categories

revealed differences in level of education ($\chi^2=43.0$; $p=0.00$), occupation category ($\chi^2=25.4$; $p=0.00$) and BMI category ($\chi^2=15.4$; 0.003) [table 1](#).

Prevalence of hypertension

A total of 216 persons among the adults in the study were found to be hypertensive, giving an overall prevalence of 23.7%. A total of 93 (43.06%) hypertensive persons were residents of communities not exposed to gas-flaring, while 123 (56.94%) hypertensive persons were residents of communities exposed to gas-flaring activities. These correspond to a prevalence of hypertension among persons resident in non-gas-flaring communities of 20.7% compared with 25.3% prevalence of hypertension among persons living in communities exposed to gas-flaring. These two prevalences were however not significantly different 2.89 (0.89). A comparison of prevalence of hypertension across socio-demographic characteristics is shown in [table 2](#). Females ($\chi^2=4.82$; 0.03), unskilled workers ($\chi^2=4.15$; 0.04) and persons aged 18–34 years ($\chi^2=5.14$; 0.02), living in communities exposed to gas-flaring activities had a higher prevalence of hypertension than those residing in communities not exposed to gas-flaring activities [table 2](#).

Table 3 Predictors of hypertension using regression models

	Unadjusted		Adjusted	
Variable	OR (95% CI)	p Value	OR (95% CI)	p Value
Study category				
Residence in a gas-flaring versus non- gas-flaring host community	1.31 (0.96 to 1.76)	0.090	*1.75 (1.11 to 2.74)	0.02
Mean age	*1.05 (1.04 to 1.06)	0.000	*1.05 (1.03 to 1.07)	0.00
Sex				
Female versus male	0.82 (0.60 to 1.12)	0.20	0.9 (0.57 to 1.42)	0.65
BMI	1.01 (1.00 to 1.02)	0.00	1.01 (0.99 to 1.02)	0.11
Levels of education				
Primary (reference)	Reference		Reference	
Junior secondary	0.2 (0.09 to 0.44)	0.000	0.51 (0.18 to 1.39)	0.19
Senior secondary	0.52 (0.34 to 0.80)	0.003	1.08 (0.59 to 1.99)	0.80
Tertiary	0.81 (0.49 to 1.32)	0.399	1.33 (0.67 to 2.64)	0.42
Marital status				
Single (Reference)	Reference		Reference	
Married/living with partner	*2.96 (1.92 to 4.55)	0.000	1.59 (0.88 to 2.89)	0.13
Widowed	*8.39 (4.03 to 17.47)	0.000	1.45 (0.28 to 7.58)	0.66
Divorced/separated	*5.87 (2.07 to 16.66)	0.001	3.95 (0.97 to 16.12)	0.06
Occupation				
Unemployed/student (reference)	Reference		Reference	
Unskilled	0.75 (0.48 to 1.17)	0.203	0.73 (0.420 to 1.25)	0.25
Skilled	0.73 (0.46 to 1.14)	0.165	0.68 (0.40 to 1.15)	0.15
Professional	1.46 (0.53 to 4.02)	0.460	0.90 (0.28 to 2.92)	0.86

*Confidence Intervals showing significant association between variables and hypertension status.

Figure 1 Predicted probability of hypertension for all ages by gas-flaring status.

Association between hypertension and residence in gas-flaring host community

We carried out bivariate and multivariate regression of the following independent variables found to be associated with hypertension status (dependent variable) at $p < 0.25$, gas-flaring status of the community of residence, age, sex, BMI, marital status, the level of education and category of occupation. Bivariate regression analysis showed that mean age (OR=1.05; 95% CI=1.04 to 1.06), the status of being married/living with partner (OR=2.96; 95% CI=1.92 to 4.55), being divorced/separated (OR=8.39; 95% CI=4.03 to 17.47) and being widowed (OR=5.87; 95% CI=2.07 to 16.66) were related to being hypertensive. Table 3 However the multivariate regression model adjusted for confounders and identified residence in a community exposed to gas-flaring (AdjOR=1.75; 95% CI=1.11 to 2.74) and mean age (AdjOR=1.05; 95%

CI=1.03 to 1.07) as the only significant predictors of hypertension among the study participants table 3.

Further analysis was done to determine the interaction effects and fit of the model between hypertension and gas-flaring status, and hypertension status and age. The interaction between hypertension status and age O.R 1.05 (1.04–1.06) was found to be significant, while that between hypertension and gas-flaring status was not O.R 1.30 (0.96–1.76). This finding was supported by the area under the ROC curve which was 0.533 for hypertension versus gas-flaring status and 0.737 for hypertension versus age. Other model fit statistics are as shown in table 4. The probability of being hypertensive was modified by age (figure 1) such that the probability of being hypertensive was higher among residents of gas-flaring host communities between 20 to 40 years and 60 to 80 years.

DISCUSSION

Principal findings

Our study compared hypertension status of persons resident in gas-flaring host communities with persons resident in communities in which oil exploration had never occurred. Findings from this study indicate that about one out of four adults interviewed in all six communities was hypertensive. About one-quarter and one-fifth of respondents were found to be hypertensive in gas-flaring and non-gas-flaring communities respectively. This observed difference was not significant. However, after adjusting for confounding variables, age and residence in a gas-flaring host community were found to be the strongest predictors of hypertension status. Residents in a gas-flaring host community were 1.75 times more likely to be hypertensive than residents in non-gas-flaring communities. The interaction effects and model fit statistics identified age as the stronger predictor of hypertension. The age-adjusted probability of being hypertensive among those 20 to 40 and 60 to 80 years, was found to be higher among residents of a gas-flaring host community compared with residents of a non-gas-flaring host community.

Implications

A previous study done in other communities in the Niger Delta identified residence in a crude oil-polluted environment as a strong predictor of hypertension. It also identified age as the most important moderating variable in the relationship between oil pollution and hypertension.²² This identified relationship and interactions between hypertension status, age and residence in a gas-flaring host community may be explained by observed interactions between exposure to air pollution and metabolic changes underlying the development of hypertension, cardiovascular disease and diabetes observed by previous air pollution studies.^{12–15 17 18 20 21} The report of the United Nations Environment Programme (UNEP) in Ogoniland in South-South Nigeria²⁴ and other research

Table 4 Interaction of demographic variables and hypertensive status

Interaction effect	Odds ratio (95% CI)	p Values
Hypertensive (vs non-hypertensive) x gas-flaring	1.30 (0.96 to 1.76)	0.09
Model fit statistics		
Area under ROC curve		0.53
Lemeshow test (p Value)		1.00
Link test (p Values)		
Collinearity diagnostic (mean VIF)		1.00
Hypertensive (vs non-hypertensive) x age	1.05 (1.04 to 1.06)	<0.001
Model fit statistics		
Area under ROC curve		0.74
Lemeshow test (p Value)		0.19
Link test (p Values)		
Collinearity diagnostic (mean VIF)		1.00

in the Ogoni axis in South-South Nigeria also support biological plausibility for this phenomenon.^{25–27}

An alternative theory explaining the observation may be an increase in the adoption of negative lifestyle changes relating to improved socio-economic status resulting from the economic advantages that go along with residing in oilproducing communities.²² Study participants residing in gas-flaring host communities were significantly better educated, had a higher proportion of skilled jobs and were more overweight and obese than study participants residing in non-gas-flaring communities. This observation is in keeping with the theory of lifestyle-related risk factors strongly associated with non-communicable diseases.^{28–34}

Limitations and strengths

This study did not evaluate the relationship between socio-economic status, modifiable risk factors, hypertension and residence in gas-flaring host communities. However, a previous study that did evaluate this relationship did not find any modifiable risk factor to have a significant association with oil pollution and hypertension.²² Our study also did not employ a longitudinal study design that is preferable for establishing temporal relationships. However, this study had a large sample size and employed rigorous sampling and statistical analysis. Finally, our study was unable, due to funding constraints, to conduct bio-monitoring of study participants to elicit the presence of specific pollutants. Further longitudinal research is needed in this regard to determine the long-term attributable effects of gas flaring on the health of persons living in the Niger Delta region of Nigeria

CONCLUSION

The implication of a significant relationship between gas-flaring and hypertension brings to the fore the need for regulatory activities that prevent and stop gas-flaring activities. There is also the need for greater efforts by government regulatory bodies such as the ministries of environment and health to scale up environmental and biological monitoring of pollutants.

Acknowledgements We acknowledge funding support from the Department for International Development (DfID) under the Climate Impacts Research Capacity and Leadership Enhancement (CIRCLE) Program. We also acknowledge the University of Port Harcourt (UNIPORT) and its teaching Hospital (UPH) for approving a one-year study leave to enable OM to undertake research activities.

Contributors OM wrote the protocol, conducted the research, did the data analysis and wrote the first draft of the manuscript. CT -W co-wrote the protocol, supported the data collection and revised the manuscript prior to submission.

Funding This research is supported by funding from the Department for International Development (DfID) under the Climate Impact Research Capacity and Leadership Enhancement (CIRCLE) programme.

Competing interests None declared.

Patient consent This was a population-based study. Client forms were signed by each study participant prior to administration of study tools.

Ethics approval Research Ethics Committee of the University of Port Harcourt.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement Patient level data and full dataset is available from the corresponding author at omosivie.maduka@uniport.edu.ng. The presented data and data set are anonymised and risk of identification is low.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Oni SI, Oyewo MA. Gas flaring, transportation and sustainable energy development in the Niger-Delta, Nigeria. *J Hum Ecol* 2011;33:21–8.
- Ikeke MO. Thomas Berry's idea of technological transformation its relevance to the management of oil technology in Nigeria's Niger delta. *Thought Pract Assoc Kenya* 2013;5:141–58.
- Egwurugwu JN, Nwafor A, Ezekwe S. Effects of prolonged exposure to gas flares on the lipid profile of humans in the Niger delta region, Nigeria. *Arch Appl Sci Res* 2013;5:98–104.
- Nigerian National Petroleum Corporation. Oil Production in Nigeria. 2015 <http://www.nnpcgroup.com/nnpcbuisness/upstreamventures/oilproduction.aspx> (accessed 7 May 2015).
- American Association for the Advancement of Science. *Geospatial Technologies and Human Rights Project: Eyes of Nigeria*, 2011.
- Friends of the Earth Limited. *Gas flaring in Nigeria*, 2004. Media Report 1649.
- World Health Organization. *Climate Change and Human Health: Risk and Responses*. World Health Organization, 2003.
- E. Ite A, J. Ibok U. Gas flaring and venting associated with petroleum exploration and production in the Nigeria's Niger delta. *Amer J Envi Prot* 2013;1:70–7.
- Vidal J. Gas flaring is wasting fuel and fueling climate change | climate central. *Guardian Newspaper* 2012 <http://www.climatecentral.org/news/gas-flaring-is-wasting-fuel-and-fueling-climate-change-15163> (accessed 7 May 2015).
- U.S Energy Information Administration (EIA). Nigeria: Country Analysis Brief Overview. US EIA Independent statistics and analysis, 2015:1–20. <http://www.eia.gov/countries/country-data.cfm?fips=ni> (accessed 7 May 2015).
- Ajugwo aO. Negative effects of gas flaring: the Nigerian experience. *J Environ Pollut Hum Heal* 2013;1:6–8.
- Foraster M, Basagaña X, Aguilera I, et al. Association of long-term exposure to traffic-related air pollution with blood pressure and hypertension in an adult population-based cohort in Spain (the REGICOR study). *Environ Health Perspect* 2014;122:404–11.
- Foraster M, Basagaña X, Aguilera I, et al. Association of long-term exposure to traffic-related air pollution with blood pressure and hypertension in an adult population-based cohort in Spain (the REGICOR study). *Environ Health Perspect* 2014;122:404.
- Sughis M, Nawrot TS, Ihsan-ul-Haque S, et al. Blood pressure and particulate air pollution in schoolchildren of Lahore, Pakistan. *BMC Public Health* 2012;12:378.
- Gobo A, Richard G, Ubong I. Health impact of gas flares on Ikwuruta/Umuochem communities in Rivers State. *J Appl Sci Environ Manage* 2009;13:27–33.
- Krewski D, Burnett R, Jerrett M, et al. Mortality and long-term exposure to ambient air pollution: ongoing analyses based on the American Cancer Society cohort. *J Toxicol Environ Health A* 2005;68:1093–109.
- Pope C, Burnett R, Thurston G, et al. Cardiovascular Mortality and Long-Term Exposure to Particulate Air Pollution. *Circulation* 2004;109:71–7.
- Coogan PF, White LF, Jerrett M, et al. Air pollution and incidence of hypertension and diabetes mellitus in black women living in Los Angeles. *Circulation* 2012;125:767–72.
- Dong GH, Qian ZM, Xaverius PK, et al. Association between long-term air pollution and increased blood pressure and hypertension in China. *Hypertension* 2013;61:578–84.
- Chuang KJ, Yan YH, Chiu SY, et al. Long-term air pollution exposure and risk factors for cardiovascular diseases among the elderly in Taiwan. *Occup Environ Med* 2011;68:64–8.

21. Chuang KJ, Yan YH, Cheng TJ. Effect of air pollution on blood pressure, blood lipids, and blood sugar: a population-based approach. *J Occup Environ Med* 2010;52:258–62.
22. Ezejimofor MC, Uthman OA, Maduka O, *et al*. The burden of hypertension in an oil- and gas-polluted environment: a comparative cross-sectional study. *Am J Hypertens* 2016;29:925–33.
23. Abramson JH. WINPEPI updated: computer programs for epidemiologists, and their teaching potential. *Epidemiol Perspect Innov* 2011;8:1.
24. United Nations Environment Programme. *Environmental assessment of Ogoniland*. United Nations Environment programme, 2011.
25. Lindén O, Pålsson J. Oil contamination in Ogoniland, Niger delta. *Ambio* 2013;42:685–701.
26. Okparanma RN, Coulon F, Mouazen AM. Analysis of petroleum-contaminated soils by diffuse reflectance spectroscopy and sequential ultrasonic solvent extraction-gas chromatography. *Environ Pollut* 2014;184:298–305.
27. Kponee KZ, Chiger A, Kakulu II, *et al*. Petroleum contaminated water and health symptoms: a cross-sectional pilot study in a rural Nigerian community. *Environ Health* 2015;14:86.
28. Kinra S, Bowen LJ, Lyngdoh T, *et al*. Sociodemographic patterning of non-communicable disease risk factors in rural India: a cross sectional study. *BMJ* 2010;341:c4974.
29. Murray CJ, Lopez AD. Global mortality, disability, and the contribution of risk factors: Global Burden of Disease Study. *Lancet* 1997;349:1436–42.
30. Beaglehole R, Yach D. Globalisation and the prevention and control of non-communicable disease: the neglected chronic diseases of adults. *Lancet* 2003;362:903–8.
31. Lopez AD, Mathers CD, Ezzati M, *et al*. Global and regional burden of disease and risk factors, 2001: systematic analysis of population health data. *Lancet* 2006;367:1747–57.
32. Ezzati M, Lopez AD, Rodgers A, *et al*. Selected major risk factors and global and regional burden of disease. *Lancet* 2002;360:1347–60.
33. Murray CJL, Lopez AD. *The global burden of disease: a comprehensive assessment of mortality and disability from diseases, injuries and risk factors in 1990 and projected to 2010*. 1. Harvard University Press, 1996:1–35.
34. World Health Organization. Noncommunicable diseases: the slow motion disaster. *WHO* 2017 <http://www.who.int/publications/10-year-review/ncd/en/index4.html> (accessed 7 May 2015).