(ST152), t186 (ST88), and t346 dominating. While many distinct strains were isolated from both health centres, genotype clustering was identified within centres pointing to possible health care-associated transmission. Phylogenomic analysis confirmed these clusters. Among the GNB, phenotype screening showed widespread resistance to ampicillin, chloramphenicol, ticarcillin-clavulanic acid, cefuroxime and sulphamethoxazole-trimethoprim. ESBL production was confirmed in 15 isolates phenotypically while 61.5% of screen-positive isolates harboured at least one ESBL-conferring gene. Carbapenem encoding genes were detected in 41% of the isolates.

Conclusions Our findings indicate that the health-care environment likely contributes to superinfection of BU wounds and calls for training in wound management and infection control techniques. The observed frequency of ESBL and carbapenem resistance indicates the need to set up surveillance networks and strictly enforce policies which guide the rational use of antibiotics.

PA-103 **DRUG RESISTANCE AND GENETIC PROFILE OF BACTERIAL SPECIES ASSOCIATED WITH BURULI ULCER WOUND INFECTIONS IN TWO DISTRICTS OF GHANA**

Grace Kpeli,¹ Isaac Otchere,¹ Araceli Lamelas,² Andrew Buultjens,³ Dieter Bulach,³ Sarah Baines,³ Torsten Seemann,³ Stefano Giuliani,³ Zuliehatu Nakobu,¹ Samuel Aboagye,¹ Evelyn Owusu-Mireku,¹ Emelia Danso,¹ Julia Hauser,⁴ Vladimira Hinic,⁵ Gerd Pluschke,⁴ Timothy Stinear,³ Dorothy Yeboah-Manu¹. ¹NMIMR, Ghana; ²Red de Estudios Moleculares Avanzados, Instituto de Ecología A.C., Mexico; ³Doherty Institute, University of Melbourne, Australia; ⁴Swiss TPH, Switzerland; ⁵University Hospital Basel, Switzerland

10.1136/bmjgh-2016-000260.133

Background We identified secondary infection of Buruli ulcer (BU) wounds as a cause of healing delay. In order to contribute to the improvement of wound management and reduction of healing delay, we initiated a study to gain understanding of the possible routes of infection and also characterised the resistant profiles of Gram negative bacteria isolated from the wounds of patients attending two health facilities in Ghana.

Methods *Staphylococcus aureus* isolates were characterised by the *spa* gene, *mecA* and the Pantone Valentine Leukocidin (PVL) toxin followed by *spa* sequencing and whole genome sequencing of a subset of isolates. Phenotypic antibiotic susceptibility testing of Gram negative clinical isolates was performed and multidrug-resistant *Pseudomonas aeruginosa* identified. The *Enterobacteriaceae* were further investigated for ESBL and carbapenem production, and some resistance conferring genes were analysed by PCR.

Results Twenty-four isolates were identified as methicillin-resistant *S. aureus* (MRSA), and *lukFS* genes encoding PVL were identified in 67 isolates. Typing and sequencing of the *spa* gene from 91 isolates identified 29 different *spa* types with t355