

Supplementary File 2

Table S2. Associations between reported pregnancy complications and attendance for at least one antenatal care visit

Variable		Attended any ANC	Crude OR	Adjusted OR ¹	95% CIca of adjusted OR	p-value from test of heterogeneity of strata ²
Persistent headache	No	2108/2489	0.80	0.50	0.25-0.98	0.826
	Yes	1015/1162				
Dizziness or blurred vision	No	2400/2798	1.08	0.79	0.65-0.96	0.303
	Yes	723/853				
Swelling of face or hands	No	2638/3069	1.22	0.96	0.89-1.04	0.301
	Yes	485/97				
Vaginal bleeding	No	2951/3432	1.68	1.43	0.90-2.26	0.000
	Yes	172/219				
Convulsions	No	3033/3547	0.92	0.77	0.40-1.49	0.084
	Yes	90/104				

1 Adjusted OR after stratification by intervention status

2 Tests significance of difference in OR for association with ANC attendance between intervention and control wards

OR=Mantel Haenszel odds ratio; 95% CIca = cluster-adjusted 95% confidence interval

Bold font indicates an association significant at the 5% level

Table S3. Associations between reported pregnancy complications and delivery in a health facility

Variable		Delivery in health facility	Crude OR	Adjusted OR ¹	95% CIca of adjusted OR	p-value from test of heterogeneity of strata ²
Persistent headache	No	619/2253	1.33	1.03	0.86-1.23	0.239
	Yes	247/1111				
Dizziness or blurred vision	No	670/2552	1.12	0.84	0.70-1.01	0.773
	Yes	196/812				
Swelling of face or hands	No	732/2816	1.09	0.86	0.72-1.03	0.723
	Yes	134/548				
Vaginal bleeding	No	801/3183	0.60	0.50	0.14-1.76	0.078
	Yes	65/181				
Convulsions	No	827/3271	0.47	0.40	0.08-1.88	0.003
	Yes	39/93				
High blood pressure ³	No	631/2466	0.87	0.76	0.50-1.17	0.193
	Yes	73/258				

1 Adjusted OR after stratification by intervention status

2 Tests significance of difference in OR for association with facility delivery between intervention and control wards

3 Among those who reported any measurement of blood pressure

OR=Mantel Haenszel odds ratio; 95% CIca = cluster-adjusted 95% confidence interval