

## Supplementary material

### Appendix 5. Participants responses to NCD survey questions by study arm in Bangladesh and Uganda

#### Bangladesh

Indicator	Control	50 Taka	100 Taka	p-value
Current smoker	144/339 (42.5%)	142/362 (39.2%)	165/351 (47.0%)	0.109
Current smokeless tobacco	62/348 (17.8%)	69/366 (18.9%)	54/357 (15.1%)	0.394
Alcohol abstainer	277/348 (79.6%)	293/362 (80.9%)	275/355 (77.5%)	0.512
Current alcohol drinker	23/348 (6.6%)	25/362 (6.9%)	27/355 (7.6%)	0.868
No fruit in past week	58/351 (16.5%)	52/366 (14.2%)	44/360 (12.2%)	0.261
No vegetables in past week	28/335 (8.4%)	27/353 (7.7%)	26/344 (7.6%)	0.914
No vigorous physical activity in past week	132/346 (38.2%)	138/363 (38.0%)	127/357 (35.6%)	0.726
No moderate physical activity in past week	108/345 (31.3%)	131/363 (36.1%)	109/351 (31.1%)	0.271
Had blood pressure measured	155/336 (46.1%)	190/362 (52.5%)	187/352 (53.1%)	0.129
Had hypertension diagnosis	38/333 (11.4%)	52/362 (14.3%)	40/351 (11.4%)	0.385
Had diabetes diagnosis	27/340 (7.9%)	19/362 (5.3%)	20/357 (5.6%)	0.281

#### Uganda

Indicator	Control	5000 UGX	10,000 UGX	p-value
Current smoker	28/345 (8.1%)	31/371 (8.4%)	34/378 (9.0%)	0.907
Current smokeless tobacco	20/358 (5.6%)	26/380 (6.8%)	33/398 (8.3%)	0.343
Alcohol abstainer	151/366 (41.3%)	163/395 (41.3%)	178/410 (43.4%)	0.776
Current alcohol drinker	108/365 (29.6%)	135/395 (34.2%)	125/409 (30.6%)	0.350
No fruit in past week	102/373 (27.4%)	100/402 (24.9%)	92/418 (22.0%)	0.219
No vegetables in past week	88/363 (24.2%)	92/399 (23.1%)	79/413 (19.1%)	0.192
No vigorous physical activity in past week	102/375 (27.2%)	89/397 (22.4%)	112/415 (27.0%)	0.219
No moderate physical activity in past week	76/370 (20.5%)	73/397 (18.4%)	60/414 (14.5%)	0.078
Had blood pressure measured	217/363 (59.7%)	212/384 (55.2%)	237/394 (60.2%)	0.302
Had hypertension diagnosis	39/363 (10.7%)	33/384 (8.6%)	41/394 (10.4%)	0.566
Had diabetes diagnosis	18/361 (5.0%)	20/393 (5.1%)	24/405 (5.9%)	0.814

Abbreviations: UGX, Ugandan Shilling