Formulating questions to explore complex interventions within qualitative evidence synthesis

Andrew Booth,¹ Jane Noyes,² Kate Flemming,³ Graham Moore,⁴ Özge Tunçalp,⁵ Elham Shakibazadeh⁶

To cite: Booth A, Noyes J, Flemming K, *et al.* Formulating questions to explore complex interventions within qualitative evidence synthesis. *BMJ Glob Health* 2019;4:e001107. doi:10.1136/bmjgh-2018-001107

Handling editor Soumyadeep Bhaumik

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/bmjgh-2018-001107>).

Received 12 August 2018
Revised 1 November 2018
Accepted 13 November 2018

ABSTRACT

When making decisions about complex interventions, guideline development groups need to factor in the sociocultural acceptability of an intervention, as well as contextual factors that impact on the feasibility of that intervention. Qualitative evidence synthesis offers one method of exploring these issues. This paper considers the extent to which current methods of question formulation are meeting this challenge. It builds on a rapid review of 38 different frameworks for formulating questions. To be useful, a question framework should recognise context (as setting, environment or context); acknowledge the criticality of different stakeholder perspectives (differentiated from the target population); accommodate elements of time/timing and place; be sensitive to qualitative data (eg, eliciting themes or findings). None of the identified frameworks satisfied all four of these criteria. An innovative question framework, PerSPECtIF, is proposed and retrospectively applied to a published WHO guideline for a complex intervention. Further testing and evaluation of the PerSPECtIF framework is required.

INTRODUCTION

Many commentators seek to define complexity in connection with complex interventions.¹ Rogers distinguishes between simple, complicated and complex: simple is encapsulated in following a recipe, complicated by sending a rocket to the moon and complex in bringing up a child.² In the first paper of this WHO series on Complex Interventions, Petticrew and colleagues turn the emphasis away from the activity itself (intervention or exposure) and towards the perspective adopted by the evaluator (in this case an ‘intervention perspective’ or a ‘systems perspective’).¹ These perspectives offer alternative evaluation ‘lenses’ to be adopted by reviewers or guideline developers even when examining the same phenomenon. For example, when examining use of a safety checklist within operating theatres³ one could either adopt an ‘intervention perspective’ or ‘lens’ to consider issues that relate to implementation within a controlled setting

Summary box

- ⇒ Qualitative evidence syntheses are useful for exploring the effects of complex interventions and their subsequent implementation.
- ⇒ Question formulation is critical to the systematic review and associated guideline processes yet current frameworks do not accommodate a complex systems or complex intervention perspective.
- ⇒ We review existing question formulation frameworks for qualitative evidence synthesis and report on their suitability for formulating questions to explore complex interventions.
- ⇒ We propose an alternative question formulation framework, the PerSPECtIF framework, to be further tested, to accommodate context, perspective, time and space within a complex system perspective

(the theatre) or adopt a ‘systems perspective’ or lens to explore the wider organisational or system culture within which the checklist is being implemented (eg, within a culture of blame or of improvement).

When making decisions about complex interventions, guideline development groups need to take account of the sociocultural acceptability of the intervention, as well as how feasible the intervention will be to implement. Complex interventions are inextricably linked to context; interventions interact with, and sometimes change, the context within which they are implemented.¹ Recognition that complex interventions are context-dependent not only holds implications for the effect of the intervention, but also for its sustainability, acceptability and feasibility. This paper examines implications of adopting a ‘systems perspective’, as opposed to an ‘interventions perspective’, when formulating questions to be addressed by qualitative research. As with the first paper in the WHO Complex Interventions series,¹ it focuses on the first part of the evidence synthesis process, defining the question. This

© World Health Organization 2019. Licensee BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Andrew Booth;
A.Booth@sheffield.ac.uk

Box 1 Complexity-related questions to be addressed in a qualitative evidence synthesis (QES)

Potential research questions for a QES

- ⇒ How do the components work along and in combination to produce effects?
- ⇒ How do they interact to produce outcomes?
- ⇒ How and why does the implementation of the intervention vary across contexts?
- ⇒ How does the system change when the intervention is introduced?
- ⇒ What are the effects (anticipated and unanticipated) which follow from this system change?
- ⇒ How do effects change over time? (Changes may relate to biological, ecological, epidemiological or social factors)
- ⇒ What explains how effectiveness of the intervention changes over time?
- ⇒ What factors enable or inhibit implementation of interventions?
- ⇒ What changes in processes and outcomes follow the introduction of this system change?
- ⇒ At what levels in the system are they experienced? (eg, individuals, families, communities)
- ⇒ To what extent do patients/beneficiaries value different health outcomes?
- ⇒ Is the intervention socioculturally acceptable to patients/beneficiaries as well as to those implementing it?
- ⇒ Is the intervention socioculturally acceptable to the public and other relevant stakeholder groups?
- ⇒ To what extent do patients/beneficiaries value different non-health outcomes?
- ⇒ How accessible—in terms of physical as well as informational access—is the intervention across different population groups?
- ⇒ What are the barriers and facilitators to implementing the intervention?

Adapted from Petticrew¹ and Rehfuess.⁸

paper reflects on frameworks for structuring systematic review questions, informed by a rapid review of existing frameworks, to evaluate their suitability when exploring complex interventions. The paper proposes an alternative framework, the PerSPECiTIF framework, for further testing.

It is increasingly recognised that systematic reviews of effects do not adequately capture how or why the effects of complex interventions differ according to context.^{4 5} Decision makers are demanding different types of synthesis to provide such evidence. Qualitative evidence synthesis (QES), for example, increasingly contributes to recommendations from WHO and other guideline development processes.^{6 7} QES can provide evidence for diverse questions beyond those that typically relate to the feasibility and acceptability of complex interventions (see Box 1)^{4 8 9} QES can potentially provide rich data relating to the context of interventions, policies or conditions and the lived experiences, views and beliefs of those involved. However, typical question frameworks for QES do not adequately account for a complexity perspective,^{10 11} in particular they do not account for the presence and assimilation of multiple stakeholder perspectives or

for the importance of contextual variation; critical if QES findings are to support holistic decision-making and if guidelines are to be applied with contextual sensitivity. As Squire and colleagues emphasise:

Such complexity ... makes the task of formulating a good review question both more important and more difficult. Furthermore, given the expected heterogeneity, systematic review questions should go beyond simple effectiveness questions (eg, 'does X work?') to consider under what circumstances X works.¹⁰

Guideline development organisations, such as WHO, the National Institute for Health and Care Excellence in the UK and other members of the Guidelines International Network (G-I-N), need to develop guideline recommendations that are feasible and acceptable to those planning, providing, implementing or receiving care. In turn, guideline development requires systematic review methodologies that explore the complexity of interventions, the context in which they are implemented and, the emphasis of this paper, the lens or evaluation frame through which they are evaluated.¹²

HOW DO GUIDELINE QUESTIONS CURRENTLY RECOGNISE COMPLEXITY?

Question formulation is critical to guideline development¹³ because it determines both the priorities to be addressed by the guideline and the types of evidence that will subsequently be admitted when addressing these priorities. Systematic reviews typically specify the elements of a question using an epidemiological design framework—population/patient; intervention; comparison; outcome(s); popularly known by the 'PICO' acronym¹⁴ or, replacing intervention with exposure, with the non-intervention formulation, PECO.¹⁵ Use of a PICO/PECO question within a guideline development process implicitly privileges an experimental/observational epidemiological model. Notwithstanding this inherent incompatibility the PICO model persists within many QES.^{16 17}

A systematic review team should routinely consider the potential added value of a complexity perspective for their review topic and work with guideline developers to decide whether a simple, complex intervention or complex system perspective is most appropriate and feasible for the review. The stage of question formulation thus offers a unique opportunity to surface and resolve issues that relate to such a complexity perspective.

As the example of a safety checklist illustrates, one cannot choose an appropriate perspective simply on the basis of intervention characteristics. Increasingly, health technology assessments, which have traditionally adopted an 'intervention perspective', are being encouraged to adopt a broader societal (and hence 'systems') perspective within the frameworks they use.^{18 19} Nevertheless, broadly speaking, questions relating to health technology assessment typically gravitate towards an 'intervention perspective' while those that arise within a public health context often adopt a 'systems perspective'. Decisions on

which overarching perspective to adopt must be based on what is most useful and most practicable. A review team can only attain partial understanding of complexity and they may implicitly or explicitly overlook something which subsequently matters.

This choice between an 'intervention perspective' and a 'systems perspective' holds clear implications for question formulation. Where an 'intervention perspective' is to be adopted, a QES typically explores barriers or facilitators to implementation, or the acceptability of the intervention to individuals and populations. A complex intervention perspective may still be accommodated within a population-based and intervention-based framework,¹⁴ or by use of frameworks that add either setting²⁰ or context.¹⁸ However, a 'complex systems perspective' for the same intervention makes the literature search and review process logistically demanding. Questions may be specified within an extended framework to specify the complexity of the decision problem, or, increasingly, review teams may articulate the problem within an a priori logic model,¹ which may be open to ongoing revision and refinement throughout the review.^{10 19}

FORMULATING COMPLEXITY-RELATED QUESTIONS TO BE ADDRESSED IN A QES

As described elsewhere within this series,^{1 8} guideline development groups consider issues that extend beyond effectiveness (what works) and cost-effectiveness (the cost-benefit of each option). Questions relate to intervention components, interactions between them, how intervention components adapt to and are modified by the context, how the system adapts to change and the role of feedback loops and emergent changes on the overall system.¹ Further questions align to the WHO-INTEGRATE framework.⁸ For example, concerns such as patient values in relation to health outcomes, safety, sociocultural acceptability, equity and equality and societal perspectives require different question frameworks and draw on diverse types of evidence. Both companion papers in this series^{1 2} rehearse why a complex systems perspective might be considered separately from a complex intervention perspective and why a systems perspective could add value to the intervention perspective. The resulting premise of this analysis is that review authors may consequently need to advance their complexity perspective beyond the typical PICO framework used in conventional reviews. Box 1 draws on earlier papers in this series to identify specific questions.^{4 8} The PICO framework does not easily accommodate a complexity perspective^{10 21} requiring an alternative framework for articulating the review question.

Previous commentators have identified the need to consider how a complex intervention is implemented (ie, feasibility) and the environment or context (ie, acceptability and meaningfulness) within which the intervention or programme is delivered. The GRADE Evidence to Decision (EtD) framework²² and the

Joanna Briggs 'FAME' framework²³ acknowledge that feasibility, appropriateness and meaningfulness must be recognised in the shift from 'what works' to 'what happens'.²¹

A guideline development group should prioritise questions relating to contextual variation that cannot be addressed in a primary study or questions which could be asked in a primary study, but are addressed robustly across multiple contexts through synthesis (Box 1). Questions may relate to *contextual variation* (variation across study contexts or between study contexts and the target context) or to *contextual sensitivity* (how the intervention must be adapted or modified to recognise contextual factors present in the target context). Questions addressed by a QES mirror those that can be answered by primary qualitative research. Clearly, questions that focus on differences in acceptability, delivery and implementation *between* interventions or programmes are better addressed through synthesis rather than by a single-context primary study. In contrast, a primary study is better equipped to identify nuances *within* a specific context. Simple PICO variants are unsuited to handle either the complexity inherent in any single complex intervention review question¹ or the wide variation in question types and evidence types required for guideline development (as illustrated by the WHO-INTEGRATE framework).⁸ The collective experience of the authors is that question formulation frameworks for qualitative evidence syntheses should include elements relating to both context and stakeholder perspective. Dissatisfaction with the suitability of PICO for constructing questions relating to feasibility or appropriateness has led to several alternatives being proposed. We therefore resolved to conduct a rapid review of existing question formulation frameworks to examine their suitability to accommodate a complexity perspective.

REVIEWING EXISTING QUESTION FORMULATION FRAMEWORKS

We performed a rapid review for currently available frameworks for formulating questions. This was not a comprehensive methodological review; we sought to map question variants and to examine their suitability for capturing a complexity perspective. In the interests of transparency, methods and results of the rapid review are reported separately in online supplementary file 1. However, the focus of this particular analysis is on specifying a candidate framework that accommodates complexity perspectives, not on the underpinning technical process by which individual items for possible inclusion were identified.

Following review of 1481 references and 113 full-text citations, we identified 38 question formulation frameworks (See online supplementary file 1 for full list of included references) which were evaluated for their suitability to accommodate a complexity perspective.

Table 1 Proposed question formulation framework for use of qualitative synthesis within guidelines

Per	S	P	E	(C)	Ti	F
Perspective	Setting	Phenomenon of interest/problem	Environment	(optional Comparison)	Time/timing	Findings

Based on the methodological literature relating to QES and complex interventions, the team considered that to be useful a question framework should recognise context (articulated as setting, environment or context)²⁴; acknowledge the criticality of stakeholder perspective (differentiated from target population)²⁵; accommodate elements of time/timing²⁶ and place; be sensitive to qualitative data (in the sense of eliciting themes or findings, rather than quantitative outcomes).⁹ None of the identified frameworks satisfied all four criteria. We therefore concluded that none of the existing frameworks was entirely suited to capture a complexity perspective when addressing questions potentially answerable by qualitative evidence syntheses.

Based on our rapid review of the elements of question formulation frameworks, we concluded that an optimal framework for use by guideline development groups would need to better incorporate elements of 'context', currently missing from existing frameworks. 'Context' encompasses temporal, spatial and societal dimensions¹ and offers an umbrella term beneath which environment and setting are subsumed. We propose that 'environment' should exploit well-established public health connotations of wider societal determinants and health service characteristics within which a service is delivered. In contrast, setting describes the point where interaction between service user and service provider takes place.⁶¹ Within these two overarching constructs, further granularity can be

accommodated; so, for example, a narrative review of contextual factors influencing health committees in low-income and middle-income countries identified four overlapping conceptual spheres—community and society which would be nested under environment and health facilities and health administration which would be articulated under setting.

An optimal framework would also capture different stakeholder perspectives, such as the views of partners or carers or, in a health systems context, of wider societal stakeholders. Finally, the question framework should encourage guideline developers to consider qualitative data, using alternatives to 'outcomes', and acknowledge contextual variation in time and space. Because no existing framework accommodates these requirements, the authors devised an alternative, PerSPeCtiF. Table 1 outlines the elements proposed in the PerSPeCtiF framework, which consolidates elements from existing question formulation frameworks.

Table 2 provides a worked example of a PerSPeCtiF question framework compared with the popular PICO and SPICE variants. Clear omissions of the latter two frameworks are flagged in the column 'missing'. So SPICE overlooks important features of the wider environment as well as temporal (time/timing) aspects while PICO omits not only these two elements but also the local context of setting.

Online supplementary table 1 illustrates the subsequent search strategy for this example. By prompting review authors to identify key areas of a complexity

Table 2 Worked example of question formulation framework (PerSPeCtiF, spice and PICO) for a qualitative synthesis

Per	S	P	E	(C)	Ti	F
From the perspective of a pregnant woman	In the setting of rural communities	How does the phenomenon of facility-based care	Within an environment of poor transport, infrastructure and geographically remote facilities	Compare with traditional birth attendants at home	In the time period up to and including childbirth	In relation to the woman's perceptions and experiences
	Setting	Perspective	Interest, phenomenon of	Comparison	Evaluation	Missing
In the setting of rural communities	From the perspective of a pregnant woman	How does the phenomenon of facility-based care	Compare with traditional birth attendants at home	In relation to the woman's perceptions and experiences?		Environment, time/timing
	Population	Intervention	Comparison	Outcome		Missing
	Pregnant women	Facility-based care	Traditional birth attendants at home	Women's perceptions and experiences		Setting environment time/timing

Box 2 Steps in the question formulation process for qualitative evidence syntheses (expanded from WHO¹³)

Step 1: generate an initial list of questions
 Step 2: decide whether a simple, complex intervention or complex system approach is most appropriate for this particular review
 Step 3: draft the key questions using a relevant question framework
 Step 4: list relevant stakeholder perspectives
 Step 5: review and revise
 Step 6: prioritise the key questions
 Step 7: order stakeholder perspectives according to their criticality to the implementation chain
 Step 8: finalise the key questions and the important and critical stakeholder perspectives

perspective beyond a typical PICO, the PerSPECtiF framework assists team members to identify appropriate search terms. Articulating the review question in this way helps to identify how eligible studies differ in time, setting, stakeholder perspectives and surrounding environment.

QUESTION FORMULATION IN GUIDELINE DEVELOPMENT

According to the WHO handbook for guideline development, question formulation requires an eight-step process (Box 2).¹³ Complex interventions generate numerous questions with implications for the scale and complexity of retrieval and synthesis. Below we use selected case studies to highlight where and how PerSPECtiF could be used.

Protocol development for a QES is iterative, not linear, starting with ‘guiding review questions’.²⁷ For the WHO smoking in pregnancy guidelines, a Technical Secretariat listed scoping questions and outcomes related to tobacco use and secondhand smoke exposure. This list was reviewed and prioritised by international stakeholders. Scoping establishes whether a review question is feasible (the logistics) and where conceptual boundaries (inclusion and exclusion criteria) lie.²⁸ Question formulation helps a guideline development group to consider what is feasible and what is meaningful. Will the QES review team mirror the scope of an accompanying effectiveness review? If so, the team may use similar subject terms for the searches for both effectiveness and qualitative data. Alternatively, will a system perspective examine the experience of a condition,²⁹ how the condition is viewed by health professionals or society at large and how implementation might occur³⁰? While QES can accommodate either a complex system or a complex intervention perspective, health technology assessments and guidelines often favour the narrower, intervention perspective, mainly for pragmatic reasons.³¹

Types of questions addressed by QES include: ‘Does evidence show that an “effective” intervention is poorly accepted within its target context?’ ‘Do studies demonstrate that an experimental intervention works suboptimally in a “real world” context?’. At this stage, all

questions are potentially valid. The guideline development group and stakeholders need to agree on terms used when articulating each question.¹⁰ Questions for the WHO smoking in pregnancy guidelines required several consultations with the guideline development group prior to identifying and synthesising the best available evidence for each key question within a systematic review.

Step 2 involves drafting the prioritised key questions using a relevant question formulation framework. Question formulation should ‘prompt review authors to identify the key components of the intervention/s and how these interact’.³² The PerSPECtiF question formulation framework encourages this complexity perspective³² by acknowledging contextual variation of both time and space, and by using alternatives to ‘outcomes’. In addition, by including ‘perspective’ as an element, the PerSPECtiF framework accommodates a ‘systems perspective’, with organisational levels and causal pathways¹⁰ (step 3), thus extending beyond the PICO format. Logic models offer an alternative way of thinking through such complexity¹ and may be revised throughout the review process, either opportunistically or at preplanned stages, as new data emerges.¹⁶

Stakeholders are key when refining questions and ‘lumping’ or ‘splitting’ interventions³³ and outcomes¹⁰ (steps 4–6). In prioritising questions to be addressed by qualitative evidence, guideline development groups need to agree which contextual factors are meaningful. This may require multiple iterations.⁷ Our rapid review revealed that current question frameworks fail to accommodate context. By incorporating ‘environment’ (the context within which a health service is delivered) and ‘setting’ (the point where interaction between service user and service provider takes place), PerSPECtiF seeks to address these limitations.

It is preferable to identify important contextual variables a priori, to inform study selection criteria. Alternatively, they may emerge when analysing included studies, determining how the data extraction form is structured. Ideally, a review team would construct a simple typology of important contextual factors. The reality is demonstrably different, for example, grouping African countries by socioeconomic factors or by predominant religion results in different groupings. By considering the extent of contextual variation early, a guideline development group not only shapes the final set of key questions but also informs data extraction and formulation of recommendations.

Having finalised the overarching guideline question (step 7), this is broken down into key questions to be addressed within the guideline. Key questions must be clearly formulated and the guideline development group should consider whether each key question is best addressed by quantitative or qualitative evidence or by a combination of both. The review team should decide whether these key questions are best addressed by evidence retrieved by an overarching guideline search, with evidence sifted and mapped to each key question,

or whether specific supplementary searches are required for each key question (step 8). Online supplementary table 2 includes what should be considered at each stage of the question formulation process.

CONSIDERATIONS WHEN FORMULATING REVIEW QUESTIONS FOR QES

In summary, systematic reviewers and guideline developers should decide whether a systematic review will adopt an 'intervention perspective' or extend a 'systems perspective' beyond the setting to include the wider environment. An intervention perspective can use population-based and intervention-based frameworks perhaps with the addition of context (eg, PICOC) or 'setting' (eg, SPICE). Where a 'systems perspective' offers insight a nuanced approach, using either the innovative PerSPECiTIF framework or a logic model structure is valuable.

PerSPECiTIF allows systematic reviewers and guideline developers to factor in contextual variation relating to time and/or space. Some variation can be anticipated; literature published prior to a particular date may be irrelevant to the decision in hand. Alternatively, the study contexts informing a particular guideline may emerge as substantively different from the target context and the review and guideline development teams may explicitly exclude such studies.³⁴ The first paper in this series offers guidance on how teams might judge contextual relevance.⁴

If a QES accompanies a systematic review of quantitative evidence, then teams must consider the extent to which the quantitative and qualitative review share the same question scope. As illustrated in the first paper of this series,¹ quantitative and qualitative approaches are typically complementary with questions addressed variously by quantitative or qualitative evidence or by a combination of both.

Finally, we acknowledge that the type of question asked is not only determined by the perspective adopted by the guideline but also, in turn, influences selection of an appropriate method for qualitative, or mixed method synthesis. Guidance on how to select a QES method based, for example, on whether a question is 'fixed or emerging' is available^{35 36} and is considered in other papers in this series.^{37 38}

We acknowledge that further work is required to test the utility of the PerSPECiTIF framework for the widest possible variety of questions addressed by QES. Similarly, there is a need to look at the extent to which this framework elicits additional information, not otherwise identified by population-based and intervention-based frameworks, and, importantly, what the implications of the framework are on the subsequent construction of search strategies.

CONCLUSION

This paper highlights the need for focused, relevant questions in qualitative evidence syntheses that address a complexity perspective. Use of the PerSPECiTIF framework, where a systems perspective is to be employed, should result in more appropriate questions addressing, for example, feasibility and

acceptability. Better formulation of questions will, in turn, lead to more informed decisions on choice of synthesis method and, ultimately, to better evidence on aspects of context that are important to consider when guidelines adopt a complexity perspective.

Author affiliations

¹School of Health and Related Research, University of Sheffield, Sheffield, UK

²School of Social Sciences, Bangor University, Wales, UK

³Department of Health Sciences, The University of York, York, UK

⁴School of Social Sciences, Cardiff University, Cardiff, UK

⁵Department of Reproductive Health and Research including UNDP/UNFPA/UNICEF/WHO/World Bank Special Programme of Research, Development and Research Training in Human Reproduction (HRP), World Health Organization, Genève, Switzerland

⁶Department of Health Education and Promotion, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

Acknowledgements The authors gratefully acknowledge input provided during the drafting process by Anayda Portela and Susan L Norris from the World Health Organization and by members of the other Complex Interventions Working Groups and by Helen Smith at the editorial stage.

Contributors AB, JN, KF and GM conceptualised the paper. ÖT and ES interpreted the paper in the context of guideline and decision-making processes. AB conducted the rapid review and devised the original PerSPECiTIF framework. JN, KF, GM and ÖT helped to revise the framework. AB wrote all drafts of the paper. All authors revised the drafts critically and approved the final version for publication.

Funding Funding provided by the World Health Organization Department of Maternal, Newborn, Child and Adolescent Health through grants received from the United States Agency for International Development and the Norwegian Agency for Development Cooperation.

Disclaimer ÖT is a staff member of WHO. The author alone is responsible for the views expressed in this publication and they do not necessarily represent the decisions or policies of WHO.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement All data supporting this article is available in the text, supporting tables within the text or the online supporting tables.

Open access This is an open access article distributed under the terms of the Creative Commons Attribution IGO License (CC BY NC 3.0 IGO), which permits use, distribution, and reproduction in any medium, provided the original work is properly cited. In any reproduction of this article there should not be any suggestion that WHO or this article endorse any specific organization or products. The use of the WHO logo is not permitted. This notice should be preserved along with the article's original URL.

Disclaimer: The author is a staff member of the World Health Organization. The author alone is responsible for the views expressed in this publication and they do not necessarily represent the views, decisions or policies of the World Health Organization.

REFERENCES

1. Wong G. Is complexity just too complex? *J Clin Epidemiol* 2013;66:1199–201.
2. Rogers PJ. Using programme theory to evaluate complicated and complex aspects of interventions. *Evaluation* 2008;14:29–48.
3. World Health Organization. *WHO surgical safety checklist and implementation manual*, 2012.
4. Petticrew M, Knai C, Thomas J, et al. Implications of a complexity perspective for systematic reviews and guideline development in health decision making. *BMJ Glob Health* 2019;4(Suppl 1):i6–i17.
5. Booth A, Moore G, Flemming K, et al. Taking account of context in systematic reviews and guidelines considering a complexity perspective. *BMJ Glob Health* 2019;4(Suppl 1):i6–i7.
6. Gülmezoglu AM, Chandler J, Shepperd S, et al. Reviews of qualitative evidence: a new milestone for Cochrane. *Cochrane Database Syst Rev* 2013;11:ED000073.

7. Langlois EV, Tunçalp Ö, Norris SL, *et al.* Qualitative evidence to improve guidelines and health decision-making. *Bull World Health Organ* 2018;96:79–79A.
8. Rehfuess EA, Stratil JM, Scheel IB, *et al.* The WHO-INTEGRATE evidence to decision framework version 1.0: integrating who norms and values and a complexity perspective. *BMJ Glob Health* 2019;4(Suppl 1):i90–i110.
9. Harris JL, Booth A, Cargo M, *et al.* Cochrane Qualitative and Implementation Methods Group guidance series-paper 2: methods for question formulation, searching, and protocol development for qualitative evidence synthesis. *J Clin Epidemiol* 2018;97:39–48.
10. Squires JE, Valentine JC, Grimshaw JM. Systematic reviews of complex interventions: framing the review question. *J Clin Epidemiol* 2013;66:1215–22.
11. Butler M, Epstein RA, Totten A, *et al.* AHRQ series on complex intervention systematic reviews-paper 3: adapting frameworks to develop protocols. *J Clin Epidemiol* 2017;90:19–27.
12. Kelly MP, Noyes J, Kane RL, *et al.* AHRQ series on complex intervention systematic reviews-paper 2: defining complexity, formulating scope, and questions. *J Clin Epidemiol* 2017;90:11–18.
13. World Health Organization. Chapter 7: formulating questions and selecting outcome. In: *Who Handbook for Guideline development*. Geneva: World Health Organization, 2014.
14. Richardson WS, Wilson MC, Nishikawa J, *et al.* The well-built clinical question: a key to evidence-based decisions. *ACP J Club* 1995;123:A12–13.
15. Collaboration for Environmental Evidence, 2013. Guidelines for systematic review and evidence synthesis in environmental Management: Environmental Evidence. Available from: <http://www.environmentalevidence.org/wp-content/uploads/2014/06/Review-guidelines-version-4.2-final.pdf>
16. Carstensen K, Lou S, Groth Jensen L, *et al.* Psychiatric service users' experiences of emergency departments: a CERQual review of qualitative studies. *Nord J Psychiatry* 2017;71:315–23.
17. Ryan C, Hesselgreaves H, Wu O, *et al.* Protocol for a systematic review and thematic synthesis of patient experiences of central venous access devices in anti-cancer treatment. *Syst Rev* 2018;7:61.
18. Petticrew M, Roberts H. *Systematic reviews in the social sciences: a practical guide*. Oxford: Blackwell Publishing, 2006.
19. Rehfuess EA, Booth A, Brereton L, *et al.* Towards a taxonomy of logic models in systematic reviews and health technology assessments: A priori, staged, and iterative approaches. *Res Synth Methods* 2018;9:13–24.
20. Booth A. Clear and present questions: formulating questions for evidence based practice. *Library Hi Tech* 2006;24:355–68.
21. Petticrew M. Time to rethink the systematic review catechism? Moving from 'what works' to 'what happens'. *Syst Rev* 2015;4:36.
22. Alonso-Coello P, Oxman AD, Moher J, *et al.* GRADE Evidence to Decision (EtD) frameworks: a systematic and transparent approach to making well informed healthcare choices. 2: clinical practice guidelines. *BMJ* 2016;353:i2089.
23. Pearson A, Wiechula R, Court A. *The JBI model of evidence-based healthcare. ... OF EVIDENCE BASED ...*, 2005.
24. Petticrew M, Rehfuess E, Noyes J, *et al.* Synthesizing evidence on complex interventions: how meta-analytical, qualitative, and mixed-method approaches can contribute. *J Clin Epidemiol* 2013;66:1230–43.
25. Thorne S, Jensen L, Kearney MH, *et al.* Qualitative metasynthesis: reflections on methodological orientation and ideological agenda. *Qual Health Res* 2004;14:1342–65.
26. Petticrew M, Anderson L, Elder R, *et al.* Complex interventions and their implications for systematic reviews: a pragmatic approach. *Int J Nurs Stud* 2015;52:1211–6.
27. Ako-Arrey DE, Brouwers MC, Lavis JN, *et al.* Health systems guidance appraisal-a critical interpretive synthesis. *Implement Sci* 2016;11:9.
28. Booth A, Sutton A, Papaioannou D. *Systematic approaches to a successful literature review*. Second edition. London: Sage, 2016.
29. Lorenc T, Pearson M, Jamal F, *et al.* The role of systematic reviews of qualitative evidence in evaluating interventions: a case study. *Res Synth Methods* 2012;3:1–10.
30. Movsisyan A, Melendez-Torres GJ, Montgomery P. A harmonized guidance is needed on how to "properly" frame review questions to make the best use of all available evidence in the assessment of effectiveness of complex interventions. *J Clin Epidemiol* 2016;77:139–41.
31. Booth A, Synthesis QE. Qualitative Evidence Synthesis. In: Facey K, Ploug Hansen H, Single A, eds. *Patient involvement in health technology assessment*. Singapore: Adis, 2017: 187–99.
32. Lewin S, Hendry M, Chandler J, *et al.* Assessing the complexity of interventions within systematic reviews: development, content and use of a new tool (iCAT_SR). *BMC Med Res Methodol* 2017;17:76.
33. Weir MC, Grimshaw JM, Mayhew A, *et al.* Decisions about lumping vs. splitting of the scope of systematic reviews of complex interventions are not well justified: a case study in systematic reviews of health care professional reminders. *J Clin Epidemiol* 2012;65:756–63.
34. Noyes J, Booth A, Lewin S, *et al.* Applying GRADE-CERQual to qualitative evidence synthesis findings-paper 6: how to assess relevance of the data. *Implement Sci* 2018;13:4.
35. Booth A, Noyes J, Flemming K. *Guidance on choosing qualitative evidence synthesis methods for use in health technology assessments of complex interventions*. INTEGRATE-HTA, 2016.
36. Booth A, Noyes J, Flemming K, *et al.* Structured methodology review identified seven (RETREAT) criteria for selecting qualitative evidence synthesis approaches. *J Clin Epidemiol* 2018;99:41–52.
37. Flemming K, Booth A, Garside R, *et al.* Qualitative evidence synthesis for complex interventions and guideline development: clarification of the purpose, designs and relevant methods. *BMJ Glob Health* 2019;4(Suppl 1):i40–i48.
38. Noyes J, Booth A, Moore G, *et al.* Synthesising quantitative and qualitative evidence to inform guidelines on complex interventions: Clarifying the purposes, designs and outlining some methods. *BMJ Glob Health* 2019;4(Suppl 1):i64–i77.