Methods Blood samples collected from consenting 133 males and 305 female participants in Ibadan were tested for HBsAg, HBeAg, HBcIgM, HBcTotal and HBsAb by ELISA technique. Samples positive for HBsAg were further analysed for HBV DNA by amplifying and sequencing the S gene. Isolates were genotyped and subtyped based on amino acid residues at position 122, 127, 134, 160 of the S gene.

Results Of the 438 subjects tested 31 (7.1%) were positive for HBsAg, 2 (6.5%) of which were HBeAg positive. Ninety-nine (22.8%) had detectable HBsAb, 3 (0.7%) were positive for HBcIgM and 195 (44.5%) were HBcTotal positive. HBV DNA was amplified and sequenced in 27 out of 31 and 4 could not be amplified due to low titres. After sequencing, 9 (33.3%) were not exploitable due to the presence of multiple peaks. Of the 18 exploitable isolates, only 15 showed significant similarity to HBV S-gene. Eleven of the 15 isolates were subtyped as ayw4 while others could not due to substitution at s122p. Phylogram showed that the 11 isolates were genotype E. Two of the 4 isolates with R122Q/P substitutions also belonged to genotype E while the other 2 which were >11% divergent from the reference genotype E sequence clustered with an isolate previously described as an Immune Escape Mutant.

Conclusion This study identified high endemicity of HBV infection, presence of markers of infection even in non-detectable HBsAg levels and circulation of genotype E ayw4 and vaccine mutants in south-western Nigeria. It therefore emphasises the risk of development of an indigenous infected population that may not be protected by the current vaccine.

PO 8276 COMMUNITY INDEX CASE APPROACH AND HIV TESTING AND COUNSELING (HTC) FOR SEXUAL PARTNERS OF HIV-POSITIVE PATIENTS LOST TO FOLLOW-UP: THE EXPERIENCE OF WORLD VISION MOZAMBIQUE

Miguelhete Lisboa*, Saraiva Culuze, Saimado Imputiua, Adolfo Cambule, Christine Latif. *World Vision Mozambique, Mozambique*

10.1136/bmjgh-2019-EDC.66

Background HIV-positive patients lost to follow-up (HP-LTFU) represent a challenge for HIV/AIDS control efforts as they are associated with higher risk of HIV transmission to their sexual partners, low viral load suppression and higher risk of morbidity and mortality than adherent patients. The SCIP-Ogumaniha programme implemented by World Vision Mozambique, has been utilising the index case approach together with systematic home-based HIV testing and counseling (hHTC) since August 2016 in 7 districts of the Zambezia province. This abstract outlines an evaluation of the contribution of this approach to HIV/AIDS care and treatment (HACT) of sexual partners of HP-LTFU in alignment with the first and second targets of the 90–90–90 UNAIDS strategy.

Methods The study involved HP-LTFU returned to HACT between October 2016 and September 2017. These patients reported to have sexual partners who had not been tested for HIV and provided informed, written consent for joint hHTC with these individuals. The hHTC package for sexual partners was offered by World Vision project counselors and those who tested HIV-positive were referred to HACT.

Results Of 7,084 patients who returned to HACT and reported to have an untested sexual partner, 63% (4,471)

provided informed, written consent for joint hHTC. Of 4264 sexual partners found and tested, 52% was female, 64% was in the 15–34 age groups, and 88% had never been tested for HIV. About 28% (1,205/4,264) was HIV-positive, 56% of the sexual partners who tested HIV-positive, was female and 98% of these was successfully referred to HACT.

Conclusion The index case approach together with hHTC has contributed to the early diagnosis of 28% of new HIV infections among sexual partners of HP-LTFU and 98% of them ensured timely linkage to the HACT. Therefore, broader promotion and adoption of this approach would make a significant contribution to achievement of the first and second targets of the 90–90–90 UNAIDS strategy.

PO 8278 BUILDING CAPACITY IN CONDUCTING CLINICAL RESEARCH IN A VIRTUAL SETTING: EXPERIENCES FROM THE EAST AFRICAN CONSORTIUM FOR CLINICAL RESEARCH (EACCR2)

¹Emily Nyanzi-Kabuye*, ¹Pontiano Kaleebu, ¹Benard Kikaire, ²Blandina T Mmbaga, ³Godfrey S Mfinanga, ⁴Norbert Peshu, ⁵Maowia Mukhtar. *¹Uganda Virus Research Institute, Entebbe, Uganda; ²Kilimanjaro Clinical Research Institute, Tanzania; ³National Institute of Medical Research - Muhimbili, Tanzania; ⁴Kenya Medical Research Institute- Wellcome Trust, Nairobi, Kenya; ⁵Institute of Endemic Diseases, University of Khartoum, Sudan*

10.1136/bmjgh-2019-EDC.67

Background EACCR2 is an EDCTP-funded, Eastern African-led network established in May 2009, with 23 regional partners from Ehtiopia, Kenya, Sudan, Tanzania, and Uganda, and 8 northern partners from Germany, Netherlands, Norway, Sweden and United Kingdom. The objective is to strengthen capacity to conduct health research to international standards with specific focus on clinical trials on poverty-relevant diseases such as HIV, TB, malaria and neglected infectious diseases. EACCR2 optimises the use of shared research infrastructures and other regional capacity building resources and opportunities.

Activities The activities of the network are implemented in five work packages cutting across ‘nodes’ in different countries. Capacity building programmes and studies funded by EDCTP are implemented by coordinators at the disease nodes. The Uganda Virus Research Institute (UVRI) hosts the secretariat of the consortium of five nodes located in the following institutions: Malaria Node in Kilifi-Kenya Medical Research Institute Wellcome Trust, Tanzania; Training Node in Kilimanjaro Clinical Research Institute, Tanzania; Tuberculosis Node at the National Institute of Medical Research- Muhimbili, Tanzania; the Neglected and Re-Emerging Tropical Diseases Node at the University of Khartoum, Sudan; and the HIV Node at UVRI, Uganda.

Coordinators form the project implementation committee which meets via skype or teleconference every quarter to assess progress, share best practice and challenges of the network. Scientific and annual meetings are arranged every year in one of the implementing institutions. During such meetings, students, the nodes and steering committee also meet to minimise travel costs while helping teams to network.

EACCR2 learns from the experiences, best practice and challenges of EACCR1 while implementing its current activities. Careful planning and consensus building from all partners has been the driving force to build and implement activities of this virtual network. EACCR2 also works closely with other EDCTP Networks of Excellence, i.e. in Central