Factors associated with head circumference and indices of cognitive development in early childhood

Laura Nicolaou,^{1,2} Tahmeed Ahmed,³ Zulfiqar Ahmed Bhutta ⁴, Pascal Bessong,⁵ Margaret Kosek,⁶ Aldo A M Lima,⁷ Sanjaya Shrestha,⁸ Ram Chandyo,⁹ Estomih R Mduma,¹⁰ Laura Murray-Kolb,¹¹ Brooks Morgan,^{1,2} Matthew R Grigsby,^{1,2} William Checkley,^{1,2} on behalf of the MALED Network Investigators

To cite: Nicolaou L, Ahmed T, Bhutta ZA, *et al.* Factors associated with head circumference and indices of cognitive development in early childhood. *BMJ Global Health* 2020;**5**:e003427. doi:10.1136/bmjgh-2020-003427

Handling editor Seye Abimbola

► Additional material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjgh-2020-003427>).

Received 16 July 2020

Revised 22 August 2020

Accepted 24 August 2020

© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to
Dr William Checkley;
wcheck1@jhmi.edu

ABSTRACT

Background While head circumference (HC) has been related to intracranial volume and brain size, its association with cognitive function remains unclear. We sought to understand the relationship among various biological and socioeconomic risk factors, HC and cognitive development.

Methods We analysed data across resource-poor settings in Bangladesh, India, Nepal, Peru, South Africa and Tanzania from the Etiology, Risk Factors and Interactions of Enteric Infections and Malnutrition and the Consequences for Child Health and Development longitudinal birth cohort study. Participating children were enrolled and followed up between 2009 and 2014. A final sample of 1210 children aged 0–24 months were included in the analyses. The main outcomes were HC for age Z-score and cognitive, gross motor and language scores from Bayley Scales of Infant Development-III tests. Length, weight and HC were measured monthly, and cognitive tests were administered at 6, 15 and 24 months of age. To disentangle the associations between risk factors and HC from linear growth and to distinguish the direct and indirect effects of these risk factors on cognitive function, we conducted mediation analysis using longitudinal models to account for all data measured during follow-up.

Results Average HC-for-age Z-score (HCAZ) was -0.54 (95% CI -0.47 to -0.62) near birth and -1.01 (95% CI -0.94 to -1.08) at 24 months. Children with higher enrolment weight ($p<0.0001$), higher socioeconomic score ($p=0.00037$) and taller mothers ($p=0.00084$) had higher HCAZ at all ages, while enteropathogen infection ($p=0.013$) and more febrile episodes ($p=0.013$) were associated with lower HCAZ. The associations between HCAZ and enrolment weight-for-age, maternal height, socioeconomic status or pathogen burden were partly mediated through their associations with length-for-age. HCAZ showed no association with cognitive, gross motor or language skills at 6, 15 and 24 months of age.

Conclusions The main risk factors associated with HC are similar to those associated with body length, and HC is not related to cognitive function.

Key questions

What is already known?

- Head circumference (HC) is a common anthropometric measurement that is used routinely in paediatric research and surveillance.
- It has been related to intracranial volume and brain size, but its association with cognitive function and the complex relationship between nutritional and environmental factors, body length, HC and cognitive outcomes remain unclear.

What are the new findings?

- Mediation analysis allowed us to disentangle associations between risk factors and HC from linear growth and to distinguish the direct and indirect effects of these risk factors on cognitive function.
- The main risk factors affecting HC were found to be similar to those affecting body length.
- We found no association between HC and cognitive function, gross motor function or language skills.

What do the new findings imply?

- Our results contribute to a growing body of literature that questions the value of HC as a marker of cognitive function in healthy children.
- These findings add to the limited literature on the relationship between HC and cognition in LMIC settings.

INTRODUCTION

Head circumference (HC) has been long considered a sensitive marker of neurodevelopment.^{1,2} Indeed, the American Academy of Pediatrics recommends measuring HC eight times during the first 2 years of life.³ In many settings in low-income and middle-income countries (LMICs) however, HC measurements are not performed regularly.⁴ Furthermore, there is some controversy about the utility of measuring HC for surveillance in healthy children beyond the neonatal period.

The United Kingdom Royal College of Paediatrics and Child Health for example currently recommends measuring HC at birth and 8 weeks.⁵

HC during early childhood development is associated with many nutritional factors, including protein energy, micronutrient intake and breastfeeding.^{6–8} Previous studies have examined the association between HC and intracranial volume, tying smaller heads to reduced brain volumes. However, it is unclear if the risk factors associated with HC growth impairment are independent from those associated with linear growth and if HC deficits are associated with cognitive impairments.^{9–11} Indeed, an analysis of 10 851 children found extreme HC measurements to be neither sensitive nor specific in identifying neurocognitive disorders.⁹ Another study of 2104 children from a birth cohort in Spain found no associations between HC or HC growth and mental or psychomotor scores at 14 months of age,¹² while an analysis of 505 full-term born children in South India found a positive association between HC at birth and subsequent cognitive abilities.¹³ While correlations have been established between nutritional and environmental factors, body length, HC and cognitive outcomes, the complex relationship between these facets remains unclear. Furthermore, there is a dearth of information on the association between HC and cognitive function in LMIC settings.

Since body length, HC and cognitive function are all inter-related,^{14–16} mediation analysis provides a useful framework to disentangle associations between risk factors and HC from linear growth, and to distinguish the direct and indirect effects of these risk factors on cognitive function¹⁷ (figure 1). We analysed data from a multi-centre cohort of children living in eight resource-poor

settings of LMICs followed from birth to 24 months. In a previous study,¹⁸ we determined the associations between risk factors and body length (Pathway 1). Factors that contributed to the odds of being in a lower body length-for-age category at 24 months were lower enrolment weight-for-age, shorter maternal height, higher number of enteropathogens in non-diarrheal stools, lower socioeconomic status and lower per cent of energy from protein. Site-specific analyses suggested that these associations were similar across settings.

Here, we examine the association between body length and HC (Pathway 2), the determinants of HC (Pathway 3) and the associations of cognitive function with HC (Pathway 4) and risk factors (Pathway 5). We also evaluate the indirect effects of body length on the associations between these risk factors and HC (Panel A) and those of HC on the associations with cognitive function (Panel B) to provide new insight into the complex relationship between all these facets. With these analyses, we seek to determine whether HC is a useful marker for cognitive function in healthy children and to establish whether any of our a priori determined risk factors affect HC growth differently from linear growth. Our hypothesis is that the same risk factors that predict body length also predict HC and that HC predicts cognition.

METHODS

Study design and participants

We analysed data from the Etiology, Risk Factors and Interactions of Enteric Infections and Malnutrition and the Consequences for Child Health and Development (MAL-ED) Network cohort study, a multidisciplinary,

Figure 1 Schematic illustrating the relationships between risk factors, body length, head circumference and cognitive function. Panels A and B are path diagrams of the mediation analyses depicting how the relationship between risk factors and HC is mediated by body length (A), and how the relationship between risk factors and cognitive function is mediated by HC (B). Path a represents the effect of the risk factors on the mediator (body length in A; HC in B). Path b represents the impact of the mediator on our dependent variable (HC in A; cognitive function in B). Together, paths a and b represent the indirect effect of the risk factors on the dependent variable through the mediator. Path c represents the direct effect of the risk factors on HC (A) or cognitive function (B), which is calculated controlling for the indirect effect. Path c represents the total (indirect plus direct) effect of the risk factors on HC (A) or cognitive function (B). HC, head circumference

prospective, longitudinal study aimed at better understanding the complex relationships among enteric infections, nutrition and other environmental exposures on child growth and development. Participating children were enrolled and followed up between 2009 and 2014 at eight field sites: Dhaka, Bangladesh (BGD); Fortaleza, Brazil (BRF); Vellore, India (INV); Bhaktapur, Nepal (NEB); Naushahro Feroze, Pakistan (PKN); Loreto, Peru (PEL); Venda, South Africa (SAV) and Haydom, Tanzania (TZH).¹⁹ After a baseline community census, enrolment began with the goal of recruiting 200 children per site. All infants were enrolled within 17 days of birth and followed continuously until 24 months.

Participant and public involvement

Neither participants nor the public were involved in the design, conduct or reporting of our study.

Anthropometric measurements and cognitive function

The main outcome for this analysis was HC for age Z-score (HCAZ) and cognitive, gross motor and language scores from the Bayley Scales of Infant Development-III (BSID-III) tests. We measured length, weight and HC in children monthly from the first 2 weeks of life. Several quality control measures were implemented, including the use of standardised techniques and instruments across all sites, same-day review of growth curves to identify unlikely measurements and duplicate anthropometric measurements on a subset of children.²⁰ We standardised all anthropometric measurements into Z-scores using the WHO growth standard.²¹

We assessed cognitive development through periodic administration of the BSID-III test for cognitive skills and gross motor function at 6, 15 and 24 months and language development at 15 and 24 months. At every site, the BSID-III test was translated, adapted to local culture and piloted on a separate group of children to ensure comparable difficulty of each item to the original item.²² Psychometric analyses were completed, and items with sufficient variability were kept (range 0–15). We excluded TZH from the cognitive development analyses, as quality control reviews revealed that the administration of cognitive tests deviated significantly from the protocol.²³ Cognitive scores from NEB at the 24-month time point were also excluded, as psychometric analyses suggested that the questions performed different compared with the other sites.²³

Biological and socioeconomic factors

Risk factors considered were weight-for-age Z-score at enrolment, total energy intake in kilocalories, cumulative proportion of days that a child was breastfed, micronutrient levels (inflammation-adjusted ferritin and retinol), altitude-adjusted haemoglobin concentration, maternal height, socioeconomic status (mean water and sanitation, wealth and assets, maternal education and household income (WAMI) score), food insecurity, diarrheal episodes, longitudinal prevalence of fever, acute lung

respiratory infection (ALRI), antibiotic use, faecal pathogen burden, alpha-1-acidglycoprotein (AGP) plasma concentration, myeloperoxidase (MPO), neopterin (NEO) and alpha-1 antitrypsin (AAT) concentrations, and lactulose:mannitol Z-score.

The methods of data collection for feeding, illness surveillance and microbiology are described elsewhere.^{24–29} Briefly, teams of field workers visited the households of participating children two times a week to attain daily reports of childhood illness (diarrhoea, acute lower respiratory infection and fever) and antibiotic use. Longitudinal prevalence was calculated as the number of days ill (or with antibiotic use) divided by the number of follow-up days.¹⁸ To assess micronutrient status, systemic inflammation, enteropathogen burden and gut inflammation and permeability markers, longitudinal visits were conducted to collect blood (at 7, 15 and 24 months), urine (at 3, 6, 9 and 15 months) and non-diarrheal stool samples (monthly during the first year of life and quarterly during the second year of life).¹⁹

From blood, we evaluated concentrations of altitude-adjusted haemoglobin, inflammation-adjusted retinol, AGP as a marker of systemic inflammation and inflammation-adjusted plasma ferritin by averaging the three measurements obtained during the study period. From non-diarrheal stools, we measured gut inflammation biomarkers, including MPO, NEO and AAT concentrations. We calculated quarterly averages from the monthly measurements collected during the first year and used the collected quarterly measurements thereafter.¹⁸ We computed enteropathogen load as the average number of enteropathogens in non-diarrheal stool divided by the total number of stool samples tested and determined cumulative enteropathogen load, assuming it was zero at enrolment and carrying forward cumulative values for months with missing data. From urine, we evaluated the lactulose:mannitol ratio as a marker of gut permeability and used the value measured at 3 months for the interval birth to 5 months, the 6-month value for the interval 6–8 months, the 9-month value for the interval 9–14 months and the 15-month value for the interval 15–24 months.^{18 27} We then transformed the lactulose:mannitol ratios to Z-scores using the BRF data as our internal reference standard. The BRF site had no deviation in anthropometry from the WHO growth standards and was therefore chosen to represent the least abnormal gut permeability values.^{28 29}

Initiation of breastfeeding was recorded and surveillance of feeding practices was conducted two times a week. We assumed days between visits to have the same status as the preceding visit³⁰ and calculated the cumulative proportion of days that a child was breastfed. Additionally, once a month beginning at 9 months, we conducted 24 hours complementary food intake recall interviews and computed cumulative energy intake from non-breast-milk foods over time, by linking the information provided in the interviews to country-specific nutritional databases. While a single recall does not provide a

highly precise estimate of usual energy intake at any given time point due to within-subject (ie, day-to-day) variation, the 16 measures available for each child from 9 to 24 months, can be combined to capture more precise intake information over time. To enhance the data collected, each child was randomly allocated to have a secondary recall done 2–7 days after one of the study visits between 9 and 24 months. This information was used to refine both group and individual estimates of energy intakes beyond the precision afforded by the monthly energy intake estimates.²⁴

Socioeconomic status (SES) was quantified using an index combining WAMI index.³¹ Every 6 months, we interviewed mothers or caregivers about their sources of water and sanitation facilities, assets, income and food security. We collected level of maternal education at baseline. As the WAMI index and food insecurity score showed small variation over time, we averaged their values across all collected time points.

Biostatistical methods

The analytical objectives were to study the association of HC growth with body length (Pathway 2, [figure 1](#)) and various biological and socioeconomic risk factors (Pathway 3); determine if the associations between these risk factors and HC were mediated by body length (Panel A); examine the association of cognitive function with HC (Pathway 4) and risk factors (Pathway 5) and last, determine if the associations between these risk factors and cognitive function were mediated by HC (Panel B).

To study the association of HC growth with body length (Pathway 2, [figure 1](#)) and various biological and socioeconomic factors (Pathway 3, [figure 1](#)), we used linear mixed-effects regression to model longitudinal HCAZ as a function of age, sex, indicator variables for sites and all of the above-mentioned biological and socioeconomic risk factors. All factors except site and sex were included as continuous variables. We modelled age using a piecewise linear spline with knots at age 3, 6 and 12 months and included interactions of the spline elements for age with all risk factors, site and sex. We included random effects at the individual level, accounting for heterogeneity in HC growth and HC at enrolment with random slopes and intercepts, respectively, and adjusted by site as a fixed effect. We built a fully adjusted model which included all the a priori risk factors as well as unadjusted models for each individual biological and socioeconomic risk factor, which included age, and interactions of the spline elements for age with that particular risk factor. We verified goodness-of-fit of the fully-adjusted model by comparing predicted and observed HCAZ trajectories with age (online supplemental efigure 1). Normality of residuals was confirmed through visual inspection (QQ plot of residuals, histogram of residuals, plot of residual versus fitted values).

To assess the contribution of linear growth on the relationship between risk factors and HC growth (Panel A), we estimated average direct effects (ADE), average

causal mediation effects (ACME) and average total effects (ATE) for each risk factor following the methods described by Imai *et al.*³² Specifically, we constructed a linear mixed-effects model for HCAZ as a function of above-mentioned variables plus length-for-age Z-score (mediator) and a linear mixed-effects model for length-for-age Z-score (LAZ) as a function of the same variables. We conducted 1000 Monte Carlo simulations to estimate the probability distributions of ADE, ACME and ATE and used a quasi-Bayesian approach to estimate mean and 95% CI.^{32 33} This method also allows for calculation of the percentage of the total association attributed to the mediator.

To evaluate Pathway 4, we modelled the indices of cognitive development at the ages of test administration (cognitive skills and gross motor function at 6, 15 and 24 months and language development at 15 and 24 months) as a function of HC and above-mentioned biological and socioeconomic factors. Since we have a single observation per participant at each time point, we adopted simple linear regression models. We used two types of HC measures: HCAZ and the slope of HCAZ (as a measure of HC growth) at the time of test administration. We estimated individual child slopes for HCAZ using a linear model of HCAZ as a function of age in the intervals between test administrations (0–6 months, 6–15 months and 15–24 months). To capture the non-linear relationships of the two HC measures with cognitive, gross motor and language scores (online supplemental efigure 2), we used truncated cubic regression splines. We also examined the relationship between cognitive function and time-lagged values of HCAZ using piecewise linear splines with knots at the 10th percentile, median and 90th percentile. The models included current (0 lag) and monthly lagged HCAZ in the interval between test administrations, that is, 0–6 months lagged HCAZ at 6 months of age and 0–8 months lagged HCAZ at 15 and 24 months of age.

Finally, using our linear models for the indices of cognitive development as a function of the above-mentioned factors plus HCAZ, and a linear model for HCAZ as a function of the same factors, we conducted mediation analyses as previously described. These analyses allowed us to evaluate the direct effects of these risk factors on cognitive function (Pathway 5) and to determine whether any of the three HC measures mediated the association between the risk factors and cognitive development (Panel B).

We conducted statistical analyses in R V.3.6.2 (Dark and Stormy Night) (<http://www.r-project.org>).

RESULTS

Study population

A total of 2145 children were included in the initial study population. Of those enrolled, 17.1% (n=367) were lost to follow-up and 9.8% (n=210) did not meet minimal criteria³ for longitudinal follow-up. Of those

Figure 2 Head circumference-for-age Z-score stratified by age and site. Site-specific head circumference-for-age trajectories with age were smoothed using natural splines. Sites include BGD, Dhaka, Bangladesh; INV, Vellore, India; NEB, Bhaktapur, Nepal; PEL, Loreto, Peru; SAV, Venda, South Africa; TZH, Haydom, Tanzania.

with complete data, participants from Pakistan ($n=277$) were excluded due to bias identified in a subset of length measurements, and participants from Brazil ($n=81$) were excluded due to bias in the HC measurements. A final sample of 1210 was included in the analyses. Those excluded from analysis varied from those included except in sex and mean WAMI score (online supplemental table 2). Among those who were included, all baseline features except for sex varied by site (online supplemental table 3).

Longitudinal trends in HC

On average, HCAZ fell with age during the first 9–12 months and remained constant thereafter, decreasing from -0.54 (95% CI -0.47 to -0.62) at enrolment to -1.01

(95% CI -0.94 to -1.08) at 24 months. Site-specific trajectories showed a decrease in HCAZ with age, except for PEL that showed an increase in the first 9 months of life followed by a plateau (figure 2). Mean HCAZs remained below the WHO standard (Z-score <0) across all sites except SAV. At enrolment, 12.6% ($n=153$) of children (range of 2.0% in SAV to 33.5% in INV) had $HCAZ < -2$, which rose to 20.8% ($n=252$) of children at 24 months (range of 1.5% in SAV to 52.5% in INV) (figure 3). Less than 1.5% of all children had $HCAZ \geq 2$.

Association between body length and HC (Pathway 2)

In online supplemental efigure 3, we show the 95% confidence ellipses for the scatter plots between HCAZ and LAZ by site at 3, 6, 12 and 24 months of age. These

Figure 3 Categories of head circumference-for-age stratified by age and country. Head circumference-for-age Z-score $HCAZ < -2$ is represented in orange, $-2 \leq HCAZ < -1$ is represented in yellow, $-1 \leq HCAZ < 2$ is represented in green and $HCAZ \geq 2$ in dark grey. Sites include BGD, Dhaka, Bangladesh; INV, Vellore, India; NEB, Bhaktapur, Nepal; PEL, Loreto, Peru; SAV, Venda, South Africa; TZH, Haydom, Tanzania.

confidence ellipses represent the 2D analogue of a CI for bivariate data, with their angle determined by the covariance of the data and the magnitude of the axes determined by the variance. As expected, there was a strong positive correlation between HCAZ and LAZ at all sites and ages. A larger variability in HCAZ compared with LAZ was observed across sites. On the other hand, LAZ displayed a larger decrease with age (mean and 95% CI across all sites: -1.04 (-0.98 to -1.10) at 3 months; -1.91 (-1.85 to -1.97) at 24 months) while HCAZ remained fairly constant beyond 6 months of age in all sites except BGD (mean and 95% CI across all sites: -0.81 (-0.74 to -0.88) at 3 months; -1.01 (-0.94 to -1.08) at 24 months).

Determinants of HC (Pathway 3)

Overall, a child had a higher HCAZ if he or she had a higher enrolment weight-for-age, was born to a taller mother or lived in a household with a higher WAMI. Pathogen burden and longitudinal prevalence of fever were negatively associated with HC (online supplemental efigure 4). The associations with WAMI index and pathogen burden were greater with age, while the associations with enrolment weight, maternal height and fever weakened with age. HCAZ was also weakly associated with higher energy intake, larger proportion of

days breastfeeding, lower prevalence of ALRIs and lower AAT and ferritin concentrations. No associations were found between HCAZ and sex ($p=0.49$; likelihood ratio test using a χ^2 distribution with 5 df (LRT with $\chi^2_{df=5}$), haemoglobin ($p=0.25$; LRT with $\chi^2_{df=5}$), retinol ($p=0.23$; LRT with $\chi^2_{df=5}$), food insecurity ($p=0.13$; LRT with $\chi^2_{df=5}$), NEO ($p=0.68$; LRT with $\chi^2_{df=5}$), MPO ($p=0.71$; LRT with $\chi^2_{df=5}$), lactulose:mannitol Z-score ($p=0.31$; LRT with $\chi^2_{df=5}$), AGP ($p=0.054$; LRT with $\chi^2_{df=5}$), prevalence of diarrheal episodes ($p=0.62$; LRT with $\chi^2_{df=5}$) or antibiotic use ($p=0.054$; LRT with $\chi^2_{df=5}$). There was substantial heterogeneity in the associations between risk factors and HCs across sites, with estimated mean differences in HCAZ between the lowest and higher quartiles of these risk factors varying in both magnitude and direction (online supplemental efigure 5).

Mediation of body length on associations between risk factors and HC (Panel A)

The effects of enrolment weight, WAMI index, maternal height and pathogen burden on HCAZ were all mediated by LAZ (figure 4). The indirect effect of body length on the association between maternal height and HCAZ

Figure 4 Mediation of body length on head circumference growth during early life development. The symbols represent the direct (ADE), length-mediated (ACME) and total effects on head circumference scaled to the interdecile range of various risk factors at enrolment, 12 months and 24 months of age. For example, the total effect of enrolment weight Z-score (WEIZ) on HCAZ at 1 month of age is 0.456. This value reflects how much one unit change in WEIZ affects HCAZ. To determine how much a difference between the 90% and 10% percentile of WEIZ affects HCAZ, we scale this per-unit effect by the IDR of WEIZ. The 10% and 90% percentiles of WEIZ at 1 month of age are -2.19 and 0.52 , respectively; the IDR is 2.71 . The total effect over the interdecile range is therefore 1.24 (0.456×2.71) as plotted in the figure. Error bars represent 95% CIs. ACME, average causal mediation effect; ADE, average direct effect; IDR, interdecile range; WAMI, water and sanitation, wealth and assets, maternal education and household income.

remained fairly constant throughout the first 2 years of life with mediation effects constituting 35%–40% of the total effects. On the other hand, the contribution of mediation effects to the total effects of enrolment weight, mean WAMI index or pathogen score on HCAZ decreased with age. Body length accounted for 25% of the total effect of enrolment weight on HCAZ in the first month of age, dropping to 20% at 12 months and 16% at 24 months. The mediation effect of body length contributed to 19%, 16% and 15% of the total effect of WAMI index on HCAZ at 1, 12 and 24 months, respectively, and 28% and 21% of the total effect of pathogen burden at 12 and 24 months, respectively. No mediation was found on the effects of fever.

Associations between HC and cognitive function (Pathway 4)

We found no associations between cognitive, gross motor or language scores and any of the three HC measures as assessed at 6, 15 and 24 months of age (online supplemental efigure 6). Furthermore, taking into account various time lags between cognitive function and HCAZ, the vast majority of differences in all three cognitive measures were not significant, and those that were statistically significant, such as the differences in cognitive score at 6 months of age with 0 and 1 month lagged HCAZ, appeared in opposite directions showing no clear association between cognitive function and HCAZ (online supplemental efigures 7 and 8).

Risk factors for cognitive impairment (Pathway 5) and mediation by HC (Panel B)

Cognitive function was positively associated with enrolment weight, maternal height and WAMI (figure 5A). The total

effects (ATEs) of WAMI on cognitive score increased with age, while those of enrolment weight and maternal height were only significant at 6 months. We also found negative associations with MPO and AGP concentrations at 15 months of age (online supplemental efigure 9). However, none of these associations were mediated by HC. Higher gross motor function scores were associated with higher enrolment weight, WAMI index and haemoglobin levels (figure 5B). None of these associations were mediated by HC. Associations between gross motor function and all other risk factors were found to be insignificant (online supplemental efigure 10). For language, we found a positive association with WAMI and haemoglobin, and a negative association with pathogen burden (figure 5C). A small positive effect was also observed for lactulose:mannitol at 15 months (online supplemental efigure 11). The causal mediation effect of HC on these associations remained non-significant. Additional mediation analyses conducted using the HCAZ slope and HCAZ/LAZ as HC measures revealed similar results (not presented).

DISCUSSION

In this analysis, we found that the previously identified biological and socioeconomic factors associated with linear growth¹⁸ are also associated with HC growth. Moreover, the effects of enrolment weight-for-age, maternal height, WAMI index and pathogen burden on HC were partly mediated through body length. HC was not associated with indicators of cognitive development at any age, whether it was language, cognitive skills or gross motor function; and, HC did not mediate the association between any of the studied risk factors and cognitive

Figure 5 Mediation of head circumference on (A) cognitive function, (B) gross motor function and (C) language skills during early life development. The symbols represent the direct (ADE), head circumference mediated (ACME) and total effects on cognitive, gross motor and language scores over the interdecile range of various risk factors at 6, 15 and 24 months of age. Error bars represent 95% CIs. ACME, average causal mediation effect; ADE, average direct effect.

development. These findings suggest that the routine measurement of HC may not provide useful information regarding cognitive development in healthy children. Our study also adds to the limited literature on the relationship between HC and cognition in LMIC settings.

Our findings are consistent with those observed by other investigators in different settings.^{9–12} Indeed, the Avon Longitudinal Study of Parents and Children (ALSPAC) study found that most children with small heads will have no underlying pathology and that the most common explanation for centile shifts in HC is measurement error.⁹ ALSPAC investigators concluded that longitudinal HC assessments were unhelpful (unless <2 SDs or >2 SDs below or above the median) and that HC surveillance should be conducted at 2–5 days after birth and then at 6 months of age. Treit *et al* studied 144 children with prenatal alcohol exposure and 145 healthy children without prenatal alcohol exposure but did not find significant associations between HC and any cognitive score and called into question the predictive value of HC at the individual-subject level.¹⁰ Our findings provide further support to the notion that longitudinal assessment of HC is neither helpful nor cost-effective for research or surveillance in healthy children.^{9 12 34} One possible exception could be screening for hydrocephalus. A population-based study conducted in Norway found that increased HC as a debut symptom was only important in children with hydrocephalus and cysts, but only during the first 10 months of life.³⁵ The authors concluded that the benefits of extending routine measurements of HC even for hydrocephalus beyond 1 year of age is questionable. Another study conducted in Netherlands found the same results.³⁶

Overall, we observed similar trends of a fall in HC with age and similar relationships between risk factors and HC as in a previous analysis conducted by our group where linear growth was the main outcome.¹⁸ HC decreased on average between birth and the first years of life at roughly the same magnitude as it did for body length. The risk factors for a lower HC gain were also similar to those identified for linear growth, namely, a lower enrolment weight, lower maternal height, lower socioeconomic status and a higher prevalence of enteric pathogens. Finally, the direction and magnitude of association of these risk factors with age on HC were similar to those documented for body length. Mediation analyses revealed that the effects between risk factors and a lower HC were explained in part by their contributions to a lower body length. Our findings are also consistent with a single-country analysis of data collected in India,⁵ where our colleagues found that HC was negatively associated with stunting at all ages and that a lower socioeconomic status was associated with a lower HC. Our analyses and findings differ from those presented in a prior study³⁷ which focused on the association between HC and cognitive function adjusted only for enrolment weight and site and point towards a potential residual confounding in the results presented by Scharf *et al*.³⁷ One concern about our analysis is the potential overfitting given the

large number of model parameters; however, our results were unchanged whether we adjusted for the five most important risk factors or for the entire set of risk factors.

An important secondary finding of this analysis was the relationship between a lower socioeconomic status and worse cognitive development in our study children. This finding is not entirely surprising as the association between childhood socioeconomic status and late-life cognition in the US, Europe and China is well documented.^{38–41} While the mechanisms are not well understood, our findings demonstrate that factors which predict linear growth and HC do not similarly predict cognition, which suggests that socioeconomic adversity affects brain development through its effects on diet and other non-nutritional factors.⁴¹ There may also be other nutrient deficiencies not considered in this study that could be related to cognition even though factors that predict growth and head size might not similarly predict cognition. Mechanisms notwithstanding, this finding suggests that one of the most important interventions that can lead to improvements in child growth and development is poverty reduction. Moreover, our analysis adds evidence on the importance of life-course socioeconomic status on cognitive development disparities.

Our analysis has several strengths. First, the MAL-ED study collected a comprehensive assortment of prenatal and postnatal factors across a wide array of settings worldwide, each of which used a standardised protocol and data collection toolset. As this was a prospective study, repeat visits allowed for accurate collection of pathogen, nutrition and illness surveillance data over time. Second, our study conducted high-quality anthropometric assessment which allowed us to examine monthly changes in body length and HC early in childhood and relate it to several biological and socioeconomic risk factors. Third, our analyses used rigorous statistical methods to help disentangle the relationships between risk factors, body length, HC and cognition. These analyses were conducted on random samples of infants in resource-limited settings across six different LMICs. Future studies in other countries should determine whether our findings are generalisable to other settings.

This analysis also has some potential shortcomings. First, the relationships between genetic, nutritional, socioeconomic and environmental factors are complex and our results may include unaccounted for confounding or feedback effects. Second, although the MALED study contributed data from eight sites, all data from Pakistan had to be dropped due to bias in length measurements, all data from Brazil had to be dropped due to bias in HC measurements and all cognitive data from Tanzania had to be dropped due to bias in the cognitive development assessment. Third, our study failed to collect information on gestational age. While our eligibility criteria excluded those with a birth weight <1.5 kg, this criterion may not adequately screen for preterm birth or severe intra-uterine growth restriction. We should also note that our analysis did not account for the role of caregiving and

early-life stimulation and play in cognitive development. Other important limitations that need to be considered are the loss of higher-risk infants from study follow-up; limited predictive value of cognitive testing at 24 months for school-age children; lack of validation for the Bayley scales in our study settings and the unknown predictive value of the newly created subscale. However, adaptation, piloting and validity assessments of the BSID-III tests were conducted at each of the study sites.²²

CONCLUSION

In conclusion, using data from one of the most comprehensive longitudinal studies of child growth and development in LMICs, and accounting for the most extensive array of confounders to date, we found that the main risk factors associated with HC were similar to those associated with body length. We did not find an association between HC and cognitive function and did not find an indirect effect of HC on the association between cognitive function and other risk factors.

Author affiliations

¹Division of Pulmonary and Critical Care, School of Medicine, Johns Hopkins University, Baltimore, Maryland, USA

²Center for Global Non-Communicable Diseases Research and Training, Johns Hopkins University, Baltimore, United States

³Division of Nutrition and Food Security, International Centre for Diarrhoeal Disease Research, Matlab, Bangladesh

⁴Centre for Global Child Health, The Hospital for Sick Children, Toronto, Ontario, Canada

⁵HIV/AIDS and Global Health Research Programme, University of Venda, Thohoyandou, Limpopo, South Africa

⁶Division of Infectious Diseases and International Health, University of Virginia, Charlottesville, Virginia, USA

⁷Clinical Research Unit and Institute of Biomedicine, Faculty of Medicine, Univ Fed Ceara, Fortaleza, Ceará, Brazil

⁸Walter Reed Armed Forces Research Institute of Medical Sciences (AFRIMS) Research Unit, Kathmandu, Nepal

⁹Department of Community Medicine, Kathmandu Medical College, Sinamangal, Kathmandu, Nepal

¹⁰Haydom Lutheran Hospital, Haydom, United Republic of Tanzania

¹¹Department of Nutritional Sciences, College of Health and Human Development, The Pennsylvania State University, University Park, Pennsylvania, USA

Correction notice This article has been corrected since it published online to fix the figures size.

Acknowledgements We particularly want to thank the children and caregivers who participated in the study for their invaluable contributions. Tom Brewer and Gretchen Meller are also thanked for their early support of the project. We are truly indebted to the following individuals for their advisory role in the MAL-ED study: Henry Binder, Maureen Black, Kathleen Braden, Robert Breiman, Susan Bull, Katherine Dewey, Christopher Duggan, Christine Grady, Gerry Keusch, Nancy Krebs, Claudio Lanata, James Nataro, Jerome Singh, Peter Smith, Phillip Tarr, Katherine Tucker and Theodore Wachs. Members of the MAL-ED Network Investigators are as follows: Angel Mendez Acosta, Rosa Rios de Burga, Cesar Banda Chavez, Julian Torres Flores, Maribel Paredes Olotegui, Silvia Rengifo Pinedo, Mery Sigua Salas, Dixer Rengifo Trigoso, Angel Orbe Vasquez (AB PRISMA, Iquitos, Peru); Imran Ahmed, Didar Alam, Asad Ali, Zulfiqar A Bhutta, Shahida Qureshi, Muneera Rasheed, Sajid Soofi, Ali Turab, Anita KM Zaidi (Aga Khan University, Karachi, Pakistan); Ladaporn Bodhidatta, Carl J Mason (Armed Forces Research Institute of Medical Sciences, Bangkok, Thailand); Sudhir Babji, Anuradha Bose, Ajila T George, Dinesh Hariraju, M Steffi Jennifer, Sushil John, Shiny Kaki, Gagandeep Kang, Priyadarshani Karunakaran, Beena Koshy, Robin P Lazarus, Jayaprakash Muliylil, Mohan Venkata Raghava, Sophy Raju, Anup Ramchandran, Rakhi Ramadas, Karthikeyan Ramanujam, Anuradha Bose, Reeba Roshan, Surjan L Sharma, Shanmuga

Sundaram E, Rahul J Thomas (Christian Medical College, Vellore, India); William K Pan (Duke University, Durham, North Carolina, USA); Fogarty International Center/National Institutes of Health, Bethesda, Maryland, USA); Ramya Ambikapathi, J Daniel Carreon, Vivek Charu, Viyada Doan, Jhanelle Graham, Christel Hoest, Stacey Knobler (Fogarty International Center/National Institutes of Health, Bethesda, Maryland, USA); Dennis R Lang (Fogarty International Center/National Institutes of Health, Bethesda, Maryland, USA); Foundation for the NIH, Bethesda, Maryland, USA); Benjamin JJ McCormick, Monica McGrath, Mark A Miller, Archana Mohale, Gaurvika Nayyar, Stephanie Psaki, Zeba Rasmussen, Stephanie A Richard, Jessica C Seidman, Vivian Wang (Fogarty International Center/National Institutes of Health, Bethesda, Maryland, USA); Rebecca Blank, Michael Gottlieb, Karen H Tountas (Foundation for the NIH, Bethesda, Maryland, USA); Caroline Amour, Eliwaza Bayyo, Estomih R Mduma, Regisiana Mvungi, Rosemary Nshama, John Pascal, Buliga Mujaga Swema, Ladislaus Yarrot (Haydom Lutheran Hospital, Haydom, Tanzania); Tahmeed Ahmed, AM Shamsir Ahmed, Rashidul Haque, Iqbal Hossain, Munirul Islam, Mustafa Mahfuz, Dinesh Mondal, Fahmida Tofail (icddr, Dhaka, Bangladesh); Ram Krishna Chandyo, Prakash Sunder Shrestha, Rita Shrestha, Manjeswori Ulak (Institute of Medicine, Tribhuvan University, Kathmandu, Nepal); Aubrey Bauck, Robert E Black, Laura E Caulfield (Johns Hopkins University, Baltimore, Maryland, USA); William Checkley (Johns Hopkins University, Baltimore, Maryland, USA); Fogarty International Center/National Institutes of Health, Bethesda, Maryland, USA); Margaret N Kosek, Gwenyth Lee, Kerry Schulze, Pablo Peñataro Yori (Johns Hopkins University, Baltimore, Maryland, USA); Laura E Murray-Kolb (The Pennsylvania State University, University Park, Pennsylvania, USA); A Catharine Ross (The Pennsylvania State University, University Park, Pennsylvania, USA); Barbara Schaefer (The Pennsylvania State University, University Park, Pennsylvania, USA); Fogarty International Center/National Institutes of Health, Bethesda, Maryland, USA); Suzanne Simons (The Pennsylvania State University, University Park, Pennsylvania, USA); Laura Pendergast (Temple University, Philadelphia, Pennsylvania, USA); Cláudia B Abreu, Hilda Costa, Alessandra Di Moura (Universidade Federal do Ceara, Fortaleza, Brazil); José Quirino Filho (Universidade Federal do Ceara, Fortaleza, Brazil); Fogarty International Center/National Institutes of Health, Bethesda, Maryland, USA); Alexandre Havt, Álvaro M Leite, Aldo AM Lima, Noélia L Lima, Ila F Lima, Bruna LL Maciel, Pedro HQS Medeiros, Milena Moraes, Francisco S Mota, Reinaldo B Oriá (Universidade Federal do Ceara, Fortaleza, Brazil); Josiane Quetz, Alberto M Soares, Rosa MS Mota (Universidade Federal do Ceara, Fortaleza, Brazil); Crystal L Patil (University of Illinois, Chicago, Illinois, USA); Pascal Bessong, Cloupas Mahopo, Angelina Maphula, Emanuel Nyathi, Amidou Samie (University of Venda, Thohoyandou, South Africa); Leah Barrett, Rebecca Dillingham, Jean Gratz, Richard L Guerrant, Eric Houpt, William A Petri, Jr, James Platts-Mills, Rebecca Scharf (University of Virginia, Charlottesville, Virginia, USA); Binob Shrestha, Sanjaya Kumar Shrestha (Walter Reed/AFRIMS Research Unit, Kathmandu, Nepal); Tor Strand (Walter Reed/AFRIMS Research Unit, Kathmandu, Nepal; University of Bergen, Norway); Erling Svensen (Haukeland University Hospital, Bergen, Norway; Haydom Lutheran Hospital, Haydom, Tanzania).

Contributors LN and WC conducted statistical analyses and wrote the first draft of the manuscript. BM and MRG contributed to the literature search and data analysis. TA, ZAB, PB, MK, AAML, SS, RC, ERM and LM-K contributed to data interpretation and reviewed and contributed to the manuscript content.

Funding The Etiology, Risk Factors and Interactions of Enteric Infections and Malnutrition and the Consequences for Child Health and Development Project (MAL-ED) is carried out as a collaborative project supported by the Bill & Melinda Gates Foundation (#OPP47075), the Foundation for the NIH and the National Institutes of Health/Fogarty International Center.

Disclaimer The Bill & Melinda Gates Foundation did not play any role in the writing of the manuscript nor did the funders of the study have any role in the study design, data collection, data analysis or interpretation of study results. The corresponding author had full access to all the data in the study and had final responsibility for the decision to submit for publication.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval Ethics approval was obtained from the Institutional Review Boards at participating institutions (online supplemental table 1).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available in a public, open access repository. Data for this study are available to others in the scientific community at https://clinepidb.org/ce/app/record/dataset/DS_5c41b87221.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iD

Zulfiqar Ahmed Bhutta <http://orcid.org/0000-0003-0637-599X>

REFERENCES

- Hack M, Breslau N, Fanaroff AA. Differential effects of intrauterine and postnatal brain growth failure in infants of very low birth weight. *Am J Dis Child* 1989;143:63–8.
- Lipper E, Lee K, Gartner LM, et al. Determinants of neurobehavioral outcome in low-birth-weights infants. *Pediatrics* 1981;67:502–5.
- American Academy of Pediatrics. *Bright futures guidelines for health supervision of infants, children, and adolescents*. 889. 4th ed, 2017. <https://ebooks.aappublications.org/content/9781610020237/9781610020237>
- Sindhu KN, Ramamurthy P, Ramanujam K, et al. Low head circumference during early childhood and its predictors in a semi-urban settlement of Vellore, southern India. *BMC Pediatr* 2019;19:182.
- RCPCH. *Health for all children*. 5th ed. Oxford University Press, 2019.
- Georgieff MK. Nutrition and the developing brain: nutrient priorities and measurement. *Am J Clin Nutr* 2007;85:614S–20.
- Miller LC, Joshi N, Lohani M, et al. Head growth of undernourished children in rural Nepal: association with demographics, health and diet. *Paediatr Int Child Health* 2016;36:91–101.
- Gewa CA, Weiss RE, Bwibo NO, et al. Dietary micronutrients are associated with higher cognitive function gains among primary school children in rural Kenya. *Br J Nutr* 2009;101:1378–87.
- Wright CM, Emond A. Head growth and neurocognitive outcomes. *Pediatrics* 2015;135:e1393–8.
- Treit S, Zhou D, Chudley AE, et al. Relationships between head circumference, brain volume and cognition in children with prenatal alcohol exposure. *PLoS One* 2016;11:e0150370.
- Peters M. Still no convincing evidence of a relation between brain size and intelligence in humans. *Canadian Journal of Experimental Psychology/Revue canadienne de psychologie expérimentale* 1993;47:751–6.
- Álamo-Junquera D, Sunyer J, Iñiguez C, et al. Prenatal head growth and child neuropsychological development at age 14 months. *Am J Obstet Gynecol* 2015;212:661.e1–661.e11.
- Veena SR, Krishnaveni GV, Wills AK, et al. Association of birthweight and head circumference at birth to cognitive performance in 9- to 10-year-old children in South India: prospective birth cohort study. *Pediatr Res* 2010;67:424–9.
- Geraedts EJ, van Dommelen P, Caliebe J, et al. Association between head circumference and body size. *Horm Res Paediatr* 2011;75:213–9.
- Wickett JC, Vernon PA, Lee DH. Relationships between factors of intelligence and brain volume. *Pers Individ Dif* 2000;29:1095–122.
- Cooke RW, Lucas A, Yudkin PL, et al. Head circumference as an index of brain weight in the fetus and newborn. *Early Hum Dev* 1977;1:145–9.
- Xie W, Jensen SKG, Wade M, et al. Growth faltering is associated with altered brain functional connectivity and cognitive outcomes in urban Bangladeshi children exposed to early adversity. *BMC Med* 2019;17:199.
- MAL-ED Network Investigators. Childhood stunting in relation to the pre- and postnatal environment during the first 2 years of life: the MAL-ED longitudinal birth cohort study. *PLoS Med* 2017;14:e1002408.
- Acosta AM, Chavez CB, Flores JT, et al. The MAL-ED study: a multinational and multidisciplinary approach to understand the relationship between enteric pathogens, malnutrition, gut physiology, physical growth, cognitive development, and immune responses in infants and children up to 2 years of age in resource-poor environments. *Clin Infect Dis* 2014;59:S193–206.
- Richard SA, McCormick BJJ, Miller MA, et al. Modeling environmental influences on child growth in the MAL-ED cohort study: opportunities and challenges. *Clin Infect Dis* 2014;59:S255–60.
- de Onis M. Measuring nutritional status in relation to mortality. *Bull World Health Organ* 2000;78:1271–4.
- Murray-Kolb LE, Rasmussen ZA, Scharf RJ, et al. The MAL-ED cohort study: methods and lessons learned when assessing early child development and caregiving mediators in infants and young children in 8 low- and middle-income countries. *Clin Infect Dis* 2014;59:S261–72.
- MAL-ED Network Investigators. Early childhood cognitive development is affected by interactions among illness, diet, enteropathogens and the home environment: findings from the MAL-ED birth cohort study. *BMJ Glob Health* 2018;3:e000752.
- Caulfield LE, Bose A, Chandyo RK, et al. Infant feeding practices, dietary adequacy, and micronutrient status measures in the MAL-ED study. *Clin Infect Dis* 2014;59:S248–54.
- Haupt E, Gratz J, Kosek M, et al. Microbiologic methods utilized in the MAL-ED cohort study. *Clin Infect Dis* 2014;59:S225–32.
- Richard SA, Barrett LJ, Guerrant RL, et al. Disease surveillance methods used in the 8-site MAL-ED cohort study. *Clin Infect Dis* 2014;59:S220–4.
- Kosek M, Guerrant RL, Kang G, et al. Assessment of environmental enteropathy in the MAL-ED cohort study: theoretical and analytic framework. *Clin Infect Dis* 2014;59:S239–47.
- Kosek MN, Lee GO, Guerrant RL, et al. Age and sex normalization of intestinal permeability measures for the improved assessment of enteropathy in infancy and early childhood: results from the MAL-ED study. *J Pediatr Gastroenterol Nutr* 2017;65:31–9.
- Kosek MN, Ahmed T, Bhutta Z, MAL-ED Network Investigators. Causal pathways from enteropathogens to environmental enteropathy: findings from the MAL-ED birth cohort study. *EBioMedicine* 2017;18:109–17.
- Ambikapathi R, Kosek MN, Lee GO, et al. How multiple episodes of exclusive breastfeeding impact estimates of exclusive breastfeeding duration: report from the eight-site MAL-ED birth cohort study. *Matern Child Nutr* 2016;12:740–56.
- Psaki SR, Seidman JC, Miller M, et al. Measuring socioeconomic status in multicountry studies: results from the eight-country MAL-ED study. *Popul Health Metr* 2014;12:8.
- Imai K, Keele L, Tingley D. A general approach to causal mediation analysis. *Psychol Methods* 2010;15:309–34.
- King G, Tomz M, Wittenberg J. Making the most of statistical analyses: improving interpretation and presentation. *Am J Pol Sci* 2000;44:347–55.
- Coon ER, Young PC, Quinonez RA, et al. Update on pediatric overuse. *Pediatrics* 2017;139:e20162797.
- Zahl SM, Wester K. Routine measurement of head circumference as a tool for detecting intracranial expansion in infants: what is the gain? A nationwide survey. *Pediatrics* 2008;121:e416–20.
- Breuning-Broers JM, Deurloo JA, Gooskens RH, et al. At what age is hydrocephalus detected, and what is the role of head circumference measurements? *Eur J Public Health* 2014;24:32–4.
- Scharf RJ, Rogawski ET, Murray-Kolb LE, et al. Early childhood growth and cognitive outcomes: findings from the MAL-ED study. *Matern Child Nutr* 2018;14:e12584.
- Zhang Z, Gu D, Hayward MD. Childhood nutritional deprivation and cognitive impairment among older Chinese people. *Soc Sci Med* 2010;71:941–9.
- Melrose RJ, Brewster P, Marquie MJ, et al. Early life development in a multiethnic sample and the relation to late life cognition. *J Gerontol B Psychol Sci Soc Sci* 2015;70:519–31.
- Singh-Manoux A, Richards M, Marmot M. Socioeconomic position across the lifecourse: how does it relate to cognitive function in mid-life? *Ann Epidemiol* 2005;15:572–8.
- Ranjitkar S, Hysing M, Kvestad I, et al. Determinants of cognitive development in the early life of children in Bhaktapur, Nepal. *Front Psychol* 2019;10:2739.