Mass media campaigns to reduce unnecessary caesarean sections: a systematic review

Maria Regina Torloni ¹, Vanessa Brizuela ², Ana Pilar Betran ²

To cite: Torloni MR, Brizuela V, Betran AP. Mass media campaigns to reduce unnecessary caesarean sections: a systematic review. *BMJ Global Health* 2020;**5**:e001935. doi:10.1136/bmjgh-2019-001935

Handling editor Seye Abimbola

► Additional material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjgh-2019-001935>).

Received 21 August 2019
Revised 20 December 2019
Accepted 24 December 2019

© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY. Published by BMJ.

¹Evidence Based Healthcare Postgraduation Program, Department of Medicine, Universidade Federal de São Paulo, São Paulo, Brazil

²UNDP/UNFPA/UNICEF/WHO/World Bank Special Programme of Research, Development and Research Training in Human Reproduction (HRP), Department of Sexual and Reproductive Health and Research, World Health Organization, Geneva, Switzerland

Correspondence to
Dr Maria Regina Torloni;
ginecologia@terra.com.br

ABSTRACT

Introduction The worldwide increase in unnecessary caesarean sections (CSs) is a major global health issue. Mass media campaigns have been used in several countries to reduce this trend. The objectives of this systematic review were to identify, critically appraise and synthesise the findings, including the barriers and enablers, of mass media campaigns directed at lay people to reduce unnecessary CS.

Methods We included any study design that reported health communication mass media campaigns directed at lay people with the specific objective of reducing unnecessary CS, created by any agent, in any format. We searched seven electronic databases without language restrictions, from inception to February 2019. Experts in the field were contacted.

Results The search yielded 14 320 citations; 50 were selected for full-text reading; and one was included. Six other reports were included. The seven campaigns were conducted in 2009–2017, mostly in Latin America. Most campaigns were independent efforts by non-governmental or activist organisations. Only one campaign conducted formative research and pretested the intervention.

All campaigns used indirect communication, mostly through internet channels; two campaigns also used direct communication with the public. None assessed their effects on CS rates. Only two campaigns measured their impacts on participants' knowledge, attitudes and birth preferences but only in the short term. The main barriers were lack of financial and human resources. The main enablers were the enthusiasm of volunteers, the participation of famous persons/celebrities and the involvement of communication professionals.

Conclusions There are few mass media campaigns directed at lay people to reduce CS. Most campaigns did not use key principles recommended for the creation and implementation of health communication interventions, and none assessed their effects in reducing CS rates. If media campaigns can play a role in modifying population views towards CS, there is a need for more rigorous studies including impact assessment.

PROSPERO registration number CRD42019120314.

INTRODUCTION

During the last decade, there has been a steady worldwide increase in the rate of caesarean sections (CSs). In 2014, the global

Key messages

What is already known?

- Scientific evidence does not support caesarean sections (CSs) for non-medical reasons, and numerous studies report increased maternal and perinatal risks associated with this type of delivery.
- The worldwide overuse of CS is a major global health issue that has led to considerable international debate.
- Mass media campaigns are recommended to change population health behaviours, but there are no systematic reviews on campaigns to reduce CS rates.

What are the new findings?

- We found four campaigns in Latin America, two in Europe and one in Asia directed at lay people with the specific objective of reducing unnecessary CS.
- Most campaigns were independent efforts by non-governmental or activist organisations, and did not use key principles recommended for the creation and implementation of health communication interventions.
- None assessed their effects on CS rates.

What do the new findings imply?

- There is a paucity of publications and data about mass media campaigns to reduce unnecessary CS.
- If mass media campaigns can play a role in modifying population views towards CS, there is a need for more rigorous studies including impact assessment.

CS rate reached 19.1%, a threefold increase from 1990, with large variations between and within countries. Currently, Latin America and the Caribbean have the highest CS rates (40.5%), followed by North America (32.3%), Oceania (31.1%), Europe (25%), Asia (19.2%) and Africa (7.3%).^{1 2} When performed for medical indications, a CS is a valuable intervention that can save the lives of mothers and babies, or avoid serious complications that can have long lasting consequences. However, scientific evidence does not support CS for non-medical reasons, and there are numerous reports of increased

maternal and perinatal risks associated with this type of delivery.^{3–8}

Maternal request, motivated by convenience, fear, misconceptions or cultural trends, is often mentioned as one of the key contributors to the increase in CS rates.^{9–18} The attitudes, beliefs and behaviours of adults are determined by numerous factors and can be influenced by various communication channels and the media.^{19–20} Reviews have shown the effectiveness of mass media campaigns targeted at reducing unhealthy habits or promoting healthy behaviours such as breast feeding or family planning, among others.^{21–27} However, significant changes in behaviour can be achieved only when campaigns are well designed and carried out at sufficient scale and intensity.^{21–28–36}

Over the last decade, there have been some campaigns targeted at the general population to reduce unnecessary CS. In view of the increasing concern about the worldwide rise in caesarean deliveries, potential negative consequences of CS for the health of mothers and infants, and the limited effectiveness of strategies tested to date to revert this trend,^{37–38} it is important to investigate the main characteristics, barriers and enablers, and the impact of these campaigns. However, to the best of our knowledge, up to the present, there has been no systematic review of these campaigns. This information gap motivated us to perform this review.

The objectives of this review were to identify, critically appraise and synthesise the information available on mass media campaigns directed at lay people to reduce unnecessary CSs. We assessed the main characteristics, the effectiveness, and the barriers and enablers to the creation and implementation of these campaigns.

METHODS

This review followed Cochrane methods and the MOOSE reporting recommendations.^{39–40} The protocol of the review was registered in the International Prospective Register of Systematic Reviews database.

Selection criteria

Types of studies

We included any type of report on mass media campaigns directed at lay people to reduce unnecessary CS or to increase vaginal delivery (VD). This included articles published in scientific journals or presented in events, as well as written, audio or video material available on the internet (including YouTube) or on social media channels (Facebook, Instagram and others). We included randomised, quasi-randomised, cluster randomised or non-randomised trials, as well as controlled before–after studies or interrupted time series studies or case reports. The reports had to describe the main components of the campaign in sufficient detail so that they could be understood and eventually adapted/replicated in other settings. We included only reports that stated that the

main objective (or primary outcomes) of the campaign was to reduce unnecessary CS or to increase VD.

Types of campaigns (exposure)

We defined a mass media campaign as an intervention that uses a set of organised communication activities to produce specific results or effects in a relatively large number of individuals, usually within a specific period of time.^{41–42} We included campaigns created or promoted by any agent, such as local or federal government authorities, national or international governmental or non-governmental organisations (NGOs), or specific associations/groups of individuals. We excluded messages created and posted/disseminated by single individuals or small groups (e.g., an activist promoting messages against CS in his/her blog or Facebook page), as these do not fulfil the classic definition of a campaign. We included campaigns whose main messages were the reduction of primary or repeat CS or the promotion of VD in otherwise healthy nulliparous or multiparous women. Campaigns using any type of communication channel were eligible. These included (1) written material disseminated through pamphlets, newspapers, magazines, mobile phones, billboards or posters; (2) audio messages disseminated on radio, television (TV) or phone (recorded messages); and (3) any form of material disseminated over the internet and social media (YouTube, Twitter, Facebook and Instagram).

We included campaigns that used either direct or indirect communication strategies. Direct communication strategies involve the transmission of messages through face-to-face interventions such as presentations, performances and group discussion sessions in clinics, schools or other public areas. Indirect communication campaigns use mass media instruments such as radio, TV, movies, newspapers or magazines to transmit their message to the general public during a specific period of time. This mode of communication is indirect because the consumer was not expecting this information, and it is presented to him/her in the context of another experience of his/her interest.^{42–44}

We included campaigns of any cost, size, duration or catchment area conducted as a stand-alone intervention or as part of a broader, multicomponent, intervention. We included campaigns with or without formative research to inform the creation and development of the campaign, as well as campaigns with or without previous research to assess the baseline and/or postintervention (before and after the campaign) outcomes of interest.

Types of participants

We included studies that described campaigns targeted at healthy pregnant women (of any parity, with or without previous CS), non-pregnant women of reproductive age or the general lay public (men and women of any age). We included campaigns targeted at laypersons of any nationality, gender, social class or educational level. We excluded campaigns aimed at specific high-risk groups,

such as people with HIV, as well as campaigns targeted exclusively at health professionals or institutions.

Types of outcomes

We included reports with or without quantitative assessments of the impact of the campaign on hard or soft outcomes. Hard outcomes were changes in the rates of CS or VD after the campaign, or in the exposed versus non-exposed groups. Soft outcomes were changes in knowledge, attitudes, beliefs or preferences for route of delivery on a sample or all participants after the campaign or in exposed versus non-exposed groups.

We collected and present the main barriers and enablers described by authors for the creation and implementation of their campaigns as part of the outcomes of this review.

Search strategy

The search strategy used the following general search words, adapted to each specific database: 'campaigns', 'birth' and 'preferences' (see online supplementary file 1 for the full search strategy). We ran the search in the following electronic databases, from inception until 1 February 2019, without language restrictions: MEDLINE (via PubMed), Embase (via Ovid), CINHALL, PsycINFO, Soclit, Web of Science, Popline, Global Index Medicus (which includes LILACS, AIM, IMSEAR, WPRIM and IMEMR) and EBSCO Multidatabase (which includes 37 databases). We also searched the website of the Center for Communication Programs (<https://ccp.jhu.edu/>). We screened the reference lists of all studies selected for full-text reading and contacted experts in the field to identify additional potentially relevant campaigns.

Process of study selection and data extraction

All citations identified from electronic databases were uploaded into Covidence (Veritas Health Innovation, Melbourne, Australia). After the exclusion of duplicates, two investigators independently screened the titles and abstracts of all citations to select potentially relevant studies. The full texts of these studies were retrieved and read; those that fulfilled the aforementioned selection criteria were included in the review. Discrepancies between reviewers were discussed until consensus was reached; if needed, a third reviewer was invited to arbitrate.

Two review authors extracted the data from each included report using a standard form created for this review (online supplementary file 2). Discrepancies were discussed until consensus was reached. We contacted authors to obtain additional information when needed.

Data analyses

We presented the main characteristics, barriers and enablers of the campaigns descriptively. If data were available, we planned to assess the effectiveness of each campaign by calculating the mean absolute and relative reduction in CS rate and 95% CIs. We planned to combine the results of similar studies and to calculate

pooled mean differences or mean standardised differences and 95% CIs. Due to lack of data, this was not possible.

Patient and public involvement statement

It was not appropriate or possible to involve patients or the public in the design, conduct, reporting or dissemination of our research.

RESULTS

The electronic search retrieved 14320 citations. After removing 118 duplicates, 14202 citations were screened; 14152 were excluded based on their title and abstract; and 50 were selected for full-text reading. We included 1 study⁴⁵; the other 49 were excluded mainly because they did not report a mass media campaign (online supplementary file 3). Six other reports were identified through contact with experts. The final review included seven reports that described campaigns conducted in Argentina (Cava S, Campaign for the Urgent Reduction of Unnecessary Caesarean Sections), Brazil,⁴⁵ Chile (Sadler M, Campaña #INNEcesareas Chile 2014–2015), Cyprus (Leontiou S, The Normal Birth Campaign 2014–2015: Activities, Challenges, and Opportunities), Iran (Akbari N, Majlesi M, Montazeri A, *et al*, 'No to Unnecessary Caesarean Section': Evaluation of a Mass Media Campaign on Women's Knowledge, Attitude and Intention for Mode of Delivery, in press), Italy (Montilla P, Merzagora F, Scolaro E, *et al*, Lessons from a Multidisciplinary Partnership Involving Women Parliamentarians to Address the Overuse of Caesarean Section in Italy, in press) and Puerto Rico (Nazario JOM, Campaña #INNEcesareas Puerto Rico 2012–2013) (online supplementary file 4). Six of the included campaigns were not described in printed journal articles; the Cyprus campaign was an oral presentation in a meeting; the campaigns conducted in Argentina, Chile and Puerto Rico were reported in webpages; and two campaigns were manuscripts in press (the Iranian and Italian campaigns). We contacted the authors of all seven reports to obtain additional details. The main characteristics of the campaign are listed in table 1.

Details of the objectives and the creation and implementation of the campaigns are presented in tables 2 and 3.

The majority of the campaigns were independent efforts of NGOs, activist organisations or associations strongly engaged in this topic. Only two (Brazil and Cyprus) had the involvement of the ministry of health, and three (Brazil, Cyprus and Italy) had the involvement of professional associations. Three of the campaigns (Chile, Cyprus and Iran) were stand-alone interventions. Only one of the campaigns (Iran) conducted formative research, and two campaigns (Brazil and Iran) used theories of behavioural change in the development of the campaign. Five of the campaigns (Argentina, Brazil, Cyprus, Iran and Italy) had communication specialists in

Table 1 Main characteristics of seven mass media campaigns to reduce unnecessary CS

Characteristic	N	Country
Region		
South America	3	Argentina, Brazil, Chile
Central America	1	Puerto Rico
Europe	2	Cyprus, Italy
Asia	1	Iran
Formative research prior to campaign		
Yes	1	Iran
No	6	Argentina, Brazil, Chile, Cyprus, Italy, Puerto Rico
Pretesting of messages		
Yes	1	Iran
No	1	Brazil
Unclear/ no information	5	Argentina, Chile, Cyprus, Italy, Puerto Rico
Target audience		
Reproductive age & pregnant women	1	Italy
General public	6	Argentina, Brazil, Chile, Cyprus, Iran, Puerto Rico
Communication strategies		
Direct*	2	Argentina, Brazil
Indirect†	7	Argentina, Brazil, Chile, Cyprus, Iran, Italy, Puerto Rico
Both	2	Argentina, Brazil
Communication channels		
Written material	3	Cyprus, Italy, Puerto Rico
Radio	1	Argentina
TV (open, paid, or close-circuit)	4	Argentina, Cyprus, Italy, Iran
Internet and social media	5	Argentina, Brazil, Chile, Italy, Puerto Rico
Celebrities/ famous spokespersons		
Yes	4	Argentina, Chile, Cyprus, Puerto Rico
No	3	Brazil, Iran, Italy
Duration of campaign		
range (min-max)	7 days - 34 months	
less than 1 month	2	Argentina, Iran
1–11 months	2	Chile, Cyprus

Continued

Table 1 Continued

12–34 months	2	Brazil, Italy
Unclear/ No information	1	Puerto Rico
Outcome measured		
None	5	Argentina, Chile, Cyprus, Italy, Puerto Rico
Knowledge about CS / VD	2	Brazil, Iran
Attitude toward CS /VD	2	Brazil, Iran
Preference for CS / VD	2	Brazil, Iran

*Use of interpersonal contact (face-to-face presentations, performance and group discussions).

†Use of mass media channels (such as TV or newspaper).
CS, caesarean section; TV, television; VD, vaginal delivery.

their creation teams, and four (Argentina, Chile, Cyprus and Puerto Rico) used authority figures/celebrities to disseminate the campaign messages. All campaigns used indirect communication: five disseminated messages through internet channels (Argentina, Brazil, Chile, Italy and Puerto Rico); three used TV spots (Cyprus, Iran and Italy); and four used pamphlets or posters (Argentina, Cyprus, Italy and Puerto Rico). Two of the Latin American campaigns also used direct communication with the target audience. The Argentinean campaign held group discussions with women. The Brazilian campaign⁴⁵ contacted the public through an interactive exhibition that combined different languages (digital art with theatrical techniques) and media (videos, photos, scenarios and panels).

Table 4 presents the main barriers and enablers during the creation and implementation of the campaigns. In both phases, lack of funding, human resources and institutional support was the main barrier, and volunteer work was one of the main enablers.

Table 5 presents the assessment of the effects of the intervention. Five campaigns did not assess any outcome (Argentina, Chile, Cyprus, Italy and Puerto Rico), and the other two (Brazil and Iran) measured changes in knowledge, attitude and preferred route of delivery among the exposed participants using written questionnaires.

Due to lack of data, we could not pool the results from individual studies into a combined estimate to assess the effectiveness of the campaigns. We therefore present a narrative description of the most important characteristics and findings of each campaign, as well as the barriers and facilitators reported by the authors.

Argentina: 'Campaign for the Urgent Reduction of Unnecessary Caesarean Sections'

There is no publication of this campaign. We obtained information through its website and interviews with the main coordinator. The campaign occurred in May 2009, during the 'International Week for Respecting Child-birth'. It was created by the Argentinean chapter of the

Table 2 Main objectives and messages of media campaigns to reduce unnecessary CS

Country	Objectives	Messages
Argentina	<ul style="list-style-type: none"> ► Raise awareness about unnecessary CS. 	<ul style="list-style-type: none"> ► Say 'no' to unnecessary CS. ► Women who deliver vaginally are an endangered species.
Brazil	<ul style="list-style-type: none"> ► Increase the public's knowledge. ► Promote cultural change to value normal birth. ► Reduce rates of CS and unnecessary interventions during childbirth. 	<ul style="list-style-type: none"> ► VD is an important and positive experience for women and children. ► VD is good for the health of women and children. ► Think critically about CS rates in Brazil.
Chile	<ul style="list-style-type: none"> ► Avoid unnecessary CS. ► Promote respectful necessary CS. ► Promote the use of Lamaze recommendations for a physiological birth. 	<ul style="list-style-type: none"> ► In Chile, there is an excess of CS without any medical indication. ► Birth is a physiological event that only rarely requires obstetric interventions like a CS. ► Follow Lamaze recommendations for a physiological birth (eg, wait for spontaneous onset of labour, continuous support during labour and avoid lithotomy position for delivery).
Cyprus	<ul style="list-style-type: none"> ► Inform and raise awareness about the mind and body benefits of normal childbirth for families and its impact on society. ► Strengthen and support couples' rights to choice in childbirth. ► Empower midwives and support their role in normal childbirth. ► Improve national perinatal indicators. 	<ul style="list-style-type: none"> ► Say 'yes' to VD. ► VD is the best choice for women, babies and families. ► VD empowers women and midwives. ► VD leads to stronger families and stronger society and improves health outcomes for all citizens.
Iran	<ul style="list-style-type: none"> ► Persuade pregnant women to choose spontaneous VD instead of unnecessary CS. 	<ul style="list-style-type: none"> ► Give birth naturally to guarantee your own health, that of the baby and of the next generation.
Italy	<ul style="list-style-type: none"> ► Raise awareness and foster action of authorities and women to reduce unnecessary CS in the country. ► Inform women about the indications and risks of CS. ► Disseminate and promote WHO's policies to achieve optimal CS rates. 	<ul style="list-style-type: none"> ► A CS is a life-saving surgical procedure that should be used when complications occur. ► For most women, giving birth should be a natural event. ► Mode of birth is your choice and should be discussed with your healthcare provider.
Puerto Rico	<ul style="list-style-type: none"> ► Prevent and reduce unnecessary CS and other unnecessary interventions during labour, delivery and the postpartum period. ► Empower Puerto Rican women to face the increasing rates of unnecessary CS as a public health issue. ► Promote and foster humanised births as the safest and healthiest option for delivery. 	<ul style="list-style-type: none"> ► You decide, be the protagonist and take charge (of your own birth). ► CS is a major surgery with risks for mothers and babies. ► Pregnancy, delivery and the postpartum period are natural events, not diseases that require medical interventions. ► Humanised childbirth is beneficial for the health of mothers and babies.

CS, caesarean section; VD, vaginal delivery.

international NGO network Red Latinoamericana y del Caribe para la Humanización del Parto y el Nacimiento (RELACAHUPAN) as part of a multicomponent intervention that took place during the week in the country. RELACAHUPAN is a Latin American network of NGOs that spans 20 countries.⁴⁶ The goals of this network are to improve the experience of birth, to promote humanised births, and to guarantee the rights of women to make informed decisions about pregnancy and childbirth, emphasising the benefits of natural birth. Volunteers from the NGO promoted group discussions about the messages of the campaign with women in the waiting areas of hospitals and public places, with poster displays. A short video featuring a famous TV and film actress (who worked pro bono) was created and broadcast on closed-circuit TV in public transportation and on the internet. The video had close-caption legends in several languages besides Spanish. Mama Cash, an international feminist

association, provided funding to RELACAHUPAN, which was used in part for the creation of this video. There is no information on the estimated cost of the campaign nor on the intensity of messaging, nor the number of persons exposed to the campaign. No outcomes were measured.

Brazil: 'The Senses of Birth Campaign'

The campaign occurred between 2015 and 2017 (34 months) in five cities in Brazil. It was created by health professionals and involved experts in communication and museum exhibits. The campaign was part of a long-term multisectorial initiative led by the municipal health department to increase VD in the city of Belo Horizonte ('BH pelo Parto Normal'), which started in 2007 and lasted 10 years. The campaign consisted of an interactive exhibition that combined digital art with theatrical techniques, videos, photos, scenarios and panels to promote an emotional experience, engage the visitors and

Table 3 Main characteristics of the creation and implementation of seven mass media campaigns to reduce unnecessary caesarean section

Characteristics	Argentina (2009)	Brazil (2015–2017)	Chile (2014–2015)	Cyprus (2014–2015)	Iran (2016)	Italy (2010–2011)	Puerto Rico (2012–2013)
Main creator	NGO (RELACAHUPAN Argentina)	Minas Gerais Federal University and Belo Horizonte Health Department	NGO (RELACAHUPAN Chile)	Council of Midwives Committee	Multi-professional expert panel	NGO (ONDA), WHO, National OB-GYN Association and female parliamentarians	Student association, Puerto Rico University Public Health School
Authorities involved in creation/support of campaign	No	Local health department, MoH and local professional association	No	National and local professional association and MoH	No	WHO, National Professional Association and parliamentarians	No
Main funding	Mama Cash (international feminist organisation) and voluntary work (for dissemination)	International, national and local scientific funding agencies, MoH and PAHO	Mostly voluntary work	Mostly voluntary work	Iran University of Medical Science	WHO Partnership for Maternal, Newborn & Child Health	NI
Type of intervention	Part of multicomponent intervention	Part of multicomponent intervention	Isolated intervention	Isolated intervention	Isolated intervention	Part of multicomponent intervention	NI
Design							
Theory used	No	Yes	No	No	Yes	No	NI
Formative research	No	No	No	No	Yes	No	No
Communication experts involved	Yes	Yes	No	Yes	Yes	Yes	NI
Prefesting	No	No	No	No	Yes	No	NI
Target public	General public	General public	General public	General public	General public	Women (pregnant and not pregnant)	General public
Medium							
Type of communication	Direct: group discussions in hospitals/public spaces Indirect: radio and video spots on closed TV in public transportation, internet (YouTube), poster and mailing lists	Direct: itinerant exhibition Indirect: internet channels	Indirect: five video clips posted on internet channels	Indirect: spot on open TV, posters and pamphlets	Indirect: spot on open TV	Indirect: spot on open and paid TV, internet channels, women's magazines and pamphlets	Indirect: video, informative posters and pamphlets on internet channels
Celebrities involved	Yes (actress)	No	Yes (actresses)	Yes (singer, actresses and first lady)	No	No	Yes (actors, singers and musicians)
Intensity	NI	IN	IN	NI	Open TV: 10 consecutive days	Paid TV: 3 times/day, 4 days/1 month. Open TV: 3 times/day, 15 days/1 month	NI

MoH, ministry of health; NGO, non-governmental organisation; NI, no information; ONDA, National Observatory for Women's Health; PAHO, Pan American Health Organisation; RELACAHUPAN, Red Latinoamericana y del Caribe para la Humanización del Parto y el Nacimiento; TV, television.

Table 4 Main barriers and enablers of seven mass media campaigns to reduce unnecessary caesarean section

Barriers	Enablers
Creation of the campaign	
1. Lack of human resources and time for volunteers to create campaign (Cyprus and Chile).	1. Involvement of communication professionals to design campaign (Brazil).
2. Lack of institutional support (Chile).	2. Support from ministry of health to allow public servants to develop the campaign during working hours (Cyprus).
3. Lack of funding (Chile).	3. Volunteer participation of celebrities in television spots (Cyprus and Argentina).
4. Difficulties of volunteers to organise meetings and plan campaigns (Chile).	4. Volunteer work of professionals in production of good quality videos (Cyprus).
5. Difficulties in finding key persons who would help with creation at no cost (Cyprus).	5. Support of other non-governmental organisations (Argentina).
Implementation of the campaign	
1. Lack of funding to transport and display exhibit (Brazil).	1. Trained mediators to help participants get in touch with their senses and emotions (Brazil).
2. Lack of local political support to promote campaign (Brazil).	2. Use of art to affect the sensibility of participants, to touch their hearts and not only their rational side (Brazil).
3. Finding key persons who would help with dissemination without charging (Cyprus).	3. Participation of entertainment celebrities (actresses and singers) in promotional material (Chile and Argentina).
4. Opposition of professional societies, authorities and universities to a campaign that was not created by them (Chile).	4. Avoidance of controversies and dissociation of campaign from extremist/radical groups initiatives or views (Chile and Argentina).
5. Hesitancy of women to accept campaign's message due to prevailing belief that medicalised births are safer for mother and baby (Chile).	5. Enthusiasm and good will of volunteers in promoting campaign (Chile).
6. Lack of funding for dissemination (Chile) and assessment of outcomes/effects of campaign (Argentina).	6. Good relationships with the key media stakeholders (Italy).
	7. No charge from owners of communication channels to disseminate campaign spots (Argentina).

encourage them to think critically. The free exhibition consisted of a 40 min guided interactive circuit that took place inside five containers parked on outdoor spaces in different locations (near shopping centres, in public parks, schools and universities). Immediately after going through the experience, visitors were invited to participate in group chats. This itinerant exhibit remained in each site for several weeks and was then packed up and transported to other locations by truck. The campaign also involved indirect communication through a website, Facebook, Instagram pages and a YouTube channel. The campaign cost approximately US\$350 000 per year to cover 12 exhibits in the five cities. It was funded by national scientific agencies, the Brazilian Ministry of Health, the Bill and Melinda Gates Foundation and the Pan-American Health Organisation.⁴⁵

A total of 42 170 persons of all ages visited the exhibits in the five cities. There were 8204 followers on Facebook, 4505 on Instagram and 2900 on YouTube. The authors used written questionnaires to measure changes in visitors' preferences for route of delivery, knowledge about risks of CS and opinion on VD. There were significant changes in these three measures (table 5). The investigators did not assess changes in the rates of CS as one of their outcomes.

Chile: 'InneCesareas'

There are no publications of this campaign. Data were obtained through interviews with one of its creators, who also provided the web content of the campaign. This national, internet-based campaign lasted 6 months (November 2013–May 2014). It was created by the Chilean chapter of the international NGO network RELACAHUPAN.⁴⁶ The campaign was a stand-alone intervention created by volunteers from RELACAHUPAN Chile. The messages were delivered through five short videos posted on YouTube, Facebook and Twitter. Famous soap opera actresses participated in the videos, as well as local authority figures (famous obstetricians and midwives), at no cost. A journalist produced the videos. The creators of the campaign promoted the initiative by appearing in talk shows and interviews. The campaign cost US\$400 for the production of the videos. This was obtained through fundraising activities among RELACAHUPAN Chile members. There were approximately 50 000 visits to the YouTube videos. The effectiveness of the campaign was not assessed.

Cyprus: 'Normal Birth Campaign'

This report was presented only as an oral communication in an international meeting. Additional details were

Table 5 Assessment of effects of seven mass media campaigns to reduce unnecessary CS

Characteristics	Argentina (2009)	Brazil (2015–2017)	Chile (2014–2015)	Cyprus (2014–2015)	Iran (2016)	Italy (2010–2011)	Puerto Rico (2012–2013)
Outcomes assessed	None	1. Preference for CS. 2. Knowledge of risks/benefits of CS. 3. Opinion on VD.	None	None	1. Knowledge about childbirth. 2. Attitude towards VD and CS. 3. Intended route of delivery.	None	None
Period of outcome assessment	NA	Immediately after exhibit	NA	NA	10 days after exposure	NA	NA
Sample assessed	NA	1. Preference for CS: n=1933 general public and n=1287 pregnant women. 2. Knowledge of risks/benefits of CS: n=1933 general public and n=1287 pregnant women. 3. Opinion about VD: n=17 501 visitors.	NA	NA	466 pregnant women (194 had seen the TV spot; 272 had not seen it). All women had no previous CS, were mostly in the second and third trimesters of pregnancy, and attending antenatal care in public and private clinics in Teheran	NA	NA
Tool used to assess effect	NA	Written questionnaires immediately before and after the exhibit			Written questionnaires before and after TV campaign		
Effects	NA	1. Decrease in preference for CS (14.7%×10.4%, p=0.006). 2. Increase in good/very good knowledge about CS risks (50.5%×71.5%, p<0.001). 3. Decrease in opinion that VD was very bad or bad (12.2%×1.9%, p<0.001).	NA	NA	Changes in 194 women exposed to campaign: 1. Increase in knowledge scores (p=0.008). 2. Increase in attitude scores towards VD (p=0.05). 3. Decrease in intention to deliver by CS (39.2%×24.7%, p=0.004). 4. No changes in attitude scores towards CS.	NA	NA

CS, caesarean section; NA, not applicable; TV, television; VD, vaginal delivery.

obtained by contacting the presenter. This national campaign took place in Cyprus between 2014 and 2015 (12 months). It was designed as a stand-alone intervention by the Council of Midwives Committee and the Ministry of Health. Communication experts from TV channels volunteered to help create the campaign's video clip. The campaign was supported by the country's first lady, national specialist societies, religious authorities, several NGOs and civil trade unions. The messages were delivered through indirect communication, including posters, banners and pamphlets in local hospitals, and press releases that were sent through emails and mass media posts. The central piece of the campaign was a short video clip, with local female celebrities, that was disseminated in local open TV channels during 12 months. There is no information on the frequency of the campaign messages. The cost of the campaign was approximately €5000, and it was funded by the Cyprus Nurses and Midwives Association, trade unions and the Nursing and Midwives Council. There is no information on the estimated number of persons exposed to the campaign. No outcomes were measured or reported. The effectiveness of the campaign was not assessed.

Iran: 'No to Unnecessary Caesarean Section'

This prepublication manuscript (obtained from the authors) describes a national campaign conducted in Iran during 10 days in April 2016. The campaign was created by an expert panel of nine persons (academic obstetricians and midwives, health education experts and advertising professionals). There is no information on the cost of the campaign. The campaign consisted of a short video clip that was broadcast on four of the country's eight open TV channels during 10 consecutive days and in antenatal care clinics (on closed-circuit TV). There is no information on the estimated number of persons exposed to the TV campaign. The authors measured changes in knowledge, attitude and preferred route of delivery of 466 pregnant women without a previous CS and living in Teheran for at least 6 months, who were managed in public and private antenatal care clinics in that city. The authors used written questionnaires to assess all participants before and after the campaign. Most of the participants (58%, n=272) stated they had not seen the TV spots when they answered the second questionnaire. The 194 exposed women had a significant increase in knowledge and attitude scores towards VD

and a significant decrease in the proportion of pregnant women who intended to have a CS, but no significant changes in attitude towards CS (table 5).

Italy: 'Caesarean Section: When and Why'

A manuscript describing this campaign is in press and was shared by the authors. Additional information was obtained by contacting representatives of various partners involved in this initiative. The nationwide campaign took place between January 2010 and September 2011 (21 months). The campaign was created by ONDa (National Observatory for Women's Health), an NGO focused on women's health, with the participation of the World Health Organization (WHO), the Italian Society of Gynaecologists and Obstetricians and a group of Italian female bipartisan parliamentarians. The campaign was part of a multicomponent intervention that included technical meetings at WHO to increase the awareness of Italian parliamentarians about the overuse of CS and to foster political action at national and regional levels to promote vaginal births and to reduce CS. The main elements of this campaign were a TV spot (created pro bono by a professional Italian marketing and advertising company), an online survey disseminated by a national women's magazine and an educational brochure about indications and risks of CS. A short TV spot was initially broadcast on open and paid TV channels three to four times/day for up to 2 weeks and then posted on YouTube and campaign websites. The largest Italian women's magazine featured a text about CS in Italy and invited readers to answer an online anonymous survey about their preferences for and knowledge about the risks of CS versus VD. The magazine also distributed a special brochure with information on CS, including indications and risks. This brochure was also available in hospitals and clinics. There is no information on the estimated number of persons exposed to the campaign. No outcomes were measured, and the effectiveness of the campaign was not assessed.

Puerto Rico: 'INNEcesareas'

There are no publications of this campaign. Data were obtained through personal contact with the campaign coordinator. This national campaign was created by the Maternal and Child Health Student Association of Puerto Rico's University School of Public Health. The messages were delivered using indirect communication through a hip-hop video posted on the campaign webpage and internet channels (YouTube, Facebook and Twitter). Celebrities (actors, musicians and other artists) acted as spokespersons pro bono. No outcomes were measured, and there is no information on the number of persons exposed to the campaign or its effectiveness.

DISCUSSION

This review identified seven mass media campaigns directed at lay people with the specific objective of reducing unnecessary CS, only one of which was published in a scientific

journal. In their creation phase, only one of the campaigns used formative research to inform the design and implementation of the intervention and obtained input from the target population (through pilot testing) before being launched. The involvement of health authorities was not frequent. None of the campaigns assessed its effects on CS rates. Only two campaigns measured their impacts on participants' knowledge, attitudes and birth preferences, but only in the short term. The main barriers to the creation and implementation of the campaigns were lack of financial and human resources. The main enablers were the enthusiasm of the volunteers, the participation of famous persons/celebrities and the involvement of communication professionals.

The lack of formative research and testing in six of the seven campaigns was surprising. Formative research is essential to assess knowledge, awareness and beliefs of the population, to understand local cultural norms, to ensure that the content and format of the campaign are appropriate and acceptable, and to identify the most credible and most persuasive channels and spokespersons.^{29 30 41–43 47}

Several campaigns depicted messages about the value of VD. Overall, in the material produced with the aim of reducing CS, the benefits of VD are often lost in the midst of messages focused on the potential adverse consequences of CS. We believe this perspective is important and should be included in future campaigns since women welcome this information and value what is best for their babies and themselves.⁴⁸

The specific types of communication used by the campaigns varied, but they were mostly indirect, that is, did not involve personal contact with the target audience. Traditionally, most mass media campaigns rely on the use of posters, handouts, public service announcements and presentations, along with information disseminated in newspapers, magazines, radio and TV.⁴⁷ Three of the campaigns included in this review (Chile, Iran and Italy) created a short film that was produced pro bono and broadcast on open TV. The costs involved in using TV can be a barrier, and funding should be secured. In recent years, with the availability and popularisation of new media technologies, mass media campaign dissemination has expanded via websites, social networking, smart phone apps and online messaging platforms.^{49 50} Studies on the effectiveness of these new media health campaigns report mixed results.^{51–54} Most of the campaigns included in our review used internet media, and the two that measured the number of accesses (Brazil and Chile) reported a high number of visitors to their sites, which is an indication of the potential outreach of and interest in these campaigns.

We could not assess the effectiveness of the campaigns in reducing the rates of CS because none of the included studies measured this outcome. Soft outcomes (knowledge, attitude, behaviour or preferences) were assessed in two campaigns (Brazil and Iran) but only immediately after exposure to the campaign. The duration of mass media effects, that is, the persistence of the

positive effects of mass media on knowledge, attitude, behaviour or preferences over time, is considered an important element in assessing the effectiveness of mass media interventions.^{27 28 44} This was not reported by any of the campaigns. The lack of an evaluation component of campaigns is not uncommon. In a systematic review on campaigns about child survival, Naugle and Hornik reported that 54% of over 100 studies did not measure and report any behavioural or health outcome.²⁷

Our review has several strong points. To our knowledge, this is the first review on this topic. We designed a broad and sensitive search strategy and used rigorous methodology to reduce bias. Extensive efforts were undertaken to contact experts in the field to identify campaigns not identified through electronic searches. Six of the seven included campaigns were uncovered using this strategy. We acknowledge that, despite our extensive search, we may have missed relevant reports. An additional limitation of this review is that many reports were incomplete and did not provide all relevant details, which can be an obstacle to the replication of similar campaigns in other settings. Due to the lack of details and outcome assessments, we are not able to infer what elements of a mass media campaign on this topic could be most associated with success.

Mass media campaigns have been recommended as a key component of comprehensive strategies aiming at behavioural change.⁴⁷ However, our findings indicate that most of the campaigns to reduce unnecessary CS are almost 'amateur' efforts relying on volunteers, and counting on people's good will and pro bono work to support their activities, without rigorous evaluation to assess whether the campaigns actually achieved their aim or not. There is a clear need for more, adequately designed and reported studies in this area, including research on cost-effectiveness.

Importantly, the success of any campaign to reduce unnecessary CS will depend on the existence of facilities that offer women safe and adequately resourced environments to encourage and support them during labour and vaginal birth. Suboptimal quality of care, disrespect and abuse, insufficient communication or perverse relationships between women and providers must be addressed for women to prefer VD.^{48 55} Successful mass media campaigns targeted at the general public can play an important role in raising awareness. However, high CS rates are not due exclusively to the preferences of women. Therefore, initiatives to reduce CS need to involve and target the other contributors to this phenomenon, such as healthcare providers, hospitals, policy makers and organisations responsible for paying for medical procedures.^{37 55}

CONCLUSIONS

This review found seven mass media campaigns directed at laypersons to reduce CS, only one of which was published. Most campaigns did not use key principles

recommended for the creation and implementation of health communication interventions, and none assessed the effects of the campaigns in reducing CS rates. There is a need for more well-designed and well-reported studies in this area, including rigorous research to assess impact.

Acknowledgements We are grateful to the authors and the experts in the field who provided information about campaigns. After discussion, we did not include some campaigns because they did not fulfil the selection criteria of this review. We are nevertheless very grateful for the time of all persons whom we contacted. We thank Ibón Olza, University of Alcalá de Henares, Spain, for providing the contacts of campaigns in Latin America; Soo Downe, Lancashire University, UK, for the information on the UK 'Campaign for Normal Birth'; Mary Giammarino, March of Dimes, for the information on the US 'Campaign Healthier Babies Are Worth the Wait'; Linda Harmon and Amanda Darvill, Lamaze International, for the information on the US campaign 'Push for Your Baby'; Sonia Cavia and Gilda Vera López, Red Latinoamericana y del Caribe para la Humanización del Parto y el Nacimiento (RELACAHUPAN) for the information on the 'Campaign for the Urgent Reduction of Unnecessary Caesarean Section' in Argentina; Sonia Lansky for additional information on the campaign 'Senses of Birth' in Brazil; Michelle Sadler, RELACAHUPAN, for the information on the '#INNEcesareas' in Chile; Javier Morales Nazario, RELACAHUPAN, for the information on the '#INNEcesareas' in Puerto Rico; Catherine Ruhl, AWHONN, for the information on the US campaign 'Don't Rush Me... Go the Full 40'; Nahid Akbari, Iran University of Medical Science, for information on the Iranian campaign 'No to Unnecessary Caesarean sections'; Stella Leontiou, Cyprus Nurses and Midwives Association, for the information on the Cyprus 'Normal Birth Campaign'; and Pilar Montilla for information on the campaign in Italy.

Contributors MRT contributed to the design of the study, acquired, analysed and interpreted the data, and drafted the first version of the manuscript. VB made substantial contributions to the acquisition, analysis and interpretation of the data, and revised the manuscript for important intellectual content. APB conceived the study, acquired, analysed and interpreted the data, and revised the manuscript for important intellectual content. All authors approved the version submitted, agreed to be accountable for all aspects of the work and ensured that questions related to the accuracy or integrity of any part of the work would be appropriately investigated and resolved.

Funding This review was funded by UNDP/UNFPA/UNICEF/WHO/World Bank Special Programme of Research, Development and Research Training in Human Reproduction. Department of Sexual and Reproductive Health and Research, WHO, Geneva, Switzerland.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement No data are available.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Maria Regina Torloni <http://orcid.org/0000-0003-4944-0720>

Vanessa Brizuela <http://orcid.org/0000-0002-4860-0828>

Ana Pilar Betran <http://orcid.org/0000-0002-5631-5883>

REFERENCES

- 1 Betrán AP, Ye J, Moller A-B, *et al*. The increasing trend in caesarean section rates: global, regional and national estimates: 1990-2014. *PLoS One* 2016;11:e0148343.
- 2 Boerma T, Ronsmans C, Melesse DY, *et al*. Global epidemiology of use of and disparities in caesarean sections. *Lancet* 2018;392:1341-8.
- 3 Ye J, Zhang J, Mikolajczyk R, *et al*. Association between rates of caesarean section and maternal and neonatal mortality in the 21st century: a worldwide population-based ecological study with longitudinal data. *BJOG: Int J Obstet Gy* 2016;123:745-53.

- 4 Betran AP, Torloni MR, Zhang JJ, *et al.* WHO statement on caesarean section rates. *BJOG: Int J Obstet Gy* 2016;123:667–70.
- 5 Gibbons L, Belizan JM, Lauer JA, *et al.* Inequities in the use of caesarean section deliveries in the world. *Am J Obstet Gynecol* 2012;206:331.e1–9.
- 6 Victora CG, Barros FC. Beware: unnecessary caesarean sections may be hazardous. *Lancet* 2006;367:1796–7.
- 7 Hansen AK, Wisborg K, Uldbjerg N, *et al.* Risk of respiratory morbidity in term infants delivered by elective caesarean section: cohort study. *BMJ* 2008;336:85–7.
- 8 Liu S, Liston RM, Joseph KS, *et al.* Maternal mortality and severe morbidity associated with low-risk planned caesarean delivery versus planned vaginal delivery at term. *Can Med Assoc J* 2007;176:455–60.
- 9 Dursun P, Yanik FB, Zeyneloglu HB, *et al.* Why women Request caesarean section without medical indication? *J Matern Fetal Neonatal Med* 2011;24:1133–7.
- 10 Fuglenes D, Aas E, Botten G, *et al.* Why do some pregnant women prefer caesarean? the influence of parity, delivery experiences, and fear. *Am J Obstet Gynecol* 2011;205:45.e1–9.
- 11 Handelzalts JE, Fisher S, Lurie S, *et al.* Personality, fear of childbirth and caesarean delivery on demand. *Acta Obstet Gynecol Scand* 2012;91:16–21.
- 12 Handelzalts JE, Fisher S, Sadan O, *et al.* Object relations, unconscious defences and fear of childbirth, as reflected in maternal-request caesarean section. *J Reprod Infant Psychol* 2017;35:91–102.
- 13 Klein MC. Caesarean section on maternal request: a societal and professional failure and symptom of a much larger problem. *Birth* 2012;39:305–10.
- 14 Barber EL, Lundsberg LS, Belanger K, *et al.* Indications contributing to the increasing caesarean delivery rate. *Obstet Gynecol* 2011;118:29–38.
- 15 Druzin ML, El-Sayed YY. Caesarean delivery on maternal Request: wise use of finite resources? A view from the trenches. *Semin Perinatol* 2006;30:305–8.
- 16 Jackson NV, Irvine LM. The influence of maternal Request on the elective caesarean section rate. *J Obstet Gynaecol* 1998;18:115–9.
- 17 Kottmel A, Hoesli I, Traub R, *et al.* Maternal Request: a reason for rising rates of caesarean section? *Arch Gynecol Obstet* 2012;286:93–8.
- 18 Marx H, Wiener J, Davies N. A survey of the influence of patients' choice on the increase in the caesarean section rate. *J Obstet Gynaecol* 2001;21:124–7.
- 19 Handfield B, Turnbull S, Bell RJ. What do obstetricians think about media influences on their patients? *Aust N Z J Obstet Gynaecol* 2006;46:379–83.
- 20 Moyer CA, Vishnu LO, Sonnad SS. Providing health information to women. The role of magazines. *Int J Technol Assess Health Care* 2001;17:137–45.
- 21 Buchbinder R, Gross DP, Werner EL, *et al.* Understanding the characteristics of effective mass media campaigns for back pain and methodological challenges in evaluating their effects. *Spine* 2008;33:74–80.
- 22 Werder O. Battle of the bulge: an analysis of the obesity prevention campaigns in the United States and Germany. *Obesity Reviews* 2007;8:451–7.
- 23 Cohen EL, Shumate MD, Gold A. Anti-Smoking media campaign messages: theory and practice. *Health Commun* 2007;22:91–102.
- 24 McDivitt JA, Zimicki S, Hornik R, *et al.* The impact of the Healthcom mass media campaign on timely initiation of breastfeeding in Jordan. *Stud Fam Plann* 1993;24:295–309.
- 25 Kaufman MR, Harman JJ, Smelyanskaya M, *et al.* "Love me, parents!": impact evaluation of a national social and behavioral change communication campaign on maternal health outcomes in Tanzania. *BMC Pregnancy Childbirth* 2017;17:305.
- 26 Shefner-Rogers CL, Sood S. Involving Husbands in Safe Motherhood: Effects of the SUAMI SIAGA Campaign in Indonesia. *J Health Commun* 2004;9:233–58.
- 27 Naugle DA, Hornik RC. Systematic review of the effectiveness of mass media interventions for child survival in low- and middle-income countries. *J Health Commun* 2014;19:190–215.
- 28 Anker AE, Feeley TH, McCracken B, *et al.* Measuring the effectiveness of Mass-Mediated health campaigns through meta-analysis. *J Health Commun* 2016;21:439–56.
- 29 Donovan RJ. Steps in planning and developing health communication campaigns: a comment on CDC's framework for health communication. *Public Health Rep* 1995;110:215–7.
- 30 Dutta-Bergman MJ. Theory and practice in health communication campaigns: a critical interrogation. *Health Commun* 2005;18:103–22.
- 31 O'Keefe DJ, Jensen JD. The relative Persuasiveness of Gain-Framed Loss-Framed messages for encouraging disease prevention behaviors: a meta-analytic review. *J Health Commun* 2007;12:623–44.
- 32 Randolph W, Viswanath K. Lessons learned from public health mass media campaigns: marketing health in a crowded media world. *Annu Rev Public Health* 2004;25:419–37.
- 33 Sellers DE, Crawford SL, Bullock K, *et al.* Understanding the variability in the effectiveness of community heart health programs: a meta-analysis. *Soc Sci Med* 1997;44:1325–39.
- 34 Soames Job RF. Effective and ineffective use of fear in health promotion campaigns. *Am J Public Health* 1988;78:163–7.
- 35 Wallack LM. Mass media campaigns: the odds against finding behavior change. *Health Educ Q* 1981;8:209–60.
- 36 O'Connor M, McGowan K, Jolivet RR. An awareness-raising framework for global health networks: lessons learned from a qualitative case study in respectful maternity care. *Reprod Health* 2019;16:1.
- 37 Chen I, Opiyo N, Tavender E, *et al.* Non-Clinical interventions for reducing unnecessary caesarean section. *Cochrane Database Syst Rev* 2018;9:CD005528.
- 38 Hartmann KE, Andrews JC, Jerome RN, *et al.* *Strategies to reduce caesarean birth in low-risk women*. Rockville (MD), 2012.
- 39 Higgins JPT, Green S. *Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 [updated March 2011]*. The Cochrane Collaboration, 2011.
- 40 Stroup DF, Berlin JA, Morton SC, *et al.* Meta-Analysis of observational studies in epidemiology: a proposal for reporting. meta-analysis of observational studies in epidemiology (moose) group. *JAMA* 2000;283:2008–12.
- 41 Public Health Service, National Institutes of Health, [Office of Cancer Communications, National Cancer Institute]. *Making health communication programs work: a Planner's guide*. Bethesda, Md: Public Health Service, National Institutes of Health, [Office of Cancer Communications, National Cancer Institute], 2002.
- 42 Rogers EM SJ. Communication campaigns. In: Berger CR CS, ed. *Handbook of communication science*. Newbury Park, California, USA: Sage, 1987: 817–46.
- 43 Snyder LB. Development communication campaigns. In: Gutykunst WB MB, ed. *Handbook of international and Intercultural communication*. 2nd ed. Thousand Oaks, Calif: Sage, 2001: 457–78.
- 44 Snyder LB. How effective are mediated health campaigns? In: Rice RAC, ed. *Public information campaigns*. 3rd ed. Thousand Oaks, Calif: Sage, 2001: 181–90.
- 45 Lansky S, Oliveira BJ, Peixoto ERM, *et al.* The senses of birth intervention to decrease caesarean and prematurity rates in Brazil. *Int J Gynecol Obstet* 2019;145:91–100.
- 46 RELACAHUPAN. Objetivos de la red Latinoamericana Y del Caribe para La Humanización del Parto Y El Nacimiento, 2000. Available: <http://www.soniacavia.com.ar/relacahupan.com.ar/lared.html>
- 47 Wakefield MA, Loken B, Hornik RC. Use of mass media campaigns to change health behaviour. *Lancet* 2010;376:1261–71.
- 48 Kingdon C, Downe S, Betran AP. Women's and communities' views of targeted educational interventions to reduce unnecessary caesarean section: a qualitative evidence synthesis. *Reprod Health* 2018;15:130.
- 49 Korda H, Itani Z. Harnessing social media for health promotion and behavior change. *Health Promot Pract* 2013;14:15–23.
- 50 Roland D. Social media, health policy, and knowledge translation. *J Am Coll Radiol* 2018;15:149–52.
- 51 Cushing CC, Steele RG. A meta-analytic review of eHealth interventions for pediatric health promoting and maintaining behaviors. *J Pediatr Psychol* 2010;35:937–49.
- 52 Firestone R, Rowe CJ, Modi SN, *et al.* The effectiveness of social marketing in global health: a systematic review. *Health Policy Plan* 2017;32:110–24.
- 53 Guse K, Levine D, Martins S, *et al.* Interventions using new digital media to improve adolescent sexual health: a systematic review. *J Adolesc Health* 2012;51:535–43.
- 54 Swanton R, Allom V, Mullan B. A meta-analysis of the effect of new-media interventions on sexual-health behaviours. *Sex Transm Infect* 2015;91:14–20.
- 55 Betrán AP, Temmerman M, Kingdon C, *et al.* Interventions to reduce unnecessary caesarean sections in healthy women and babies. *Lancet* 2018;392:1358–68.