

**Supplementary Information for Manuscript: “Do evidence-based practices during childbirth prevent perinatal mortality? A post-hoc analysis of 3,3274 births in Uttar Pradesh, India”**

**Table A: Observed Mortality by Seven Days Postpartum**

<b>Mortality N=3274</b>	<b>N(%)</b>
Perinatal mortality	166 (5.07)
Stillbirth	56 (1.71)
Early Neonatal Death	110 (3.36)

**Table B: Stillbirth Rate (stillbirths per 1000 births) by Practice Adherence in 30 facilities in Uttar Pradesh, India**

	Stillbirth Rate: Deaths per 1000 births	
	Practice Performed (deaths/cases)	Practice Not Performed (deaths/cases)
<b>Birth companion present</b>	18.5 (50/2702)	0.0 (0/6)
<b>Partograph Started</b>	0.0 (0/14)	18.6 (50/2694)
<b>Proper Hand Hygiene**</b>	18.8 (8/426)	16.9 (41/2425)
<b>Oxytocin Not Administered Before Birth</b>	12.7 (16/1262)	20.8 (33/1589)
<b>Clean Towel Available</b>	15.3 (23/1508)	19.4 (26/1343)
<b>Clean Blade Available</b>	16.9 (40/2371)	18.8 (9/480)
<b>Cord Tie Available</b>	16.9 (48/2838)	76.9 (1/13)
<b>Mucus Extractor Available</b>	16.2 (44/2721)	38.5 (5/130)
<b>Neonatal Bag Available</b>	15.8 (43/2731)	50.0 (6/120)
<b>Clean Pads Available</b>	16.7 (35/2099)	18.6 (14/752)
<b>Oxytocin Administered Postpartum</b>	N/A	N/A
<b>Birth Companion Present (Postpartum)</b>	17.2 (49/2843)	0.0 (0/7)
<b>Baby Weight Taken</b>	N/A	N/A
<b>Baby's Temperature Taken</b>	N/A	N/A
<b>Skin-to-Skin Warming Initiated</b>	N/A	N/A
<b>Skin-to-Skin Warming Maintained for One Hour</b>	N/A	N/A
<b>Breastfeeding Initiated</b>	N/A	N/A
<b>Mother's Blood Pressure Taken (At Any Time)</b>	15.2 (18/1188)	18.5 (38/2060)
<b>Mother's Temperature Taken (At Any Time)</b>	11.9 (11/923)	19.4 (45/2325)