

Improved knowledge on substandard and falsified (SF) medical products through a dedicated course for pharmacy students at three universities in sub-Saharan Africa

Zuzana Kusynová,^{1,2} Youssra Bais,^{1,2} Hendrika A van den Ham,¹ Aukje K Mantel-Teeuwisse,¹ Gisele Etame-Loe,³ Eliangiringa Kaale,⁴ Serigne Omar Sarr,⁵ Fatima Guiet-Mati,⁶ Pernelle Bourdillon-Esteve⁶

To cite: Kusynová Z, Bais Y, van den Ham HA, *et al*. Improved knowledge on substandard and falsified (SF) medical products through a dedicated course for pharmacy students at three universities in sub-Saharan Africa. *BMJ Global Health* 2023;**6**:e009367. doi:10.1136/bmjgh-2022-009367

Handling editor Seye Abimbola

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjgh-2022-009367>).

Received 14 April 2022
Accepted 7 November 2022



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Hendrika A van den Ham;
H.A.vandenHam@uu.nl

ABSTRACT

Background Too few pharmacists receive formal training on substandard and falsified (SF) medical products. Strengthening knowledge across pharmacists is considered a moral and ethical duty of academia, that is, to build the health systems' capacities to combat this global health threat these poor-quality products represent. This study therefore aimed to evaluate whether a dedicated educational course for undergraduate pharmacy students can improve their knowledge on these products.

Methods A survey was conducted at three sub-Saharan universities. Knowledge was assessed through scores on a 20-point questionnaire with questions related to the course content. Scores were compared before and after the course, and a linear mixed-effects model analysis was used to analyse score differences. Students were furthermore asked for feedback and self-assessment. In addition, teachers were interviewed on the context of the course introduction. These data were analysed descriptively.

Results Among 335 out of 355 students who completed the survey (n=41/53 in Cameroon, n=244/252 in Senegal and n=50/50 in Tanzania), knowledge of SF medical products was enhanced, with increase in all countries, overall, by 3.5 (95% CI 3.1 to 3.9) score points. Students improved in all offered modules in each country. Students confirmed their improvement through self-assessment. The course was well received among students and teachers. Barriers included time constraints and access to practical means (equipment availability, room allocation, internet accessibility and affordability). These barriers can be overcome by key enablers such as the support from university leadership and early involvement of the university in the course design.

Conclusions The course improved students' knowledge on SF medical products. These findings encourage further full implementation of this course in existing curricula beyond the pilot and can inform possible future scale-up. This has a potential for reinforcing the capacity of health systems to protect communities from SF medicines, by empowering all pharmacist across the health systems to intervene.

WHAT IS ALREADY KNOWN ON THIS TOPIC

- ⇒ Substandard and falsified (SF) medicinal products are a serious threat to public health and are especially found in those countries that have under-resourced regulatory authorities.
- ⇒ Although pharmacists can play an essential role in protecting the public against SF medicines, too few are trained well enough to fulfil this responsibility.

WHAT THIS STUDY ADDS

- ⇒ A jointly developed educational course on SF medicinal products for undergraduate pharmacy students showed an improvement of knowledge in three pilot countries: Cameroon, Senegal and Tanzania.
- ⇒ Teachers that were involved in the implementation of the course indicated that the readily available teaching materials were considered very helpful.
- ⇒ A barrier for the successful implementation was the availability of sufficient time for preparation and the deployment of the course.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

- ⇒ This study can inform other universities that wish to implement a dedicated course on SF medicines in their pharmacy curriculum.
- ⇒ Empowering all pharmacist across the health systems to address the socially and ethically unacceptable issue of SF medical products has a potential for reinforcing health systems' capacity and safeguarding public health.

INTRODUCTION

Substandard and falsified (SF) medical products are a threat to public health.^{1 2} In addition to the obvious harm caused to patients exposed to these poor-quality medical products, they also contribute to antimicrobial resistance, undermine immunisation

programmes, erode confidence in healthcare professionals, medicines and health systems, and waste precious financial resources through prolonged treatment.^{3–6} They are morally unacceptable because they are not ordinary consumer goods and patients have the right to expect that the products they receive are safe and efficacious.

Countries with under-resourced regulatory authorities tend to be most vulnerable to SF medical products reaching patients.^{7–9} Between September 2012 and February 2022, the WHO Global Surveillance and Monitoring System has received 3169 case reports from 139 countries.^{8 10} Thirty-eight per cent of reports originate from the African region.⁷ Most serious incidents of SF medicines are often reported by healthcare professionals, in particular pharmacists.^{8 11} However, too few pharmacists receive formal training on this issue during their undergraduate education.^{12–14}

Pharmacists are important players at the interface between the health system and the patient (end-users) community. They constitute the last defence/checkpoint between patients and their treatment and thus have both ethical and professional responsibility to minimise any health risks. In order to address the training gap and advance the essential role of pharmacists in pharmaceutical and health systems, a course on SF medical products for undergraduate pharmacy students was developed as part of a pilot project funded by the European Commission, designed and implemented by the WHO and the International Pharmaceutical Federation (FIP) in 2019–2021 in close collaboration with the pilot universities.^{15 16} This course was then deployed as a new component into the curriculum for pharmacists in the sub-Saharan African region. It includes detailed information relating to the root causes of SF medical products, the products most at risk, early warning signals of their presence in the supply chain, how to avoid, detect and report SF medical products, and advise patients and consumers.¹⁵

The main objective of this study was to assess the change in knowledge about the topic of SF medical products in undergraduate pharmacy students who participated in this course, which was delivered at three selected pilot pharmacy schools in sub-Saharan Africa (Cameroon, Senegal and Tanzania). The study also looked at self-assessment of the students' knowledge prior to and after the course, and whether they found the course useful and worthy to recommend to other students. In addition, teachers were interviewed on the context of the course introduction and motivation of the students as well as main enablers and barriers of its implementation.

METHODS

Course characteristics

The course was developed by an informal expert group, composed by pharmacy experts from academia and practice from FIP, pilot universities, project partners of the Commonwealth Pharmacists Association (CPA) and La

Conférence Internationale des Ordres de Pharmaciens Francophones (CIOPF), local African experts with regulatory background and WHO, who sponsored the pilot project. It was developed through a consultative process; several meetings with multiple rounds of review and consultation were held until a consensus was reached. The consultation took into account multicultural dimension. The course is available in both English and French language. It is designed to cover comprehensive information about SF medical products in order to teach pharmacy students how to avoid, detect and report SF medical products, and how to advise affected patients. The course is composed of six modules on SF medical products: module A on general introduction; module B on identification of medical products at risk; module C on prevention (from entering the supply chain); module D on detection strategies; module E on reporting; and module F on intervention after coming in contact with them.¹⁵ As this was a generic course, the pilot universities in Cameroon, Senegal and Tanzania were responsible for considering necessity of adjustment of the content, using the 'adopt and adapt principle'¹⁵ to contextualise the content and for flexibly incorporating the course into universities' existing curriculum based on their needs.¹⁵

The teaching materials included case studies. The module on detection (module D) also included practical lessons, with universities teaching visual inspection (all) and analytical inspection with qualitative and semi-quantitative investigations of medicines (Senegal and Tanzania).

The course was ready for deployment as of 2021. It was deployed face-to-face despite challenges and delays related to the COVID-19 pandemic. Each university deployed all six modules. An overview of the characteristics of the participating universities is included in [table 1](#).

Study sample

A total of 355 students participated (n=53 in Cameroon, n=252 in Senegal and n=50 in Tanzania) in this study. The pilot universities were chosen based on recommendations from the WHO and FIP, based on academic leadership and willingness to carry out the project. Two other universities who were also part of this pilot but did not yet deploy the course were excluded from the study. The students were at least in the third year of their study programme. The participation of the students in the evaluation was voluntary, and they were excluded from the analysis if they did not enter any information in the questionnaire.

Interviews with teachers

To understand how the course was deployed, a follow-up interview was conducted with the participating teachers from each university. These teachers were all directly involved in the implementation of the course after following a teaching (Train-the-Trainer) course themselves. A semistructured interview was used to assess enablers and barriers in deployment of the course and

Table 1 Characteristics of participating universities

Country	Teaching language	Total number students	Total hours	Deployment period	Pharmacy school year	Teachers (n)	Compulsory course	Standalone/part of another course
Cameroon	French	53	40	Nov 2020–Feb 2021	4th	4	Yes	Standalone
Senegal	French	252	16	Feb 2021–Apr 2021	3rd	3	No	Part of physico-chemical tests and drug control course
Tanzania	English	50	15	Jul 2021–Aug 2021	3rd	2	Yes	Part of quality assurance course

motivation of the students. An interview guide was developed with a list of ten open-ended questions. The list of the open-ended questions that were used for the semistructured interview and shared with the teachers beforehand in the interview guide can be found in the online supplemental appendix.

Student survey

In the pre-assessment, students were asked in an open question to write out their prior knowledge about SF medical products. The answers were coded based on their correspondence to the course's six modules.

In the post-assessment, students were asked (1) to what extent their knowledge on SF medical products had improved after participation in the course, (2) to indicate in which areas (linked to each of the six modules) their knowledge had improved, (3) to what extent this course will be helpful in their professional life and (4) if they would recommend this course to all pharmacy students and/or pharmacists. Answers were analysed by using descriptive analysis.

Knowledge assessment

No suitable questionnaire or assessment was found in the available literature that could be used for knowledge assessment in the context of this study. Therefore, an assessment questionnaire was developed and validated by the expert group who developed the course. The assessment comprised of 20 multiple choice questions related to the learning objectives of the curriculum. Each correctly answered question yielded one point. No points were deducted for incorrectly answered questions. Assessment responses from the students were received via computers or mobile devices with a web-based software tool, QuestionPro. None of the questions were mandatory. Fully uncompleted assessments, however, were excluded from the analysis.

The same assessment questions were used in the pre-test and post-test to ensure the compatibility of the results. The students were asked to fill out an assessment questionnaire before the course and after the course. The study approach is visualised in the online supplemental appendix.

Data analysis

The effects of the course on assessment scores were examined via linear mixed-effects model analysis. This method was chosen to account for both within-person and across-person variability. The statistical model contains both fixed effects and random effects. In addition, it is an appropriate method of analysis when not all data can be matched. Some participants only completed the pre-test or post-test leading to missing data, and linear mixed-effects model analysis allows to use all the data. One model was built with the assessment score as dependent variable and with fixed effects of time (pre-course or post-course). The country and subject number (student number) were considered as random effects. The validity of the statistical model was evaluated by using residuals to verify the normality assumption. The two-tailed significance level was set at $\alpha=0.05$. All statistical analyses were performed with IBM SPSS Statistics, V.25 and Microsoft Excel V.2016.

The questionnaire aimed to cover all modular areas, but the emphasis was not given to an equal distribution; therefore, some questions were linked to more modules than others. Module E on reporting focuses on a national situation and is therefore not suitable for a uniform questionnaire, thus was not part of the assessment. Initially, it was not the intention to assess improvement per module, but a post hoc descriptive analysis was undertaken to explore this. The assessment scores were stratified by module and the total obtained score (percentages) per module was determined, both before (pre-test) and after (post-test) the course. The percentages were determined by dividing the obtained total score per module over the theoretical maximum score (number of questions per module multiplied by the number of students who completed the assessment) multiplied by 100%.

Data from the student survey and interview responses from the semistructured interviews were clustered into four themes (enablers, barriers, motivation of students and important environmental factors) with important quotes highlighted.

Table 2 Deployment of SF course at different universities

Factors	Description	Quotes
Enablers	<ul style="list-style-type: none"> ▶ Participation in course development and ownership ▶ Support from WHO, FIP and other partners ▶ Adequate preparation of teachers with Train-the-Trainers training ▶ Availability of course materials; bilingually available materials ▶ Sufficient measures for alignment with existing pharmacy courses ▶ Course development coincided with overall University curriculum revision— support from university leadership ▶ With support from university leadership course is standalone and compulsory 	<ul style="list-style-type: none"> ▶ “Often when we prepare a lesson, we try to manage the documents with which we are going to prepare a training support, but here it is already prepared in advance, with updated data, excellent examples and very concrete cases. (...) We participated in the development providing our input and insights of what are likely to be the needs of our students.” ▶ “The project’s start and course development coincided with major undergraduate pharmacy curriculum revision at the university. (...) Because we were involved in the development of the new course from the beginning, we were able to take sufficient measures to ensure that the course aligned well with other obligatory pharmacy courses at our university.”
Barriers	<ul style="list-style-type: none"> ▶ Time constrains for course delivery; time constraints for pre-deployment preparation ▶ Lack of equipment for practical case studies, for example, provision of reference standards for Mini-Labs as university could not buy the reference tablets and equipment for practical case studies (eg, use of Global Pharma Health Fund Minilabs, detection tools, terrain work, etc) ▶ Lack of adequate teaching space (room size) ▶ Lack of internet access (and affordability), for example, some students had to buy internet passes to connect for the assessment ▶ Lack of support from university leadership to run the course as mandatory part of pharmacy curriculum ▶ Lack of support from the local authorities 	<ul style="list-style-type: none"> ▶ “The level of difficulty is in relation to the timing. (...) If we had the authorisation of the deanery to integrate the modules in the courses, it could facilitate the management of the timing, so that we would have an agenda with the necessary time to meet the objectives.” ▶ “We thought the course will take a shorter time, but once we started teaching, we realised we needed more time. When you are teaching, you are not simply reading the slides, you have to explain to the students, and give examples as it is a new topic for them, it was time more consuming.” ▶ “Students (...) need practicals. For example, using kits to demonstrate so that after graduation, they know what is about. We have a good experience of Minilab, we would have used it for this course but the reference standards have expired.” ▶ “Solvents we can easily replace but we are not able to buy the reference tablets anymore.”
Motivation of students	<ul style="list-style-type: none"> ▶ Lot of interest from the students in this topic, with main feedback of it being useful for their future professional life ▶ Students could relate the teaching material to media reports ▶ The classes were full 	<ul style="list-style-type: none"> ▶ “Students were very participative during the lectures.” ▶ “Students were very interested, (...) because it is directly related to the daily life as pharmacists.”
Important environmental factors	<ul style="list-style-type: none"> ▶ COVID-19-related restrictions, for example, social distance, students wearing masks in class ▶ Context of the revival of the pharmaceutical industry as part of stimulation of local production to achieve therapeutic sovereignty and consequently, pharmacy reforms are expected ▶ Overall university curriculum revision 	<ul style="list-style-type: none"> ▶ “We taught in the circumstances of COVID-19, for example, students had to wear masks.” ▶ “(...) there is a revival of the local pharmaceutical industry, which interests the students a lot. This course allows them to better communicate and inform the population.”

SF, substandard and falsified.

RESULTS

Pre-assessment versus post-assessment knowledge scores

A total number of 335 students (response rate 94.4%) filled out the questionnaire. Overall, as displayed in [table 2](#), student knowledge scores improved by 3.5 points (95% CI 3.1 to 3.9) out of 20 points after they took the course. When stratified by country, students showed a

statistically significant improvement of 3.5 points (95% CI 2.2 to 4.8) in Cameroon, 3.4 points (95% CI 3.0 to 3.8) in Senegal and 5.3 points (95% CI 2.9 to 7.7) in Tanzania ([figure 1](#)).

Students improved in each measured module across all countries ([figure 2](#)). In Senegal, the biggest improvement (15.6%) was observed in module C, covering




Figure 1 Assessment score pre-test versus post-test.

information about the prevention of SF medical products from entering the supply chain. In Cameroon, the largest improvements (14.0%) were observed in module D on detection and in Tanzania (43.2%) in module B related to the identification of medical products at risk.

Self-assessment by students

A total number of 246 students described their prior knowledge of SF medical products (Cameroon $n=51$, Senegal $n=167$, Tanzania $n=28$, response rate 69.3%). Among them, 36 students (14.6%) indicated no previous knowledge. If any knowledge was indicated, students in all three countries indicated that they mostly had some prior knowledge of the general aspects about SF medical products (52.6% in Senegal; 42.9% Cameroon; 60.0%

Tanzania). Prior knowledge per module per country can be found in the online supplemental appendix.

Students overall declared a similar improvement in all modules, approximately around 20% per module. The online supplemental appendix lists country-specific improvements per module. Most of the students indicated that their knowledge improved because of participating in the course (figure 3). Also, the large majority found the course useful for their professional life and would recommend the course to all pharmacy students and/or pharmacists around the globe. No large differences were observed between countries.

Interview sessions with teachers

Interviews with the teachers helped to understand the context and key enablers and barriers of course deployment. These are listed in table 2 with selected quotes included.

DISCUSSION

This jointly developed educational course on SF medical products for undergraduate pharmacy students improved their knowledge on these poor-quality medical products. Knowledge increased overall and for each individual module in all countries, both as assessed through the questionnaire and according to student self-assessment. The course was well received among students and faculty, and enablers and barriers of deployment were identified.




Figure 2 Percentages of questions answered correctly, stratified by modular theme in Cameroon, Senegal and Tanzania. 'Introduction' corresponds to module A and $n=5$ questions; 'Identification' corresponds to module B and $n=3$ questions; 'Prevention' corresponds to module C and $n=1$ questions; 'Detection' corresponds to module D and $n=4$ questions; 'Intervention' corresponds to module F and $n=7$ questions.




Figure 3 Students' attitudes towards knowledge improvement, future recommendation to others and usefulness of the course.

This may encourage further full implementation of this course in existing curricula beyond the pilot and inform possible future scale-up. It also indicates that prevention and detection of SF medical products should move from being a top-down exercise held by specialists to being integrated into the pharmaceutical system through training of practitioners in a bottom-up approach.

The course was building on rather poor (self-reported) prior knowledge of students. Nearly one-sixth of the responding students reported no previous knowledge, while others reported some knowledge, mainly related to general aspects about SF medical products. This does not come as a surprise given earlier observations from WHO and others.^{3 8 12} While of high importance, current pharmacy school curricula largely lack systematic teaching on SF medical products.¹³ When introducing a new topic in a healthcare curriculum, the implementation typically undergoes different phases, from the exploration phase where the topic is being explored via early-adopting programmes (such as this pilot programme) to the initiation phase during which adoption across healthcare educational programmes is enacted. This phase will require continuous feedback to adjust appropriate educational delivery mechanisms. After this follows a standardisation phase in which knowledge is spread throughout the profession.¹⁷

To inform such process, it is important to identify key barriers and enablers of course implementation.¹⁸ One of the key barriers observed was insufficient time. In all countries, teachers reported that they would welcome more hours to be either dedicated to this course or to the preparation that preceded the deployment. Insufficient time in pharmacy or medical-related curricula is a well-known barrier to a new course introduction.^{13 19-21} This barrier can be overcome by support from university leadership,¹² which was observed in Tanzania in the present study where the course development coincided with overall university curriculum revision, which resulted in more support for this course as well. Furthermore, the course would benefit from removing barriers of access to practical means, and more resources to provide for

equipment, sufficient teaching space, internet accessibility and affordability, etc.

On the enablers side, the readily available teaching materials were considered very helpful resources for the teachers to facilitate the delivery. As previous studies found, availability of ready-to-adopt course materials is an important enabler.¹³ Participation of universities in the course development, providing input and insights of what is likely to be the needs of their students and overall ownership and commitment in the pilot project was a crucial enabler of success as has been reported previously.²²

The course was developed as a modular one. When looking at individual themes, even though the universities eventually decided not to carry out the course per module and instead run the modules interchangeably, students improved in all measured modules in each country. The results therefore indicate the students improved in various areas of the course content. This was also reflected in the overall improvement across all modules in the student's self-assessment. This indicates that the course was effective in reaching its objective of increasing student's knowledge on SF medical products. Yet, this is not reason to stand still. As Dizon *et al* showed, even if students (momentarily) improve their knowledge through training, they still need to continue developing such knowledge and skills.²³ The importance of continuous (lifelong) learning is consistently highlighted.²⁴⁻²⁶ This calls for efforts to integrate training on SF medical products into continuous education for pharmacists.

Strengths and limitations

This is the first study bringing insight into the education on SF medical products and deployment of the comprehensive and dedicated undergraduate pharmacy course in sub-Saharan Africa, but there are several limitations to note. First, not all participating students completed both the pre-assessment and post-assessment questions due to practical issues. Therefore, a linear mixed-effects model was used to be able to include all the available data.

Second, the number of questions was not equally distributed across modules—balancing between optimising the length (and respective time burden on students) with a total of 20 questions only, and full representation of the course content. Moreover, each university had the possibility to adjust the teaching content to their needs. Therefore, knowledge questions related mainly to common aspects across modules.

Third, as the same assessment questions were used in the pre-test and post-test, students could theoretically have memorised the questions in favour of the post-test scores. However, the assessment answers were not shared after completing the pre-test, making this assumption less plausible. Fourth, although the course has led to a statistically significant increase in knowledge, it is uncertain how sustainable the knowledge acquired is. Also, it is unclear how long this increase will remain, which is subject for further research. It was also not the aim of the study to compare the countries given the differences in the implementation context. Although there were differences across schools, it is difficult to draw any conclusions.

Despite the limitations cited, the study presents supporting evidence for the success in this pilot course implementation with the aim to close the gap in SF medical products education.

Future directions

The findings of this study can inform future pilots and institutionalisation of the training. As soon as the two other universities who were initially involved at the pilot project but have not yet introduced the course (Nigeria and Uganda) will deploy the course, findings can be compared across the whole pilot. This would also allow to investigate further the bilingual (francophone and anglophone) and cultural aspect of the curriculum, as this has been a unique feature of this course. The results could be compared against insight studies on awareness/risks/access in the similar socioeconomic segments of the general population in each country. Recent studies conducted by the WHO in Uganda and Nigeria can serve as a starting point.

Future insights could be given into long-term knowledge acquisition (namely, whether the score would change in 2–5 years). Furthermore, it could be investigated whether the students used the acquired knowledge in their practice-oriented professional careers. Continuous professional development for the (wider) workforce also needs consideration.

While this course was implemented in sub-Saharan Africa, clear and consistent themes emerged that can be applied in expanding this course to other geographical regions—a step towards providing pharmacists with the knowledge to contribute to minimising of the threat to the populations around the globe. Adaptations of this course (language, scope, etc) could then be deemed elsewhere.

CONCLUSIONS

The undergraduate pharmacy students who participated in the dedicated educational course improved their knowledge of SF medical products. Knowledge improved in all countries across all measured modules, both as assessed through an assessment and according to student self-assessment. The course was well received among students and teachers, who found the course useful and would recommend the course further. Enablers and barriers to deployment were identified, which can inform further full implementation of this course in existing curricula beyond the pilot and possible future scale-up to other pharmacy schools. Empowering all pharmacists across the health systems—instead of specialist training only—to address the socially and ethically unacceptable issue of SF medical products has a potential for reinforcing health systems' capacity and safeguarding public health.

Author affiliations

- ¹Utrecht Centre for Pharmaceutical Policy and Regulation, Division of Pharmacoepidemiology and Clinical Pharmacology, Utrecht Institute for Pharmaceutical Sciences (UIPS), Utrecht University, Utrecht, The Netherlands
²International Pharmaceutical Federation, The Hague, The Netherlands
³University of Douala, Douala, Cameroon
⁴Muhimbili University of Health and Allied Sciences, Dar es Salaam, Tanzania, United Republic of
⁵Universite Cheikh Anta Diop de Dakar, Dakar, Senegal
⁶Access to medicines and health products, World Health Organization, Geneva, Switzerland

Twitter Zuzana Kusynová @ZKusynova and Eliangiringa Kaale @eliangiringa1

Acknowledgements The authors would like to thank Dr Ir Irma Meijerman, Utrecht Institute for Pharmaceutical Sciences (UIPS), Utrecht University, Utrecht, The Netherlands, for her input to study design, Ms Svetlana Belitser, statistician at the Utrecht Institute for Pharmaceutical Sciences (UIPS), Utrecht University, Utrecht, The Netherlands, for her insights in the statistical analysis and Dr Catherine Duggan, the CEO of the International Pharmaceutical Federation (FIP) for the final review of this article. Given this research was conducted from international partnerships, a structured authors' reflexivity statement was added in a supplementary appendix to the paper. It summarises the measures authors put in place in their efforts to promote equitable authorship in this publication.

Contributors All authors were involved in conceptualising the study. ZK and YB collected and analysed the data with the local help of GE-L, EK and SOS. ZK and YB drafted the manuscript and all authors reviewed the paper. AM-T served as the overall guarantor for the study.

Funding The development and implementation of the course (six university modules based on a competency framework) was financially supported by the European Commission and the WHO.

Disclaimer The views expressed in this article are the personal views of the authors and must not be understood or quoted as being made on behalf of any of the affiliated organisations.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval The study was conducted under Dutch legislation that indicates ethical approval is not required for this type of study. The study was conducted according to the principles of the Directive 95/46/EC General Data Protection Regulation (GDPR). Informed, specific and explicit consent was freely given by the students who participated in the study. Furthermore, informed, specific and explicit consent was freely given by the teachers who were interviewed, and the interview guide was shared at least one week beforehand. Participation by students and teachers was completely voluntary and could be terminated at any moment.

The data were handled anonymously and none of the findings is traceable to the individual participant. Data are stored on a secure location and only accessible to the core research team. Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as supplementary information. All data relevant to the study are included in the article or uploaded as supplementary information. Contact the corresponding author if you wish to receive more information.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.


REFERENCES

- Nayyar GML, Breman JG, Mackey TK, *et al*. Falsified and substandard drugs: stopping the pandemic. *Am J Trop Med Hyg* 2019;100:1058–65.
- Barry J. Fake medicines: a global threat. *Nurs Manag* 2014;21:17.
- World Health Organization (WHO). Substandard and falsified medical products: scope, scale and harm. Available: <https://www.who.int/news-room/questions-and-answers/item/substandard-and-falsified-medical-products-scope-scale-and-harm> [Accessed 09 Mar 2022].
- Venhuis BJ, Keizers PHJ, Klausmann R, *et al*. Operation resistance: a snapshot of falsified antibiotics and biopharmaceutical injectables in Europe. *Drug Test Anal* 2016;8:398–401.
- Janvier S, De Spiegeleer B, Vanhee C, *et al*. Falsification of biotechnology drugs: current dangers and/or future disasters? *J Pharm Biomed Anal* 2018;161:175–91.
- Mori AT, Meena E, Kaale EA. Economic cost of substandard and falsified human medicines and cosmetics with banned ingredients in Tanzania from 2005 to 2015: a retrospective review of data from the regulatory authority. *BMJ Open* 2018;8:e021825.
- Ozawa S, Evans DR, Bessias S, *et al*. Prevalence and estimated economic burden of substandard and falsified medicines in low- and middle-income countries: a systematic review and meta-analysis. *JAMA Netw Open* 2018;1:e181662.
- World Health Organization (WHO). WHO global surveillance and monitoring system for substandard and falsified medical products <https://apo.who.int/publications/i/item/978-92-4-151342-5>
- Newton PN, Green MD, Fernández FM. Impact of poor-quality medicines in the ‘developing’ world. *Trends Pharmacol Sci* 2010;31:99–101.
- World Health Organization (WHO). Substandard and falsified medical products, 2022. Available: https://www.who.int/health-topics/substandard-and-falsified-medical-products#tab=tab_1 [Accessed 09 Mar 2022].
- Renschler JP, Walters KM, Newton PN, *et al*. Estimated under-five deaths associated with poor-quality antimalarials in sub-Saharan Africa. *Am J Trop Med Hyg* 2015;92:119–26.
- Ferrario A, Orubu ESF, Adeyeye MC, *et al*. The need for comprehensive and multidisciplinary training in substandard and falsified medicines for pharmacists. *BMJ Global Health* 2019;4:e001681.
- Kusynová Z, Kicken M, Van den Ham HA, *et al*. Identifying the teaching content on substandard and falsified medical products in global pharmacy education as critical public health issue. *Pharmacy Education* 2021;21:504–16.
- World Health Organization (WHO). WHO member state mechanism, 2022. Available: <https://www.who.int/teams/regulation-prequalification/incidents-and-SF/mechanism> [Accessed 09 Mar 2022].
- International Pharmaceutical Federation (FIP). Curriculum for pharmacy students on substandard and falsified medicines. Available: <https://www.fip.org/file/4917> [Accessed 09 Mar 2022].
- World Health Organization (WHO). *Report on the tenth meeting of the member state mechanism on substandard and falsified medical products*. Geneva, 2021. https://apps.who.int/gb/sf/pdf_files/MSM10/A_MSM10_11-en.pdf
- Aungst TD, Patel R. Integrating digital health into the Curriculum—Considerations on the current landscape and future developments. *J Med Educ Curric Dev* 2020;7:238212051990127.
- Leithwood KA. Managing the implementation of curriculum innovations. *Knowledge* 1981;2:341–60.
- Aiyer M, Hemmer P, Meyer L, *et al*. Evidence-Based medicine in internal medicine clerkships: a national survey. *South Med J* 2002;95:1389–95.
- Meats E, Heneghan C, Crilly M, *et al*. Evidence-based medicine teaching in UK medical schools. *Med Teach* 2009;31:332–7.
- Blanco MA, Capello CF, Dorsch JL, *et al*. A survey study of evidence-based medicine training in US and Canadian medical schools. *J Med Libr Assoc* 2014;102:160–8.
- Kirk D, MacDonald D. Teacher voice and ownership of curriculum change. *Journal of Curriculum Studies* 2001;33:551–67.
- Dizon JMR, Grimmer-Somers KA, Kumar S. Current evidence on evidence-based practice training in allied health: a systematic review of the literature. *JBI Evidence Implementation* 2012;10:347–60.
- Collins J. Lifelong learning in the 21st century and beyond. *Radiographics* 2009;29:613–22.
- Duyff RL. The value of lifelong learning: key element in professional career development. *J Am Diet Assoc* 1999;99:538–43.
- Ei Mawas N, Muntean C H. *Supporting lifelong learning through development of 21 ST century skills. 10th International Conference on education and new learning technologies*. Palma de Mallorca, Spain, 2018.

SUPPLEMENTARY APPENDIX

Visualization of the timeline of the methodology

Figure 1. Visualization of the timeline of the methodology



Prior knowledge of SF products, self-assessed by students

A total number of 246 students filled out this open question (Cameroon n=51, Senegal n=167, Tanzania n=28, response rate 69.3% from 355 total participants) to describe their prior knowledge of SF medical products. There were no students who completed only part of the assessment, the assessment was either fully completed or not completed. A total number of 36 students (14.6% from total of 246 students who filled out the open question in the survey) indicated no previous knowledge (Cameroon n=18, Senegal n=13, Tanzania n=5). If any knowledge was indicated, students (n=144, 58.5% from total of 246 students who filled out the open question in the survey) in all three countries indicated that they mostly had some prior knowledge of the general aspects about SF medical products (Cameroon n=24, Senegal n=102, Tanzania n=18). No students in Cameroon and Tanzania indicated any prior knowledge about identification of medical products most at risk being falsified (module B), reporting SF medical products (module E) and counseling of patients exposed to falsified medical products (module F). The latter two were similar in Senegal, with no student indicating prior knowledge in these.

Pre-test: The distribution of the coded answers is presented in Table 1 and Figure 2.


Table 1. Distribution of coded responses to the question about prior knowledge of SF medication as self-assessed by students, stratified by country

	Percentage (%) of responses		
	Cameroon	Senegal	Tanzania
Introducing the problem /general information about SF medical products (module A)	42.9	52.6	60.0

Identification (module B)	0.0	2.1	0.0
Prevention (module C)	3.6	8.8	10.0
Detection (module D)	10.7	11.9	13.3
Report (module E)	0.0	0.0	0.0
Advisement (module F)	0.0	0.0	0.0
Nothing/no prior knowledge	32.1	6.7	16.7
Knowledge about other courses	10.7	18.0	0.0

Caption: The sum of the percentages as presented may not be 100 due to rounding.

Figure 2. Distribution of coded responses (%) on question about prior knowledge among students from Cameroon, Senegal and Tanzania



The number of responses is higher compared to the number of students who completed this open question, because multiple modules (i.e., codes) could correlate to the given answer.

Areas of improvement: self-assessment

A total number of 281 students (response rate 79.2% from total of 355 participants) answered the question on self-assessment per module (Cameroon: $n_{\text{students}}=8$, $n_{\text{responses}}=36$; Senegal $n_{\text{students}}=215$, $n_{\text{responses}}=998$; Tanzania $n_{\text{students}}=57$, $n_{\text{responses}}=291$). Students overall declared a similar improvement in all modules, approximately around 20% per module.

Specifically, students from Senegal indicated that they improved the most in module A ($n_{\text{responses}}=197$; 19.7% out of 998), which gives a general introduction to SF medical products. The students indicated that they improved the least in module F ($n_{\text{responses}}=111$; 11.1% out of 998), which is about interventions after encountering SF medical products.

In Cameroon, a different distribution was observed, with students reporting an equal amount of knowledge improvement related to the content of modules A, B and C ($n_{\text{responses}}=8$; 22.2% each out of 36). For module D, E and F students reported less knowledge improvement ($n_{\text{responses}}=6$; 16.7%, $n_{\text{responses}}=2$; 5.6% and $n_{\text{responses}}=4$; 11.1%, respectively, thus, n responses divided over total of 36 responses).


In Tanzania, students indicated that they improved the most in module B ($n_{\text{responses}}=52$; 17.9% out of 291), which is about identification of medical products at risk. Comparable to the results in Senegal, the students from Tanzanian students self-assessed that they improved the least in module F ($n_{\text{responses}}=44$; 15.1% out of 291). Table 2 and Figure 3 summarizes the distribution.

Table 2. Distribution of responses to the question about gained knowledge of SF medical products as self-assessed by students, stratified by country

	Percentage (%) of responses		
	Cameroon	Senegal	Tanzania
Introducing the problem of SF medical products/general information (module A)	22.2	19.7	17.5
Identification (module B)	22.2	17.3	17.9
Prevention (module C)	22.2	16.8	16.5
Detection (module D)	16.7	17.9	16.5
Report (module E)	5.6	17.0	16.5
Advisement (module F)	11.1	11.1	15.1

Caption: The sum of the percentages as presented may not be 100 due to rounding.

Figure 3. Distribution of responses to the question about gained knowledge of SF medical products as self-assessed by students, stratified by country



The assessment questionnaire

The assessment questionnaire can be requested by contacting the corresponding author on confidential basis. Given the programme is ongoing in other African universities, it is important that students who may read the published article are not aware in advance of the questions they will be asked.

The list of open-ended questions used in the semi-structured interview

The following open-ended questions were used for the semi-structured interview and shared with the teachers beforehand in the interview guide:

Question 1 – Preparation

Were you adequately prepared for the teaching of the course (1) and what feedback would you give for improvement of the preparation? (2)

What tools were helpful in your teaching preparation?

- Was the Train the Trainers useful in your preparation?
- Was the Curriculum Guide useful in your preparation?
- Was the Competency Framework useful in your preparation?
- Was the Moodle platform useful in your preparation?

Question 2 – Time investment

How long did the course last?

- Did you think this was enough time to properly teach all the modules?
- If not, which modules/parts needed more time?

Question 3 - Time path course

Have you been able to complete the entire course (in accordance with the predefined learning objectives)?

- What modules (Modules A-F) did you deliver?
- If not, what were the reasons for this? Which parts have been excluded and which considerations played a role in this?

Question 4 – Obligations

Was it a compulsory course or an elective course for the students?

- Are the modules of the course part of another course or was it a stand-alone course?
- Was there an attendance requirement for the students? If not, can you indicate something about the degree of presence (compared to other (obligatory) courses)?

Question 5 – Quality of the course

What feedback would you give for improvement of the course?

Do you think there are any differences in the quality of the SF medicines course compared to other (longer existing) Pharmacy courses?

- Is the course set up in a similar way to other Pharmacy courses (teaching methods used, teaching material used, class size, level of difficulty etc.)?
- Do you think the students experienced the teaching material as challenging (enough)?

Question 6 – Motivation among students

Do you think that the students were motivated for the course?

- What impression did the student have on you?
- What do you think can help students to become more motivated to take the course (1) and to complete the assessment (2)?

Question 7 – Limitations

Did you lack any materials, support or knowledge in the implementation of the course (1) or during the evaluation phase (2) (when students took the assessment survey)?

Question 8 – Completion of the course

How is the course completed?

- In case of a compulsory final exam, what form of testing was used? Can the teachers share some information about average mark and/or pass percentage?
- In case of a compulsory final exam, did the teacher think the students were good enough informed about the learning objectives of the course?
- In other cases, why did the university choose not to have a compulsory final exam for this course?

Question 9 – Unforeseen circumstances

Are there any specific circumstances that may have influenced the education and/or implementation of the course?

- How does the COVID-19 situation influence student activities at your university?
- Any political, economic, social developments in the region?
- Any technical issues (Internet connection, face-to-face access to University, etc.)

Question 10 – Future perspectives

Would you recommend the course to other universities/countries around the globe?

- What tip/advice would you give other universities/teachers if they also strive for a successful implementation of the SF medicine curriculum?

Structured authors' reflexivity statement

Given this research was conducted from international partnerships, a structured authors' reflexivity statement summarizes the measures authors put in place in their efforts to promote equitable authorship in this publication.

Table 3: Structured authors' reflexivity statement

Study conceptualisation	<p>1. How does this study address local research and policy priorities?</p> <p>The main objective of this study was to assess the change in knowledge regarding substandard and falsified (SF) medical products as understood by undergraduate pharmacy students. The paper focuses on the course delivered in three selected pilot pharmacy schools in Sub-Saharan Africa (Cameroon, Senegal and Tanzania). This topic is of great importance in these countries given it is a major public health issue.</p> <p>2. How were local researchers involved in study design?</p>
--------------------------------	---

	<p>The protocol for the study design was developed in collaboration with local university teachers / professors (from LMIC) who were responsible for the deployment of the courses in their universities. Multiple meetings were held to discuss the design of the content, progress of the course, and status of the research project.</p>
<p>Research management</p>	<p>3. How has funding been used to support the local research team(s)?</p> <p>The local researchers from LMIC were the same university teachers who helped to design and implement the course. They are also staff members receiving salary from their universities. The funding benefitted them by providing them with material to use in their daily practice (e.g., the guide for teachers). The funding was invested to build their capacity and deployment capability (e.g., though Train-the-Trainer course).</p>
<p>Data acquisition and analysis</p>	<p>4. How are research staff who conducted data collection acknowledged?</p> <p>These are all co-authors of the paper that resulted from the research and are acknowledged in the curriculum guide that was developed to also implement the course.</p> <p>5. How have members of the research partnership been provided with access to study data?</p> <p>Involved local university teachers from LMIC are all experts within the field of SF medical products and those responsible within their respective universities for the deployment of the course within the pharmacy curriculum. They have been involved in conceptualising the study, collecting the data, evaluating the results and reviewing multiple versions of the manuscript. The research team relied heavily on their interpretation of the data and providing the context to the results. All local university teachers were included as co-authors in the manuscript. For ethical reasons and protection of personal information, the pooled study data was only accessible to the core team.</p>

	<p>6. How were data used to develop analytical skills within the partnership?</p> <p>The local university teachers from LMIC were involved in conceptualising the study and provided input at different stages. The data analysis was done by the first and the second author with supervision and help of senior university professors. A statistician was onboarded to ensure correct performance of the analysis. Results were presented to all co-authors and critically discussed and reviewed.</p>
Data interpretation	<p>7. How have research partners collaborated in interpreting study data?</p> <p>Results were reviewed by all co-authors and interpretation of the results was done using the local context, with valuable input from the local university teachers.</p>
Drafting and revising for intellectual content	<p>8. How were research partners supported to develop writing skills?</p> <p>The first author drafted the paper, and it was critically reviewed by all co-authors in multiple rounds.</p> <p>9. How will research products be shared to address local needs?</p> <p>Results from this study will be used to improve subsequent editions of the course within the universities and serve as guide for other universities to “adapt and adopt” for implementation. The programme is intended to run continuously in the future.</p>
Authorship	<p>10. How is the leadership, contribution and ownership of this work by LMIC researchers recognised within the authorship?</p> <p>Involved university teachers from LMIC are all experts within the field of SF medical products and those responsible within their respective universities for the deployment of the course within the pharmacy curriculum. They have been involved in</p>

	<p>conceptualizing the study, collecting the data, evaluating the results and reviewing multiple versions of the manuscript. The team relied heavily on their interpretation of the data and providing the context to the results. All researchers were included as co-authors in the manuscript.</p> <p>11. How have early career researchers across the partnership been included within the authorship team?</p> <p>The first and the second author of this paper are early career researchers, a PhD student and a master student respectively.</p> <p>12. How has gender balance been addressed within the authorship?</p> <p>There is an ensured gender balance with co-authors being both female and male, with majority of females (7 out of total 9 co-authors).</p>
Training	<p>13. How has the project contributed to training of LMIC researchers?</p> <p>This project trained the involved LMIC teachers in educational research, which provides evidence of the impact of new courses within the pharmacy curriculum. Scientific advice on educational research was obtained from a well-known expert at Utrecht University who was acknowledged in the relevant section of the paper. It should be noted that in the context of this project, the research partners based in LMIC gave guidance to the authors of this paper (i.e. researcher training was bi-directional) in particular to help adapt, contextualise, and interpret several elements.</p>
Infrastructure	<p>14. How has the project contributed to improvements in local infrastructure?</p> <p>The project demonstrated the need for specific demonstration equipment in universities - especially for the module on detection, and namely screening devices. This research and publication will be used as evidence for WHO and other partners to supply field screening devices to the pilot universities, so that the students can</p>

	apply the theoretical knowledge and practice with adequate equipment.
Governance	<p>15. What safeguarding procedures were used to protect local study participants and researchers?</p> <p>Study participants are students and evaluations are expected as part of the curriculum as for any course. Ethical advice was sought at Utrecht University. Informed consent was given by study participants and data are presented anonymously where possible. No personal data was used in the research and participants were able to withdraw their participation at any point in time.</p>