

HIV and SARS-CoV-2: the interplay of two wicked problems

Anneleen Kiekens ¹, George Msema Bwire,^{1,2} Catherine Decouttere,³ Michael R Jordan,^{4,5} Ally Mangara,⁶ Idda H Moshia,⁷ Tobias Rinke de Wit,⁸ Raphael Zozimus Sangeda ², Omary Swalehe,⁹ Nico Vandaele,³ Japhet Killewo,¹⁰ Anne-Mieke Vandamme ^{1,11}

To cite: Kiekens A, Bwire GM, Decouttere C, *et al.* HIV and SARS-CoV-2: the interplay of two wicked problems. *BMJ Global Health* 2022;**7**:e009105. doi:10.1136/bmjgh-2022-009105

Handling editor Seye Abimbola

Received 17 March 2022

Accepted 29 May 2022

Concern is justified observing the link between the AIDS and COVID-19 pandemics. COVID-19 outcomes are significantly worse in many people living with HIV (PLHIV), even when vaccinated, because of their impaired immune system. Moreover, CD4 T-cells are affected by both HIV and SARS-CoV-2.¹⁻³ SARS-CoV-2 variants can evolve in immunosuppressed patients due to prolonged viral replication in the context of an inadequate immune response.⁴ Accelerated intrahost evolution of SARS-CoV-2 was reported in a South African HIV patient with antiretroviral therapy (ART) failure.^{5,6} With 25 million HIV patients in sub-Saharan Africa (SSA) of whom an estimated 8 million are not virologically suppressed, this potentially creates a reservoir for future variants. Such variants, arising in PLHIV anywhere in the world, can spread to other continents, as has been reported for variants of concern (VoCs) (Beta, Omicron) and variants of interest (B.1.6.20, B.1.640.2) that arose in Africa.⁷⁻⁹ Conversely, the COVID-19 pandemic impacts HIV treatment programmes, due to supply chain issues, overburdening of healthcare systems, limiting access to testing, treatment and prevention programmes and further increasing inequalities.¹⁰ Modelled COVID-19 disruptions of HIV programmes in SSA included decreased functionality of HIV prevention programmes, HIV testing and treatment, healthcare services such as viral load testing, adherence counselling, drug regimen switches and ART interruptions, which may lead to selection of drug-resistant HIV.¹¹ A 6-month interruption affecting 50% of the population would lead to a median number of excess deaths of 296 000, during 1 year. Scientists advocate for the AIDS and COVID-19 pandemics in Africa to be addressed simultaneously,

Summary box

- ⇒ The AIDS and COVID-19 pandemics are intersecting at multiple levels.
- ⇒ Both pandemics have previously been described as wicked problems and can be studied as complex adaptive systems.
- ⇒ We recently designed a qualitative model of all known factors influencing HIV drug resistance (HIVDR) in sub-Saharan Africa and analysed its complexity as a complex adaptive system. Our detailed systems map featured three main feedback loops driving HIVDR.
- ⇒ Visualising both pandemics in one systems map confirms the urgency to address both simultaneously in order to mitigate a prolongation of the current COVID-19 pandemic and a rise in drug-resistant HIV.
- ⇒ Also at local level, a visualisation of the effects of the COVID-19 pandemic and response measures on the HIVDR system is valuable in gaining an understanding of its system-wide impact when rapid public health action is needed.

by increasing African access to COVID-19 vaccines, prioritising research on the interaction between HIV care and COVID-19, maintaining high-quality HIV services and integrating health services for both viruses.⁷ Both the COVID-19 and the AIDS pandemic, more specifically the issue of HIV drug resistance (HIVDR), have previously been described as wicked problems which are best studied as complex adaptive systems (CASs).¹²⁻¹⁵ Wicked problems consist of diverse interconnected factors and require complexity-informed and locally adapted solutions rather than one solution that fits all. We recently designed a qualitative model of all known factors influencing HIVDR in SSA and analysed its complexity.¹³ Our detailed systems map featured three main feedback loops driving HIVDR, representing (1) the alternation between adherence and non-adherence,



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Anne-Mieke Vandamme; annemie.vandamme@kuleuven.be




Figure 1 Interlinkage of the HIV and COVID-19 CAS at the international level, as supported by literature (see main text). The blue, pink and purple loops are feedback loops identified in the HIVDR CAS.¹³ The yellow elements are related to the COVID-19 pandemic and illustrate the interlinkage between both complex problems. Other links between the COVID-19 on the HIVDR CAS are to be expected, such as the likely increase in transmission of HIVDR due to increased sexual violence during lockdown.²¹ The 60% increase in teenage pregnancies in South Africa during the pandemic points in the same direction.²² ART, antiretroviral therapy; CAS, complex adaptive system; HIVDR, HIV drug resistance.

(2) the impact of an overburdened healthcare system and (3) the importance of sustaining global efforts of tackling HIVDR even when new antiretroviral drugs with high genetic barriers become available. These HIV-related feedback loops are interconnected with COVID-19 pandemic impact (in yellow, figure 1). The loop starts from PLHIV with an unsuppressed viral load, which weakens the immune system and may in turn slow down immune clearance of SARS-CoV-2, allowing prolonged replication and mutation of the virus in the context of an inadequate immune response. Prolonged viral clearance facilitates the selection of immune escape SARS-CoV-2 variants. Variants may emerge that have a selective advantage and therefore may spread through populations due to increased transmissibility (with possibly increased virulence), thereby creating an additional burden on the healthcare system, putting the overall healthcare system and the HIV care at risk. These stressors on the healthcare system lead to a higher risk of unsuppressed viral load in PLHIV, increasing the risk of HIVDR. Figure 1 shows the need to address both wicked problems simultaneously and to do so in a complexity-informed manner

as they are inevitably linked and influence each other. Evidently, the exact interconnections between both pandemics need to be locally assessed. For instance, a study in South Africa showed that while lockdown severely impacted HIV testing and ART initiation, ART provision was largely maintained, indicating that the strength of the connection between the virological suppression-related loop and the pandemic, indicated in figure 1, are context-dependent.¹⁶

At a local level we can visualise the impact of the COVID-19 pandemic and measures impacting HIV care and the lives of PLHIV. We did so based on a systems map of factors influencing HIVDR in the Ukonga and Gongolamboto areas of Dar es Salaam region, Tanzania.¹⁴ In the period between March and June 2020 several COVID-19 related restrictions were implemented in Tanzania. These included banning social gatherings, avoiding non-essential travel, obligatory face masks at health facilities, and the provision of 6-month ART for all PLHIV, instead of the usual programmatic recommendation of 1-month and 3-month supplies for patients without and with recent viral suppression, respectively.^{17 18} In figure 2,

⁵Division of Geographic Medicine and Infectious Diseases, Tufts Medical Center, Boston, Massachusetts, USA

⁶Dar es Salaam Urban Cohort Study, Dar es Salaam, Tanzania, United Republic of

⁷Department of Behavioural Sciences, Muhimbili University of Health and Allied Sciences, Dar es Salaam, Tanzania, United Republic of

⁸Amsterdam Institute for Global Health and Development (AIGHD), Amsterdam, The Netherlands

⁹Department of Business Studies, Mzumbe University, Dar es Salaam, Tanzania, United Republic of

¹⁰Department of Epidemiology and Biostatistics, Muhimbili University of Health and Allied Sciences, Dar es Salaam, Tanzania, United Republic of

¹¹Center for Global Health and Tropical Medicine, Instituto de Higiene e Medicina Tropical, Universidade Nova de Lisboa, Lisboa, Portugal

Twitter Raphael Zozimus Sangeda @rsangeda

Contributors Conceptualisation, AK, A-MV, IHM, GMB, AM, CD, NV, OS, RZS, MRJ, TRdW and JK; writing—original draft preparation: AK; writing—review and editing, AK, IHM, GMB, AM, NV, CD, OS, RZS, JK, MRJ, TRdW and A-MV; visualisation, AK; Supervision, A-MV, NV, JK and RZS; Funding Acquisition, A-MV, AK, RZS and JK.

Funding This work was funded by VLIR-UOS, grant numbers TZ2019SIN263 and TZ2020J0I032A101.

Competing interests A-MV declares consultancy fees from Gilead.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement No data are available.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Anneleen Kiekens <http://orcid.org/0000-0001-5083-9144>

Raphael Zozimus Sangeda <http://orcid.org/0000-0002-6574-5308>

Anne-Mieke Vandamme <http://orcid.org/0000-0002-6594-2766>

REFERENCES

- Spinelli MA. SARS-CoV-2 vaccination in people with HIV. *Lancet HIV* 2020;19–21.
- Bhaskaran K, Rentsch CT, MacKenna B, et al. HIV infection and COVID-19 death: a population-based cohort analysis of UK primary care data and linked national death registrations within the OpenSAFELY platform. *Lancet HIV* 2021;8:e24–32.
- Peng X, Ouyang J, Isnard S, et al. Sharing CD4+ T cell loss: when COVID-19 and HIV collide on immune system. *Front Immunol* 2020;11:1–11.
- Corey L, Beyrer C, Cohen MS, et al. SARS-CoV-2 variants in patients with immunosuppression. *N Engl J Med* 2021;385:562–6.
- Cele S, Karim F, Lustig G, et al. SARS-CoV-2 prolonged infection during advanced HIV disease evolves extensive immune escape. *Cell Host Microbe* 2022;30:154–62.
- Karim F, Moosa M, Gosnell B, et al. Persistent SARS-CoV-2 infection and Intra-Host evolution in association with advanced HIV infection. *medRxiv* 2021:1–13.
- Msomu N, Lessells R, Misana K, et al. Africa: tackle HIV and COVID-19 together. *Nature* 2021;600:33–6.
- Dudas G, Hong SL, Potter B, et al. Travel-Driven emergence and spread of SARS-CoV-2 lineage B. 1. 620 with multiple VOC-like mutations and deletions in Europe. *MedRxiv* 2021.
- Colson P, Delecre J, Burel E, et al. Emergence in southern France of a new SARS-CoV-2 variant of probably Cameroonian origin harbouring both substitutions N501Y and E484K in the spike protein. *medRxiv* 2021.
- The global fund results report 2021. ReliefWeb. Available: <https://reliefweb.int/report/world/global-fund-results-report-2021>
- Jewell BL, Mudimu E, Stover J, et al. Potential effects of disruption to HIV programmes in Sub-Saharan Africa caused by COVID-19: results from multiple mathematical models. *Lancet HIV* 2020;7:e629–40.
- Angeli F, Montefusco A. Sensemaking and learning during the Covid-19 pandemic: a complex adaptive systems perspective on policy decision-making. *World Dev* 2020;136:105106.
- Kiekens A, Dierckx de Casterlé B, Pellizzer G, et al. Exploring the mechanisms behind HIV drug resistance in sub-Saharan Africa: conceptual mapping of a complex adaptive system based on multi-disciplinary expert insights. *BMC Public Health* 2022;22:455.
- Kiekens A, Moshia IH, Zlatić L, et al. Factors associated with HIV drug resistance in Dar ES Salaam, Tanzania: analysis of a complex adaptive system. *Pathogens* 2021;10:1535.
- Klaschke B. After COVID-19: what can we learn about wicked problem governance? *Soc Sci Humanit Open* 2021;4:100173.
- Dorward J, Khubone T, Gate K, et al. The impact of the COVID-19 lockdown on HIV care in 65 South African primary care clinics: an interrupted time series analysis. *Lancet HIV* 2021;8:e158–65.
- Tanzania bans all public gatherings, closes school, suspends the premier League over coronavirus, 2020. The Citizen. Available: <https://www.thecitizen.co.tz/tanzania/news/national/tanzania-bans-all-public-gathering-closes-schools-suspends-the-premier-league-over-coronavirus-2705760>
- Bajaria S, Abdul R. Preparedness of health facilities providing HIV services during COVID-19 pandemic and assessment of their compliance to COVID-19 prevention measures: findings from the Tanzania service provision assessment (SPA) survey. *Pan Afr Med J* 2020;37:18.
- Wilkinson L, Grimsrud A. The time is now: expedited HIV differentiated service delivery during the COVID-19 pandemic. *J Int AIDS Soc* 2020;23.
- Hoffman RM, Moyo C, Balakasi KT, et al. Multimonth dispensing of up to 6 months of antiretroviral therapy in Malawi and Zambia (INTERVAL): a cluster-randomised, non-blinded, non-inferiority trial. *Lancet Glob Health* 2021;9:e628–38.
- Fawole OI, Okedare OO, Reed E. Home was not a safe haven: women's experiences of intimate partner violence during the COVID-19 lockdown in Nigeria. *BMC Womens Health* 2021;21:1–7.
- Teen pregnancies in South Africa jump 60% during COVID-19 pandemic, 2021. ReliefWeb. Available: <https://reliefweb.int/report/south-africa/teen-pregnancies-south-africa-jump-60-during-covid-19-pandemic>