




Service disruptions, trauma and burnout during the COVID-19 pandemic among healthcare providers delivering immunisation and maternal and child health services in Indonesia

Madeleine Randell ¹, Tri Yunis Miko Wahyono,² Michelle Dynes,³ Adeline Tinessia,¹ Mu Li,¹ Margie Danchin,^{4,5} Oktarinda,² Fitriyani Fitriyani,² Lintang Dian Saraswati,⁶ Kylie Jenkins,¹ Khin Devi Aung,⁷ Abdul Khalil Noorzad,⁷ Mrunal Shetye,⁷ Lulu Dewi,⁸ Prima Yosephine,⁸ Julie Leask ¹, Meru Sheel ¹

To cite: Randell M, Wahyono TYM, Dynes M, *et al*. Service disruptions, trauma and burnout during the COVID-19 pandemic among healthcare providers delivering immunisation and maternal and child health services in Indonesia. *BMJ Glob Health* 2024;**9**:e014318. doi:10.1136/bmjgh-2023-014318

Handling editor Seema Biswas

► Additional supplemental material is published online only. To view, please visit the journal online (<https://doi.org/10.1136/bmjgh-2023-014318>).

Received 23 October 2023
Accepted 7 June 2024



© Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Meru Sheel;
meru.sheel@sydney.edu.au

ABSTRACT

Background The COVID-19 pandemic resulted in extreme strain on health systems including the health workforce, essential health services and vaccination coverage. We examined disruptions to immunisation and maternal and child health (MCH) services, concerns of personal well-being and delivery of healthcare during the pandemic as well as factors associated with self-reported trauma or burnout among healthcare providers (HCPs).

Methods In March–April 2022, we conducted a cross-sectional survey among HCPs in two provinces of Indonesia. HCPs involved in COVID-19 or routine immunisation and MCH services were randomly selected from district/city health office registration lists. We descriptively analysed service disruptions experienced by HCPs as well as trauma, burnout and concerns of personal well-being and delivery of healthcare during the pandemic. Multivariate logistic regression analyses were undertaken to identify factors associated with trauma or burnout.

Results We recruited 604 HCPs. Mobilisation of staff from routine health services to COVID-19 response duties was a key reason for service disruptions (87.9%). Strategies such as community outreach and task shifting were implemented to overcome disruptions. Trauma or burnout during the pandemic was reported by 64.1% HCPs, with 23.5% reporting worse mental or emotional health. Factors associated with trauma or burnout included delivery of COVID-19 immunisation (adjusted OR (aOR) 2.54, 95% CI 1.08 to 5.94); and delivery of both COVID-19 immunisation and routine immunisation compared with no involvement in vaccination programmes (aOR 2.42, 95% CI 1.06 to 5.52); poor treatment in the workplace (aOR 2.26, 95% CI 1.51 to 3.38) and lower confidence to respond to patient queries on COVID-19 immunisation (aOR 1.51, 95% CI 1.03 to 2.22).

Conclusion HCPs experienced service disruptions, trauma and burnout and implemented strategies to minimise disruptions to service delivery and improve patient experiences. Our study highlights the need to ensure that workforce resilience and strategies to protect and support

WHAT IS ALREADY KNOWN ON THIS TOPIC

- ⇒ Routine immunisation and maternal and child health services were severely disrupted during the COVID-19 pandemic.
- ⇒ These service disruptions resulted in extreme strain on the health systems including the health workforce.

WHAT THIS STUDY ADDS

- ⇒ This study provides new insights into the perspectives and experiences of healthcare providers (HCPs) delivering essential routine services during the pandemic.
- ⇒ These HCPs work at the subdistrict to village level, providing critical insights at the grass-roots level (ie, *puskesmas* and *posyandu*) in two provinces during the pandemic in Indonesia.
- ⇒ Numerous service disruptions due to the COVID-19 pandemic, including redeployment of staff to COVID-19 relief and lower caregiver attendance to services were experienced by HCPs.
- ⇒ Over a third of HCPs reported being treated poorly by community members in the workplace and nearly two-thirds reported trauma or burnout during the pandemic.
- ⇒ HCPs had increased odds of trauma or burnout if they reported being involved in the delivery of COVID-19 or routine immunisation services, were treated poorly in the workplace or had lower confidence in their ability to respond to patient questions about COVID-19 vaccines.

HCPs are considered for pandemic planning, preparedness and management.

INTRODUCTION

Indonesia is one of the most populated countries in Southeast Asia and was significantly impacted by COVID-19, directly and through

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

- ⇒ Despite constraints during the pandemic, HCPs demonstrated innovation and resilience in the workplace by designing local strategies to minimise service disruptions.
- ⇒ These findings highlight that workforce planning should be prioritised for pandemic planning.
- ⇒ Furthermore, it is important for Indonesian and global policymakers to consider the support system for HCPs as part of future pandemic preparedness and planning.

indirect effects on the broader health system.¹ Indonesia has a decentralised health system that grants district-level governments considerable autonomy to deliver healthcare services.² This impacts availability and quality of services, as well as the capacity of local governments and the workforce to deliver care.² As a result, Indonesia's health system was already experiencing challenges in service delivery when the COVID-19 pandemic was declared. Early in the pandemic, many essential services in Indonesia were interrupted or ceased operation. Health services were further hampered by the diversion of facilities and health workforce to the COVID-19 response.^{1 3 4} Social distancing, lockdowns and fear of contracting COVID-19 were key barriers that contributed to reduced service provision and vaccine uptake.^{4 5} Among the services impacted by disruptions were routine immunisation and maternal and child health (MCH) services. Notably, in April 2020, 84% of government health facilities in Indonesia reported significant disruption to routine immunisation services.¹

MCH services in Indonesia can be classified into three services: antenatal care services, delivery services and services for infants and toddlers, which includes routine immunisation.² Healthcare providers (HCPs) working in MCH services often work across these services of which immunisation is a core component.⁶ MCH services in Indonesia are provided in various levels of health facilities in Indonesia: in the *puskesmas* (primary healthcare centre) and its networks such as *posyandu* (integrated health post) and *polindes* (village maternity cottage) at the community level.² However, during the pandemic, MCH services were centralised from *posyandu* towards *puskesmas* and prioritised for women in their first or third trimester or emergencies. There were limitations on MCH services provided, dwindling number of HCPs and limitation on visitations.⁷

Disruptions across a range of primary healthcare services had a number of negative outcomes for those receiving care.⁸ Sick child services experienced over 50% disruption, while routine outreach immunisation services and facility-based immunisation services experienced 5%–25% disruption levels.⁸ Other MCH services such as well-child visits, antenatal care, postnatal care and facility-based births experienced 5%–25% disruption levels.⁸ For maternal health, the proportion of women accessing four or more antenatal appointments dropped

from 89% nationally in 2019 to 75% in 2021.⁵ According to WHO/UNICEF estimates of national immunization coverage (WUENIC) data, the proportion of children who received the third dose of diphtheria-tetanus-pertussis, hepatitis B and *Haemophilus influenzae* type b (DTP-HB-Hib3) vaccine dropped from 85% nationally in 2019 to 67% in 2021 and the proportion of children who received first dose of DTP-HB-Hib1 dropped from 90% to 74%, and resulted in increased number of zero dose children in 2021.⁹

The impact on the health workforce globally was also significant. A systematic review of global health workers' turnover intention amid COVID-19 pandemic found that the pandemic resulted in excessive job demands and tumultuous work environments for HCPs.¹⁰ In Indonesia, during the first year of the pandemic, 53 000 health workers were trained in detecting, referring and managing COVID-19 confirmed and suspected cases and 38 000 health workers were trained in infection prevention and control.⁵ Despite HCPs reporting institutional barriers and constraints in managing the pandemic, shortages of personal protective gear and infrastructure constraints (eg, patient control and quarantine zones) were faced.^{5 11} These challenges contributed to over 1500 recorded deaths among HCPs from COVID-19 infection in Indonesia in the first 18 months of the pandemic.¹²

There is also clear evidence that the COVID-19 pandemic affected the psychological well-being of the health workforce.^{13 14} In Indonesia, high rates of depressive symptoms (22.8%), anxiety (28.1%) and burnout (26.8%) were reported among HCPs, particularly those dealing directly with the COVID-19 response.^{15–19} Additional studies from Indonesia reported burnout and other psychological distress, contributing to poorer HCP performance and service provision, and increased absenteeism and resignation rates.^{15 18} While there is available data on disruptions to immunisation and MCH services during the pandemic, much of the existing data on personal well-being and concerns of delivering healthcare among HCPs in Indonesia comes from hospital-based settings or multiprovince surveys, with a focus on HCPs actively engaged in the COVID-19 response.^{13 15 16} There is limited data on the experiences and perceptions of HCPs providing routine services such as immunisation and MCH.

In this study, we aimed to examine the disruption to immunisation and MCH services, concerns of personal well-being and delivery of healthcare during the pandemic as well as factors associated with self-reported trauma and burnout among HCPs during the COVID-19 pandemic, in the two Indonesian provinces of Central Java and West Nusa Tenggara.

METHODS

Study location

This study was conducted in two urban and two rural locations in both Central Java and West Nusa Tenggara.

Locations were purposively selected based on the incidence of COVID-19 at the time of study design, the feasibility to implement the study during the pandemic,²⁰ and to capture the urban and rural settings across the provinces. In Central Java, the study was conducted in Surakarta, Semarang, Purbalingga and Demak. In West Nusa Tenggara, the study was conducted in Bima City, Mataram, Central Lombok and Sumbawa. Central Java Province is on the island of Java, with a total population of nearly 37 million across 29 districts and 6 cities, making it the third most populated province in the country out of 38 provinces.²¹ Health and social services account for 1.4% of the overall workforce and it has the fourth highest Gross Domestic Product (GDP).²² West Nusa Tenggara Province comprises eight districts and two cities across several of the Lesser Sunda Islands. It has a population of around 5.4 million, with health and social services accounting for 1.5% of the overall workforce.^{23 24} It is the 13th most populated province with the 25th highest GDP in Indonesia.^{21 23} At the time of data collection, Central Java had 44908 active cases of COVID-19 while West Nusa Tenggara had 2298 cases. In both provinces however, over 80% of adults had received a first dose of COVID-19 vaccination.^{25 26}

During the survey period, certain areas in Central Java were subject to different levels of restrictions. Under level 3 restrictions, the operations and activities of *posyandus* were limited, while under level 4 restrictions, *posyandus* were shut down.²⁷ Conversely, in West Nusa Tenggara, pandemic restrictions varied between level 1 and level 2. Under these regulations, *posyandus* were permitted to resume operations with enhanced health protocols.²⁷

Study design

We conducted a cross-sectional survey among HCPs. Two-stage sampling was used, and the overall sample size was calculated using the formula by Lwanga and Lemeshow for estimating proportions.²⁸ We applied a design effect=2 to account for variance during multistage survey sampling.²⁹ The sample size was estimated to be 282 and was rounded to 300 HCPs per province for recruitment purposes. Detailed methodology is reported elsewhere.³⁰

We invited HCPs aged ≥ 18 years including doctors, nurses, midwives (*bidan*) and village midwives (*bidan desa*) from the public and private sectors working across COVID-19 or routine immunisation and other MCH services. For each location, we sourced and used the district/city health office's HCP registration lists to guide recruitment. We applied simple random sampling to this list to identify participants. The number of substitute participants was dependent on the sample size for each group of HCPs based on location.

Questionnaire design and data collection

The questionnaire covered four topics: the impact of COVID-19 on routine immunisation and MCH services; COVID-19 vaccine perceptions; concerns of personal well-being and delivery of healthcare during the pandemic

and HCP self-efficacy. Survey items were derived from the validated WHO/UNICEF Behavioural and Social Drivers of COVID-19 Vaccination Healthcare Worker survey and relevant items from other surveys, including the WHO Pulse Survey.^{31 32} Within the first topic of the questionnaire, participants who were involved in the provision of routine immunisation were asked several questions on disruptions to these services. Further questions regarding disruptions and strategies to overcome these were in reference to the participant's primary health service of work and did not further differentiate by type of MCH service. Concerns of personal well-being and about delivery of healthcare during the pandemic comprise a number of items which sought to understand HCPs' mental or emotional state (eg, self-reported trauma or burnout), their treatment during service provision by community members and COVID-19-specific concerns about delivery of care such as risk of infection, or confidence to answer questions about the COVID-19 vaccine. Mental and emotional health during the pandemic was measured on a 5-point Likert scale adapted from the CAIR Pandemic Impact Questionnaire.³³ Self-efficacy was measured using the Generalised Self-Efficacy Scale.²⁰ Self-efficacy is defined as a person's belief about their capability to 'produce designated levels of performance that exercise influence over events that affect their lives'.³⁴ The questionnaire is available in online supplemental annex 1.

All study materials were translated into Bahasa Indonesia and pilot tested. Cognitive interviews were conducted with seven HCPs in Depok (Jakarta) and Semarang (Central Java) to test the understanding of survey items, which were then modified as needed. An in-person 2-day training, with some virtual presentations, was conducted for field coordinators and enumerators in Depok and at the survey sites. HCPs were interviewed at their place of work using paper-based questionnaires. Data were collected during 21 March–8 April 2022.

Data analysis

Completed paper-based questionnaires were cross-checked by enumerators and field coordinators. Once complete, data were manually entered into EpiData software by research assistants from Universitas Indonesia. Ten per cent of surveys were double entered for quality assurance. For multiple choice questions, enumerators were provided a list of potential responses but did not prompt participants on these. Participants responded independently and enumerators chose the response(s) that were closest to the predefined options.

Descriptive statistics were used to summarise the data by demographic characteristics and key variables related to service disruptions, personal well-being, interpersonal experiences and concerns about delivering healthcare during the pandemic. Demographic characteristics are stratified by urban and rural study locations.

We used logistic regression to examine the relationship between *self-reported trauma or burnout* among HCPs and key independent variables: HCPs' age, involvement

in COVID-19 or routine immunisation, self-efficacy, perception of increased risk of exposure to COVID-19 in workplace, poor treatment in the workplace, stock-outs of COVID-19 vaccines, confidence to respond to patient questions about COVID-19 vaccines, delivery of COVID-19 vaccines taking away from other priorities, number of service disruptions experienced and number of strategies to overcome disruptions experienced. We created a binary variable for *self-reported trauma or burnout*, by recoding the responses (yes=1 and no/not sure=0) to a single item “Would you say you’ve experienced trauma or burnout related to the COVID-19 pandemic?”

Self-efficacy items were scored using means. *Self-efficacy* was measured using a 10-item scale with four response options on a 4-point scale (not true at all=1; hardly true=2; moderately true=3 and exactly true=4). All scores were totalled and scores ≤ 30 were labelled as ‘moderate self-efficacy’ and >30 as ‘higher self-efficacy’. Responses were then recoded into a binary variable (moderate self-efficacy=0 and high self-efficacy=1). *Confidence to respond to patient questions about COVID-19* used a single item with four response options; responses were recoded into a binary variable (not at all/a little/moderately confident=0 and very confident=1). *Risk perception* was measured using a single item “Do you feel like your job has increased risk of exposure to COVID-19?” (yes=1 and no=0).

Bivariate analyses were first conducted to examine the association between *self-reported trauma or burnout* and potential associated factors. Factors with a p value of <0.20 or with a known association with *self-reported trauma or burnout* were included in the multivariate regression to estimate adjusted OR (aOR) and 95% CIs. All analyses were conducted using STATA V.16.1.

Study governance and ethical approvals

At the start of the study, we formed a Technical Advisory Group to advise on study design and implementation, including members from the Indonesian Ministry of Health, UNICEF Indonesia and EAPRO Offices, the US Centers for Disease Control and Prevention (CDC), Indonesia CDC and WHO Country Offices and the University of Sydney research team. Key approvals and letters of permission from relevant local and national government authorities were obtained prior to study implementation.

All field activities were carried out in a culturally appropriate manner with locally engaged field teams. As the study was conducted during the pandemic, all field activities were conducted in line with a COVID-19 safety protocol developed by the research team. An author reflexivity statement is available in online supplemental annex 2.

Patient and public involvement

The questionnaire was co-designed by local and international research team members with inputs from the Technical Advisory Group. Study findings were disseminated through the Ministry of Health as well as to participants

who requested a summary of results at the time of data collection.

RESULTS

We recruited 604 HCPs (target=600) to participate in the study, 304 (50.3%) were from Central Java and 300 (49.7%) from West Nusa Tenggara). Detailed characteristics of HCPs who participated in the study are presented in [table 1](#). Four hundred and fifty HCPs (74.5%) were from rural areas. Most HCPs were female (84.6%; 511/604), aged 31–50 years (44.5%; 269/604) and held a diploma as their highest level of education (75.5%; 456/604). The majority of HCPs were midwives (40.2%, 243/604) followed by nurses (30.8%, 186/604). Most HCPs had >10 years of experience (56.8%; 343/604). The primary place of work reported by HCPs (78.1%, 472/604) was the *puskesmas* (subdistrict primary healthcare clinic). Of the 604 HCPs, 470 (77.8%) were involved in the delivery of COVID-19 immunisation and 384 (63.6%) were involved in routine immunisation. Additionally, just under 50% provided MCH services such as labour and birth (294/604; 48.7%) or other baby and child service such as growth monitoring (285/604; 47.2%).

Service disruptions experienced by HCPs during the pandemic

All HCPs reported experiencing service disruptions during the pandemic. Among HCPs who provided immunisation services, 62.8% (355/565) reported stockouts of routine vaccines. Of these, 63.9% reported stockouts of inactivated polio vaccine (227/355) ([table 2](#)). Just over a third of HCP (35.4%; 214/604) reported stockouts of COVID-19 vaccines. Mobilisation of staff to COVID-19 clinical management was the most reported reason for service disruption (87.9%; 531/604), followed by changes to vaccination policies (60.1%; 363/604). HCPs implemented numerous vaccination strategies to attempt to overcome service disruptions, including community outreach (75.5%; 456/604) and phone calls to parents or caregivers (72.4%; 437/604), among others ([table 2](#)).

Trauma, burnout and concerns of personal well-being and delivery of healthcare experienced by HCPs delivering immunisation and MCH services during the pandemic

Of the 604 HCPs, 64.1% (387/604) reported experiencing trauma or burnout during the pandemic; 23.5% (142/604) felt that their mental or emotional health had worsened by the pandemic ([table 3](#)). Of those who experienced trauma or burnout, longer working hours were the most common reason (60.2%; 233/387). For interpersonal relations, 35.3% (213/604) reported poor treatment (by the community members) in their workplace because they were health workers. Among them, 23% described patient anger at longer wait times (49/213).

Ninety-three per cent (565/604) perceived an increased risk of exposure to COVID-19 in their workplace, 66.4% (401/604) were concerned about getting sick with COVID-19 and 37.3% (225/604) had heard something worrying about the COVID-19 vaccine. Despite

Table 1 Demographic and work-related characteristics of healthcare providers in Central Java and West Nusa Tenggara, Indonesia, March–April 2022 (n=604)

	Total n (%) N=604	Urban n (%) N=154 (25.5)	Rural n (%) N=450 (74.5)
Location			
Central Java	304 (50.3)	120 (77.9)	184 (40.9)
West Nusa Tenggara	300 (49.7)	34 (22.1)	266 (59.1)
Gender			
Female	511 (84.6)	134 (87.0)	377 (83.8)
Male	93 (15.4)	20 (13.0)	73 (16.2)
Age (years)			
18–30	157 (26.0)	35 (22.7)	122 (27.1)
31–40	269 (44.5)	58 (37.7)	211 (46.9)
41–50	122 (20.2)	33 (21.4)	89 (19.8)
51 and older	56 (9.3)	28 (18.2)	28 (6.2)
Educational attainment			
Diploma	456 (75.5)	100 (64.9)	356 (79.1)
Undergraduate/Professional degree	138 (22.8)	48 (31.2)	90 (20.0)
Masters/PhD	10 (1.7)	6 (3.9)	4 (0.9)
Current role			
Doctor	40 (6.6)	22 (14.3)	18 (4.0)
Nurse	186 (30.8)	53 (34.4)	133 (29.6)
Midwife	243 (40.2)	78 (50.7)	165 (36.7)
Village midwife	135 (22.4)	1 (0.6)	134 (29.8)
Years in role			
<2	35 (5.8)	8 (5.2)	27 (6.0)
2 to <5	88 (14.6)	22 (14.3)	66 (14.7)
5 to <10	138 (22.8)	28 (18.2)	110 (24.4)
10 or more	343 (56.8)	96 (62.3)	247 (54.9)
Primary place of work*			
Puskesmas	472 (78.1)	123 (79.9)	349 (77.6)
Poskesdes	42 (7.0)	0 (0.0)	42 (9.3)
Independent midwife practice	33 (5.5)	1 (0.6)	21 (4.7)
Polindes	22 (3.6)	1 (0.6)	16 (3.6)
Pustu	17 (2.8)	23 (14.9)	10 (2.2)
Other	14 (2.3)	0 (0.0)	4 (0.9)
Private clinic	4 (0.7)	6 (3.9)	8 (1.8)
Secondary place of work† ‡			
None	429 (71.0)	126 (81.8)	303 (67.3)
Puskesmas	45 (7.5)	4 (2.6)	41 (9.1)
Poskesdes	21 (3.5)	0 (0.0)	21 (4.7)
Pustu	8 (1.3)	0 (0.0)	8 (1.8)
Independent midwife practice	78 (12.9)	7 (4.5)	71 (15.8)
Private clinic	19 (3.1)	9 (5.8)	10 (2.2)
Other	28 (4.6)	10 (6.5)	18 (4.0)
HCP area of work†			
COVID-19 immunisation	470 (77.8)	120 (77.9)	350 (77.8)

Continued

Table 1 Continued

	Total n (%) N=604	Urban n (%) N=154 (25.5)	Rural n (%) N=450 (74.5)
Routine child immunisation	384 (63.6)	97 (63.0)	287 (63.8)
Antenatal care	376 (62.3)	86 (55.8)	290 (64.4)
Family planning	356 (58.9)	83 (53.9)	273 (60.7)
Postnatal care	338 (56.0)	71 (46.1)	267 (59.3)
Labour and birth	294 (48.7)	56 (36.4)	238 (52.9)
Baby and child services (weighing, vitamin A supplementation, etc)	285 (47.2)	82 (53.2)	203 (45.1)
Other	187 (31.0)	56 (36.4)	131 (29.1)

*Puskesmas—primary healthcare centre; poskesdes—village health post; polindes—village maternity post; pustu—auxiliary health centre.
†Multiple response options allowed.
‡Secondary workplace referred to HCPs working in multiple health facilities.
HCP, healthcare provider.

this, 63.3% (382/604) reported feeling very confident to answer patient questions about the COVID-19 vaccine.

Factors associated with self-reported trauma or burnout

HCPs aged 30 to <40 years had higher odds of self-reported trauma or burnout compared with those aged 18 to <30 years (aOR 1.73, 95% CI 1.06 to 2.83) (table 4). HCPs involved in the delivery of COVID-19 immunisation alone (aOR 2.54, 95% CI 1.08 to 5.94) or COVID-19 and routine immunisation (aOR 2.42, 95% CI 1.06 to 5.52) had higher odds of self-reported trauma or burnout compared with those not involved in any vaccination programme. HCPs who reported poor treatment in the workplace had significantly higher odds of self-reported trauma or burnout compared with HCPs who did not report being treated poorly in the workplace or were not sure (aOR 2.26, 95% CI 1.51 to 3.38). HCPs who were not at all, a little or moderately confident to respond to patient questions about the COVID-19 vaccine had higher odds of self-reported trauma or burnout compared with HCPs who were very confident to respond to questions (aOR 1.51, 95% CI 1.03 to 2.22).

DISCUSSION

In this study, we present perspectives of HCPs on disruption to immunisation and MCH services and concerns of personal well-being and delivery of healthcare during the COVID-19 pandemic in Central Java and West Nusa Tenggara provinces, Indonesia. To the authors' knowledge, this is the first paper to provide these insights into HCP experiences working at the grassroots level (ie, *puskesmas* and *posyandu*) in the delivery of essential routine health services. Self-reported trauma and burnout among HCPs was significant. Several factors were associated with higher odds of self-reported trauma or burnout including poor treatment in the workplace by the community, involvement in delivery of COVID-19 or routine immunisation and confidence to respond to patient questions about COVID-19 vaccine.

Around one in five (20.7%) HCPs reported that COVID-19 immunisation had disrupted other priorities and health services. Other service disruptions included staff mobilisation to COVID-19 response (87.9%), change in vaccination policies (60.1%) and fewer caregivers attending clinics (55.0%). Despite experiencing disruptions, HCPs reported using local solutions to overcome these disruptions including community outreach activities, follow-up phone calls to caregivers and task shifting. Stockout of routine vaccines was reported by 62.8% of HCPs.

Globally, in 2021, when this study was initiated, the vaccination coverage for DTP dose 1 dropped to 86% compared with 90% in 2019.^{35 36} In Indonesia, first dose of DTP dropped to 74% in 2021 from 90% in 2019.^{35 36} As highlighted in The State of the World's Children 2023,³⁵ the main reason for this drop in vaccination was disruption to routine health services with overstretched primary healthcare and health systems. Global reports found that already limited resources were redirected from primary care, with 49% (35/72) of countries in the 2021 WHO Pulse survey reporting routine vaccination programmes were being disrupted by the need to respond to the pandemic.³² Reasons varied across the countries, including challenges related to closure of clinics, lack of medicines, diagnostics and vaccines. Vaccine stockouts across health services were reported previously in Indonesia,¹ but the variation across different types of vaccines suggested province-specific challenges with vaccine supply.¹ Supply issues were also reported in a multicountry study with 8/11 member states in the WHO South-East Asia region.³⁷ One of the key lessons learnt through the pandemic was its impact on the health workforce, with health worker availability and capacity as a key reason for disruption to health services globally.^{13 14 32}

Besides routine immunisation, other MCH services were also impacted by COVID-19 globally. In India in 2021, compared with 2019, antenatal care registration declined 18% and obstetric care declined by 40.9%.³⁸

Table 2 Service disruptions to routine immunisation and maternal and child health services experienced during the COVID-19 pandemic by healthcare providers in Central Java and West Nusa Tenggara, Indonesia, March–April 2022 (n=604)

	N (%)
Routine vaccine stockout (n=565)*	
No/Not sure	210 (37.2)
Yes	355 (62.8)
Stockouts for routine vaccines (n=355)*†	
IPV	227 (63.9)
DTP-HB-Hib	191 (53.8)
BCG	136 (38.3)
MR	131 (36.9)
HepB	39 (11.0)
PCV	61 (17.2)
OPV	36 (10.1)
Other	34 (9.6)
Vaccine stockout for COVID-19 vaccines	
No/Not sure	378 (62.6)
Yes	214 (35.4)
Do not stock	12 (2.0)
Formal training in COVID-19 immunisation	
No	287 (47.5)
Yes	317 (52.5)
COVID-19 immunisation disrupted other priorities	
Did not deliver COVID-19 immunisation	72 (11.9)
Not at all/Slightly	407 (67.4)
Moderately	95 (15.7)
Very	30 (5.0)
Service disruptions experienced by HCP†‡	
Staff providing COVID-19 relief	531 (87.9)
Change in vaccination policies	363 (60.1)
Fewer parents/caregivers attending for other reasons	332 (55.0)
Lockdowns hindering parent/caregiver access	211 (34.9)
Closure of service	202 (33.4)
Insufficient PPE	185 (30.6)
Not enough staff	184 (30.5)
Unavailability/Stockouts of vaccines or injecting equipment at health facility	169 (28.0)
Other	23 (3.8)
Strategies to overcome disruptions†	
Community outreach	456 (75.5)
Phone calls to parents/caregivers	437 (72.4)
Task shifting/role delegation	387 (64.1)
Parents/Caregivers seen outdoors from facility	358 (59.3)
Redirection of parents/caregivers to alternative services	230 (38.1)
Government removal of user fees	154 (25.5)

Continued

Table 2 Continued

	N (%)
Other (eg, home visits, online consultations)	43 (7.1)

*Only participants who provided routine immunisation services answered this question.
 †Multiple response options allowed.
 ‡100% of respondents reported some type of service disruption.
 DTP-HB-Hib, diphtheria-tetanus-pertussis, hepatitis B and *Haemophilus influenzae* type b; HepB, hepatitis B; IPV, inactivated polio vaccine; MR, measles and rubella; OPV, oral polio vaccine; PCV, pneumococcal conjugate vaccine; PPE, personal protective equipment.

In Ethiopia, MCH services were disrupted, and services became inaccessible and low quality. Additionally, HCPs experienced scarcity of personal protective equipment (PPE), increased workload and shortages in resources.³⁹ Similarly in Indonesia, MCH services experienced challenges in operating, including disruption to services, shortages of human and financial resources, difficulty in travelling and overall closure of health facilities.⁴⁰

Similar to our study, other studies in Indonesia found diversion of staff to the COVID-19 response, as well as challenges with caregiver access to services or fear of attendance, have been reported in other MCH services in Indonesia.⁴⁰ Routine MCH services responded proactively throughout the pandemic to minimise the impact of disruptions, demonstrating resilience and motivation. Telemedicine, mobile phone chats and community outreach with door-to-door home visits have been reported as strategies to minimise these disruptions.⁴⁰ Overall, there is clear recognition of the need to strengthen primary healthcare and health systems to improve care during routine times and to prevent shocks to essential health services during emergencies. Pandemic plans often focus on responding to the infectious diseases outbreak, and need to consider how to strengthen resilience in the health system to maintain routine services.⁴¹

A large proportion (64.1%) of the 604 HCPs reported experiencing trauma or burnout during the pandemic, resulting from long work hours (60.2%), shifting work duties (26.9%) and changes in service delivery (23.8%). In our study, 35.3% HCPs reported being treated poorly in the workplace during the pandemic. Health workers who reported poor treatment were at 2.3 times greater odds of experiencing trauma and burnout compared with those who did not experience poor treatment. HCPs involved in delivery of COVID-19 immunisation alone or COVID-19 and routine immunisation had 2.4–2.5 greater odds of burnout compared with HCPs not involved in any vaccination programme. Similarly, those who reported not feeling confident in responding to patient questions about COVID-19 vaccines were at 1.5 times greater odds of experiencing trauma and burnout than those who felt confident to answer questions.

Table 3 Trauma, burnout and concerns of personal well-being and delivery of healthcare experienced by HCPs delivering immunisation and MCH services during the pandemic in Central Java and West Nusa Tenggara, Indonesia, March–April 2022 (n=604)

	N (%)
Experienced trauma or burnout during pandemic	
No/Not sure	217 (35.9)
Yes	387 (64.1)
Reasons for trauma or burnout (n=387)*†	
Long work hours	233 (60.2)
Shifting work duties	104 (26.9)
Changes in service delivery	92 (23.8)
Use of PPE	65 (16.8)
Fear of making family ill	57 (14.7)
Patient distress	55 (14.2)
Fear of illness	52 (13.4)
Other personal factors	10 (2.3)
Other service-delivery factors	18 (4.7)
Other patient-related factors	17 (4.4)
Other	15 (3.9)
Mental/Emotional health worsened by pandemic	
Not at all/Slightly	462 (76.5)
Moderately/Very/Extremely	142 (23.5)
Treated poorly during pandemic	
No/Not sure	391 (64.7)
Yes	213 (35.3)
Description of poor treatment (n=213)*‡	
Patients angered by longer wait times	49 (23.0)
Patient perceptions of COVID-19 or pandemic	42 (19.7)
Patients believe I am a COVID-19 risk	35 (16.4)
Patients angered by request for medical exemption	30 (14.1)
Service delivery issues	21 (9.9)
Patients angered by the vaccine brands offered	21 (9.9)
Other patient COVID-19 fears or stigma	19 (8.9)
Negative perceptions of healthcare providers	10 (4.7)
Challenges with patient adherence to safety protocols	10 (4.7)
Patients angry at service closures	5 (2.3)
Other	15 (7.0)
Self-efficacy	
Moderate self-efficacy	198 (32.8)
High self-efficacy	406 (67.2)
Concern about getting COVID-19	
Not at all/A little concerned	203 (33.6)
Moderately concerned	170 (28.2)

Continued

Table 3 Continued

	N (%)
Very concerned	231 (38.2)
Confidence to answer questions about COVID-19 vaccine	
Not at all/A little confident	8 (1.3)
Moderately confident	214 (35.4)
Very confident	382 (63.3)
Heard something worrying about COVID-19 vaccine	
No/Not sure	379 (62.7)
Yes	225 (37.3)
Perception of increased risk of exposure to COVID-19 at work	
No	39 (6.5)
Yes	565 (93.5)

*Multiple response options allowed.
 †For those who reported yes to experiencing trauma or burnout during the pandemic.
 ‡Of those who responded to yes for being treated poorly because they are a healthcare worker.
 HCP, healthcare provider; MCH, maternal and child health; PPE, personal protective equipment.

Our finding of HCPs experiencing trauma and burnout and related factors are not surprising. Prior to the COVID-19 pandemic, some studies examined burnout among nurses in several provinces in Indonesia.^{42 43} The COVID-19 pandemic has, however, placed a spotlight on the specific issues of mental health, trauma, violence or abuse against HCPs by patients or relatives of patients.^{28 44–46} Findings on poor treatment in the workplace also align with global reports where healthcare workers experienced stigmatisation and discrimination.⁴⁷ Other studies have reported links between symptoms of depression and burnout with increased contact with COVID-19 patients or involvement in the COVID-19 response.^{18 19} In addition, conflict of balancing work and family, emotional demands (eg, aggressive or troublesome patients) and job demands (eg, lack of preparedness in role) were also associated with burnout.⁴⁸ Our findings regarding concerns of personal well-being and delivery of healthcare during the pandemic have been reflected among wider HCP populations in Indonesia. Reports that looked at HCPs working in COVID-19 response also found that these HCPs experienced increased demands, high patient loads, long working hours and limited PPE.⁴⁹ HCPs also reported experiencing stigma and discrimination from the communities, some resulting in deprivation of essential services, social alienation, even physical assault.⁵⁰ This study did not report on physical assault, however patient anger at longer wait times was a commonly reported problem among HCP in our sample. In the Philippines, HCPs were required to stay in facilities to prevent risk of COVID-19 transmission to their family, while enduring more frequent and longer hours

Table 4 Factors associated with self-reported trauma or burnout among healthcare providers in Central Java and West Nusa Tenggara, March–April 2022

	N (%)	OR (95% CI)	P value	aOR (95% CI)	P value
Gender of HCP					
Male	93 (15.4)	Reference			
Female	511 (84.6)	1.03 (0.65 to 1.63)	0.89		
Age (years)*					
18 to <30	114 (18.9)	Reference		Reference	
30 to <40	295 (48.8)	1.52 (0.97 to 2.36)	0.07	1.73 (1.06 to 2.83)	0.03
40 to <50	132 (21.9)	1.56 (0.93 to 2.62)	0.09	1.73 (0.98 to 3.05)	0.06
50 and older	63 (10.4)	1.01 (0.54 to 1.87)	0.99	1.47 (0.74 to 2.89)	0.27
Current role					
Doctor	40 (6.6)	Reference			
Nurse	186 (30.8)	1.23 (0.60 to 2.50)	0.57		
Midwife	243 (40.2)	0.89 (0.45 to 1.77)	0.74		
Village midwife	135 (22.4)	1.28 (0.62 to 2.68)	0.51		
Years in role					
Between 2 and <5	88 (14.6)	Reference			
<2	35 (5.8)	0.90 (0.40 to 2.01)	0.80		
5 or more years	481 (79.6)	1.10 (0.69 to 1.76)	0.70		
Involvement in RI or COVID-19 immunisation*†					
Neither RI or COVID-19 immunisation	39 (6.5)	Reference		Reference	
Just RI	95 (15.7)	1.68 (0.79 to 3.55)	0.18	2.14 (0.94 to 4.90)	0.07
Just COVID-19 immunisation	181 (29.9)	2.13 (1.06 to 4.29)	0.03	2.54 (1.08 to 5.94)	0.03
Both RI and COVID-19 immunisation	289 (47.9)	2.46 (1.25 to 4.83)	0.009	2.42 (1.06 to 5.52)	0.03
Self-efficacy*					
Moderate	198 (32.8)	Reference		Reference	
High	406 (67.2)	0.71 (0.50 to 1.02)	0.07	0.83 (0.56 to 1.23)	0.35
Perception of increased risk of exposure to COVID-19 in workplace*					
No	39 (6.5)	Reference		Reference	
Yes	565 (93.5)	2.20 (1.14 to 4.22)	0.02	1.76 (0.88 to 3.52)	0.11
Treated poorly in workplace during COVID-19 pandemic*					
No/Not sure	391 (64.7)	Reference		Reference	
Yes	213 (35.3)	2.62 (1.79 to 3.82)	<0.001	2.26 (1.51 to 3.38)	<0.001
Stockouts of COVID-19 vaccines (n=592)*‡					
No/Not sure	378 (63.9)	Reference		Reference	
Yes	214 (36.1)	1.69 (1.18 to 2.42)	0.01	1.29 (0.86 to 1.92)	0.21
Heard worrying information about COVID-19 vaccine					
No	379 (62.7)	Reference			
Yes	225 (37.3)	1.13 (0.80 to 1.59)	0.50		
Confidence to respond to patient questions about COVID-19 vaccine*					
Very confident	382 (63.3)	Reference		Reference	
Not at all/A little confident/Moderately	222 (36.7)	1.36 (0.96 to 1.93)	0.09	1.51 (1.03 to 2.22)	0.04
Concerned about getting COVID-19					
Not at all/A little/Moderately concerned	373 (61.8)	Reference			
Very concerned	231 (38.2)	1.00 (0.71 to 1.41)	0.99		
Delivering COVID-19 vaccines has taken away from other priorities*					
Did not deliver COVID-19 vaccine	72 (11.9)	Reference		Reference	

Continued

Table 4 Continued

	N (%)	OR (95% CI)	P value	aOR (95% CI)	P value
Not at all/Slightly	407 (67.4)	1.29 (0.78 to 2.13)	0.33	0.79 (0.39 to 1.59)	0.51
Moderately/Very	125 (20.7)	2.65 (1.42 to 4.94)	0.01	1.27 (0.56 to 2.91)	0.57
Number of service disruptions experienced*					
3 or less	299 (49.5)	Reference		Reference	
4 or more	305 (50.5)	1.52 (1.09 to 2.13)	0.01	1.20 (0.82 to 1.75)	0.34
Number of vaccination strategies to overcome disruptions experienced*					
3 or less	309 (51.2)	Reference		Reference	
4 or more	295 (48.8)	1.59 (1.14 to 2.22)	0.01	1.29 (0.87 to 1.89)	0.18

*Variable included in multivariable analysis due to bivariate results or known association to outcome.
†This variable only looks at routine immunisation and COVID-19 immunisation. Participants may have also been involved in other maternal and child health services (table 1).
‡This variable excludes participants whose place of work did not stock COVID-19 vaccines (n=12).
HCP, healthcare provider; RI, routine immunisation.

of working, increasing demand, medical supply shortages and harassment, bullying and abuse.⁵¹ In Nepal, HCPs reported lack of PPE and experiences of stigma lead to increased burnout, fatigue and psychological distress.⁵² In Jordan, 65.5% of HCPs in a public hospital reported exposure to workplace verbal and physical violence during the pandemic.⁴⁵

The association between trauma and burnout with delivery of COVID-19 immunisation and the ability to respond to patient queries might also relate to the demand on the health workforce to rapidly gain and implement new concepts related to COVID-19 vaccines and vaccination processes; this highlights the public's reliance on health workers for trusted information. Prior work by some of the study team members found that HCP recommendation is a strong motivator for getting a vaccination.^{53 54} Interestingly, in our study, 47.5% HCPs reported not having received formal training in COVID-19 immunisation. These findings highlight the need for mechanisms and innovative ways to ensure the availability of timely, tailored and targeted training and upskilling for the health workforce during acute emergencies.

The pandemic also negatively impacted the mental health of HCPs globally. A 2021 systematic review found pooled prevalence of anxiety and depression resulting from the COVID-19 pandemic to be 37% and 36%, respectively.¹⁴ Other Southeast Asian countries had similar experiences; in Vietnam, 18%, 11.5% and 7.7% of participants reported symptoms of depression, anxiety and stress, respectively.⁵⁵ Interestingly, despite a large proportion of HCPs experiencing trauma and burnout from the pandemic in this study, only 23.5% HCPs reported worsened mental and emotional health. It is possible that HCPs may be more willing or perceive it to be more socially acceptable to report physical reasons for trauma or burnout (eg, 'long work hours') rather than to report a decline in emotional or mental health. Disclosure of mental health conditions in the workplace

is often difficult due to barriers and perceived risk such as discrimination.⁵⁶ Prior studies exploring the negative impact of the COVID-19 pandemic on the mental health of HCPs in Indonesia found clinical factors (eg, lack of PPE), personal (eg, fear of transmitting COVID-19) and social factors (eg, stigma) related to HCPs' psychological well-being.^{15 17 18} These findings highlight the need for HCPs to be better supported in the health system, particularly during emergency settings when the burden is inevitably higher. Previous studies in Indonesia used support systems to strengthen the resilience of HCPs, provision of psychological support (eg, therapy), the encouragement to access these services and incentives and compensation for working in a high-risk setting.¹⁶⁻¹⁸ In some low-resource settings, such as Sierra Leone, Liberia, India, Bangladesh and the Democratic Republic of Congo, telephone hotlines, care coordination teams and use of psychological support applications were also used to support HCPs' mental well-being during the COVID-19 pandemic.^{57 58} This additional support to the workforce, as well as other modifications to factors such as management of waiting processes at the clinic, may also help to improve patient experience. The pandemic has also highlighted the need to re-imagine the incorporation of health workforce planning as a crucial component of the health system. A recent Organisation for Economic Co-operation and Development report found that the health workforce in many countries is understaffed, under pressure and undervalued.⁵⁹ Health workforce shortages existed even prior to the pandemic and were only magnified through the pandemic. Health workers also played a crucial role in absorbing the shocks of the pandemic and risking their own and family's health and well-being. However, the future requires revived momentum that focuses on sustainable and resilient capacity to ensure sufficient numbers, and proper mix and distribution of health workers to respond to population needs. The WHO Roadmap for the public health workforce highlights the need to train a workforce

that can perform essential public health functions, with a focus on developing national workforce strategies that can cater to routine services, health functions and provide surge capacity during an acute emergency.^{60–62} These approaches can also be expanded to include immunisation and MCH specialist skills.

Future research is needed that further explores the reasons for trauma and burnout among HCPs in Indonesia, including the relationship with ability to respond to patient questions related to COVID-19 immunisation, and reasons for discordance between HCP reports of trauma and burnout and mental and emotional health. Our study had multiple strengths including the use of validated and field-tested survey tools^{31 32} and relatively large sample sizes within the selected provinces. Furthermore, to our knowledge, this is the only paper to explore perspectives and experiences of HCPs working at the provincial level during the pandemic in Indonesia. The study also had some limitations. Our study provides measurement of associations between HCP-reported influences on self-reported trauma or burnout at a single point in time, therefore trends could not be monitored. As the study was only conducted in two provinces, and given Indonesia's decentralised health system, study results cannot be generalised. Our HCP survey sampling frame did not draw from the overall population of HCP in each study location, but rather focused on those working in immunisation and MCH services. As a result, we could not apply population weights to adjust for the survey design. Due to questionnaire design and the structure of co-delivery of immunisation and other MCH services in Indonesia, analysis was not stratified by type of service. Lastly, we used a single questionnaire item to capture data on HCP experiences and perspectives which was not designed to measure trauma or burnout. Our analysis was exploratory, results should be interpreted with caution. Future studies would benefit from the use of a validated scale of trauma or burnout and prospectively collected data to confirm our findings.^{16 17 19} Insights from our study can help contribute towards policies to improve HCP experiences during an emergency.

CONCLUSION

Our study reported on experiences and perspectives of the HCP on the disruption to immunisation and MCH services, which are critical components of the primary healthcare system. We found HCPs experienced service disruption, trauma and burnout. However, we also identified local solutions aimed at minimising disruption to service delivery and improving patient experiences, particularly in immunisation and MCH. We highlight the need to ensure that workforce resilience is placed at the centre of the health system and is considered as an integral part of pandemic planning, preparedness and management to ensure continuity of essential health services.

Author affiliations

¹The University of Sydney, School of Public Health, Faculty of Medicine and Health, Sydney, New South Wales, Australia

²Universitas Indonesia, Department of Epidemiology, Faculty of Public Health, Jakarta, Indonesia

³UNICEF East Asia and Pacific Regional Office, Bangkok, Thailand

⁴Murdoch Children's Research Institute, Parkville, Victoria, Australia

⁵The University of Melbourne, Melbourne School of Population and Global Health, Melbourne, Victoria, Australia

⁶Universitas Diponegoro, Department of Epidemiology, Faculty of Public Health, Semarang, Indonesia

⁷UNICEF Country Office Indonesia, Jakarta, Indonesia

⁸Ministry of Health of the Republic of Indonesia, Jakarta, Indonesia

X Madeleine Randell @maddierandell1, Julie Leask @julieleask and Meru Sheel @merusheel

Acknowledgements We thank the data enumerators and study participants for their contributions to this study. We would like to thank members of the Technical Advisory Group for their input into study design and implementation. We also thank Dr Shibani Kulkarni from the US Centers for Disease Control and Prevention for critical review of the manuscript.

Contributors JL, MShe, TYMW, MR, MDA, AT, ML and MDy designed the overall study. MR, MDy, MS, JL conceptualised and designed this paper. TYMW, OO, FS and LDS supervised and implemented field work and data collection and entry. MR and MShe analysed the data. JL, MR, TYMW, OO, KJ, MShe, MDy, FS and LDS supported the survey training. KDA, AKN, LD and PY provided guidance for study design, recommendations and implications. MShe and MR wrote the initial drafts of the manuscript. All authors critically reviewed and provided input the manuscript. JL was responsible for overall supervision and management of the project. MShe is responsible for the overall content as the guarantor of this work.

Funding The study was funded by the UNICEF East Asia and Pacific Regional Office.

Competing interests Julie Leask receives funding from the World Health Organization and has received reimbursement for travel to an overseas meeting from Sanofi.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research. Refer to the Methods section for further details.

Patient consent for publication Not applicable.

Ethics approval This study was approved by University of Sydney Human Research Ethics Committee (2022/010) and the University of Indonesia (Ket-23/UN2.F10.D11/PPM.00.02/2022). Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request. All data relevant to the study are included in the article or uploaded as supplementary information. Aggregated data are available however, data files are not available to be shared unless on reasonable request. Please send requests for data to human.ethics@sydney.edu.au who will confirm with data custodians.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Madeleine Randell <http://orcid.org/0000-0002-5264-5295>

Julie Leask <http://orcid.org/0000-0001-5095-1443>

Meru Sheel <http://orcid.org/0000-0001-5453-070X>

REFERENCES

- UNICEF. *Impact of COVID-19 on routine immunization in Indonesia*. Jakarta, Indonesia: UNICEF, 2022. Available: <https://www.unicef.org/indonesia/reports/impact-covid-19-routine-immunization-indonesia>

- 2 Mahendradhata Y, Trisnantoro L, Listyadewi S, et al. *The Republic of Indonesia health system review*. WHO Regional Office for South-East Asia, 2017. Available: <https://apps.who.int/iris/handle/10665/254716>
- 3 World Health Organization. Indonesia: WHO Coronavirus disease (COVID-19) dashboard with vaccination data. 2023. Available: <https://covid19.who.int>
- 4 Downey LE, Gadsden T, Vilas VDR, et al. The impact of COVID-19 on essential health service provision for endemic infectious diseases in the South-East Asia region: a systematic review. *Lancet Reg Health Southeast Asia* 2022;1:100011.
- 5 UNICEF. Solidarity, resilience and hope in the COVID-19 response. 2022. Available: <https://www.unicef.org/indonesia/reports/solidarity-resilience-and-hope-covid-19-response>
- 6 Maharani A, Kuroda Y. Determinants of immunization status among 12- to 23-month-old children in Indonesia (2008-2013): a multilevel analysis. *BMC Public Health* 2018;18:288.
- 7 Saputri NS, Anbarani MDA, Toyamah N, et al. *The Impact of the COVID-19 Pandemic on Nutrition and Maternal and Child Health (MCH) Services: case study in five regions in Indonesia*. Jakarta, Indonesia: The SMERU Research Institute, 2020. Available: <https://smeru.or.id/en/content/impact-covid-19-pandemic-nutrition-and-maternal-and-child-health-mch-services-case-study>
- 8 World Health Organization. Tracking continuity of essential health services during COVID-19 pandemic. 2021. Available: <https://www.who.int/teams/integrated-health-services/monitoring-health-services/global-pulse-survey-on-continuity-of-essential-health-services-during-the-covid-19-pandemic/dashboard>
- 9 WHO/UNICEF Estimates of National Immunization Coverage (WUENIC). Immunization data. Diphtheria Tetanus Toxoid and pertussis (DTP) vaccination Coverage- Indonesia. 2021. Available: <https://immunizationdata.who.int/global/wiise-detail-page>
- 10 Poon YSR, Lin YP, Griffiths P, et al. A global overview of healthcare workers' turnover intention amid COVID-19 pandemic: a systematic review with future directions. *Hum Resour Health* 2022;20:70.
- 11 Fauk NK, Seran AL, Raymond C, et al. Why do we not follow lifesaving rules? Factors affecting Nonadherence to COVID-19 prevention guidelines in Indonesia: healthcare professionals' perspectives. *Int J Environ Res Public Health* 2022;19:8502.
- 12 Ekawati LL, Arif A, Hidayana I, et al. Mortality among healthcare workers in Indonesia during 18 months of COVID-19. *PLOS Glob Public Health* 2022;2:e0000893.
- 13 Sheraton M, Deo N, Dutt T, et al. Psychological effects of the COVID 19 pandemic on Healthcare workers globally: a systematic review. *Psychiatry Res* 2020;292:113360.
- 14 Sun P, Wang M, Song T, et al. The psychological impact of COVID-19 pandemic on health care workers: a systematic review and meta-analysis. *Front Psychol* 2021;12:626547.
- 15 Agustina TS, Rarastanti PD, Hidayat AF. Stress and job performance of healthcare workers amidst COVID-19 pandemic: the mediating role of burnout. *Shirkah* 2021;6:315-35.
- 16 Daryanto B, Rahmadiani N, Amorga R, et al. Burnout syndrome among residents of different surgical specialties in a tertiary referral teaching hospital in Indonesia during COVID-19 pandemic. *Clin Epidemiol Glob Health* 2022;14:100994.
- 17 Setiawati Y, Wahyuhadi J, Joestandari F, et al. Anxiety and resilience of healthcare workers during COVID-19 pandemic in Indonesia. *J Multidiscip Healthc* 2021;14:1-8.
- 18 Sunjaya DK, Herawati DMD, Siregar AYM. Depressive, anxiety, and burnout symptoms on health care personnel at a month after COVID-19 outbreak in Indonesia. *BMC Public Health* 2021;21:227.
- 19 Soemarmo DS, Basrowi RW, Chandra Khoe L, et al. Prevalence and determinant factors of health workers burnout during COVID-19 pandemic in Indonesia. *Safety and Health at Work* 2022;13:S211.
- 20 Schwarzer R, Jerusalem M. Generalized self-efficacy scale. In: Weinman J, Wright S, Johnston M, eds. *Causal and control beliefs. (Measures in health psychology: a user's portfolio)*. Windsor, UK: NFER - Nelson, 1995: 35-7.
- 21 Badan Pusat Statistik Jawa Tengah. Jawa Tengah Dalam Angka 2022. Semarang: Badan Pusat Statistik Jawa Tengah, 2023. Available: <https://jateng.bps.go.id/publication/2022/02/25/431f4f4bbe02b47866b357cc/provinsi-jawa-tengah-dalam-angka-2022.html>
- 22 Badan Pusat Statistik Jawa Tengah. Profil Ketenagakerjaan Provinsi Jawa Tengah: Hasil Sakernas Agustus 2021. Semarang: Badan Pusat Statistik Jawa Tengah, 2022. Available: <https://jateng.bps.go.id/publication/2022/06/08/706b8f88e68be79428a7213a/profil-ketenagakerjaan-provinsi-jawa-tengah-hasil-sakernas-agustus-2021.html>
- 23 Badan Pusat Statistik Nusa Tenggara Barat. Nusa Tenggara Barat Dalam Angka 2022. Mataram: Badan Pusat Statistik Nusa Tenggara Barat, 2022. Available: <https://ntb.bps.go.id/publication/2022/02/25/81b407c481be37afd75d6f5/provinsi-nusa-tenggara-barat-dalam-angka-2022.html>
- 24 Badan Pusat Statistik Nusa Tenggara Barat. Profil Ketenagakerjaan Provinsi Nusa Tenggara Barat: Berdasarkan Hasil Sakernas Agustus 2021. Mataram: Badan Pusat Statistik Nusa Tenggara Barat, 2022. Available: <https://ntb.bps.go.id/publication/2024/05/31/52bf8f10786671b01ed2b052/profil-ketenagakerjaan-provinsi-nusa-tenggara-barat-berdasarkan-hasil-sakernas-agustus-2023.html>
- 25 Kementerian Kesehatan Indonesia. Vaksinasi COVID-19 Nasional. 2024. Vaksin dashboard. n.d. Available: <https://vaksin.kemkes.go.id/#/vaccines>
- 26 Kementerian Kesehatan Republik Indonesia. Dashboard Laporan Harian COVID-19. 2022. Available: <https://dashboard.kemkes.go.id/views/LaporanHarianCovid-19v2/Topik?%3Aembed=y&%3Aiid=2&%3AisGuestRedirectFromVizportal=y>
- 27 Saptohutomo AP, KOMPAS.com. Daftar Lengkap Daerah PPKM level 1 Sampai 4 se-Indonesia Mulai 15 Maret 2022 Halaman all. 2022. Available: <https://nasional.kompas.com/read/2022/03/15/09550811/daftar-lengkap-daerah-ppkm-level-1-sampai-4-se-indonesia-mulai-15-maret-2022>
- 28 Brahmi N, Singh P, Sohal M, et al. Psychological trauma among the healthcare professionals dealing with COVID-19. *Asian J Psychiatry* 2020;54:102241.
- 29 World Health Organization. *Vaccination coverage cluster surveys: reference manual*. Geneva: WHO, 2018. Available: <https://www.who.int/publications-detail-redirect/WHO-IVB-18.09>
- 30 Randell M, Sheel M, Dynes M, et al. Influence of the COVID-19 pandemic on Caregiver beliefs and experiences of routine childhood Immunisation in Indonesia. *Vaccine* 2022;42:812.
- 31 World Health Organization. Behavioural and Social Drivers of Vaccination: Tools and Practical Guidance for Achieving High Uptake. Geneva: WHO, 2022. Available: <https://apps.who.int/iris/handle/10665/354459>
- 32 World Health Organization. *Third round of the global pulse survey on continuity of essential health services during the COVID-19 Pandemic*. Geneva: WHO, 2022. Available: https://www.who.int/publications-detail-redirect/WHO-2019-nCoV-EHS_continuity-survey-2022.1
- 33 Lang J. Complementary and integrative research (CAIR) lab. CAIR pandemic impact questionnaire (C-PIQ). NIH public health emergency and disaster research response. 2020. Available: https://www.phenxtoolkit.org/toolkit_content/PDF/CAIR_PIQ.pdf
- 34 Bandura A. Self-efficacy. In: Ramachandran V, ed. *Encyclopedia of Human Behavior*. Academic Press, 1994: 71-81.
- 35 UNICEF. The State of the World's Children 2023: For every child, vaccination. Florence: UNICEF Innocenti- Global Office of Research and Foresight, 2023.
- 36 World Health Organization. COVID-19 pandemic fuels largest continued backslide in vaccinations in three decades. 2022. Available: <https://www.who.int/news/item/15-07-2022-covid-19-pandemic-fuels-largest-continued-backslide-in-vaccinations-in-three-decades>
- 37 Shet A, Carr K, Danovaro-Holliday MC, et al. Impact of the SARS-CoV-2 pandemic on routine Immunisation services: evidence of disruption and recovery from 170 countries and territories. *Lancet Glob Health* 2022;10:e186-94.
- 38 Sharma S, Singh L, Yadav J, et al. Impact of COVID-19 on utilization of maternal and child health services in India: health management information system data analysis. *Clin Epidemiol Glob Health* 2023;21:101285.
- 39 Tilahun B, Nigusie A, Zelalem M, et al. Effect of COVID-19 pandemic on maternal and child health services and strategies for effective service implementation in Ethiopia. *J Multidiscip Healthc* 2022;15:2781-95.
- 40 Helmyati S, Dipo DP, Adiwibowo IR, et al. Monitoring continuity of maternal and child health services, Indonesia. *Bull World Health Organ* 2022;100:144-154A.
- 41 Matenge S, Sturgiss E, Desborough J, et al. Ensuring the continuation of routine primary care during the COVID-19 pandemic: a review of the International literature. *Fam Pract* 2022;39:747-61.
- 42 Nursalam N, Fibriansari RD, Yuwono SR, et al. Development of an empowerment model for burnout syndrome and quality of nursing work life in Indonesia. *Int J Nurs Sci* 2018;5:390-5.
- 43 Rusca Putra K. Prevalence of burnout syndrome among nurses in general hospitals in provincial East Java: cross-sectional study. *Enfermería Clínica* 2019;29:362-6.
- 44 Bhatti OA, Rauf H, Aziz N, et al. Violence against Healthcare workers during the COVID-19 pandemic: a review of incidents from a lower-middle-income country. *Ann Glob Health* 2021;87:41.
- 45 Ghareeb NS, El-Shafei DA, Eladl AM. Workplace violence among healthcare workers during COVID-19 pandemic in a Jordanian

- governmental hospital: the tip of the iceberg. *Environ Sci Pollut Res Int* 2021;28:61441–9.
- 46 Costa B, Pinto IC. Stress, burnout and coping in health professionals: a literature review. *J Psychol Brain Stud* 2017;1:1–8.
- 47 Nabavian M, Rahmani N, Seyed Nematollah Roshan F, et al. Nurses' experiences of the social stigma caused by the COVID-19 pandemic: a qualitative study. *J Res Nurs* 2023;28:104–15.
- 48 Cotel A, Golu F, Pantea Stoian A, et al. Predictors of burnout in healthcare workers during the COVID-19 pandemic. *Healthcare* 2021;9:304.
- 49 Mahendradhata Y, Andayani NLPE, Hasri ET, et al. The capacity of the Indonesian healthcare system to respond to COVID-19. *Front Public Health* 2021;9:649819.
- 50 Lazarus G, Meyer M, Depfenhart M, et al. Indonesian medical frontliners during the Coronavirus disease 2019 pandemic: have we been protecting them enough? *J Clin Transl Res* 2021;7:558–62.
- 51 de Guzman AB, de Castro BV, Laguilles-Villafuerte S, et al. Portrait of Filipino healthcare workers' discrimination experiences during the early part of the COVID-19 pandemic. *J Med Imaging Radiat Sci* 2022;53:396–403.
- 52 Khanal P, Devkota N, Dahal M, et al. Mental health impacts among health workers during COVID-19 in a low resource setting: a cross-sectional survey from Nepal. *Global Health* 2020;16:89.
- 53 Brewer NT, Chapman GB, Rothman AJ, et al. Increasing vaccination: putting psychological science into action. *Psychol Sci Public Interest* 2017;18:149–207.
- 54 Wiley KE, Massey PD, Cooper SC, et al. Pregnant women's intention to take up a post-Partum pertussis vaccine, and their willingness to take up the vaccine while pregnant: a cross sectional survey. *Vaccine* 2013;31:3972–8.
- 55 Le Thi Ngoc A, Dang Van C, Nguyen Thanh P, et al. Depression, anxiety, and stress among frontline health workers during the second wave of COVID-19 in Southern Vietnam: a cross-sectional survey. *PLOS Glob Public Health* 2022;2:e0000823.
- 56 Stratton E, Choi I, Peters D, et al. Co-designing a web-based decision aid tool for employees disclosure of mental health conditions: a participatory study design using employee and organizational preferences. *JMIR Form Res* 2020;4:e23337.
- 57 Yakubu K, Musoke D, Chikaphupha K, et al. An intervention package for supporting the mental well-being of community health workers in low, and middle-income countries during the COVID-19 pandemic. *Compr Psychiatry* 2022;115:152300.
- 58 López-Pineda A, Carrillo I, Mula A, et al. Strategies for the psychological support of the Healthcare workforce during the COVID-19 pandemic: the ERNST study. *Int J Environ Res Public Health* 2022;19:5529.
- 59 OECD. *Equipping health workers with the right skills: skills anticipation in the health workforce*. Paris: Organisation for Economic Co-operation and Development, 2022. Available: https://www.oecd-ilibrary.org/employment/equipping-health-workers-with-the-right-skills_9b83282e-en
- 60 World Health Organization. *The impact of COVID-19 on health and care workers: a closer look at deaths*. Geneva: World Health Organization, 2021. Available: <https://apps.who.int/iris/bitstream/handle/10665/345300/WHO-HWF-WorkingPaper-2021.1-eng.pdf>
- 61 Squires N, Garfield R, Mohamed-Ahmed O, et al. Essential public health functions: the key to resilient health systems. *BMJ Glob Health* 2023;8:e013136.
- 62 Hunter MB, Ogunlayi F, Middleton J, et al. Strengthening capacity through competency-based education and training to deliver the essential public health functions: reflection on roadmap to build public health workforce. *BMJ Glob Health* 2023;8:e011310.

Impact of COVID-19 on immunisation and maternal child health services in Indonesia

Survey for healthcare workers

	DATA COLLECTION GENERAL INFORMATION	Answer Code
A1	Province 1. Central Java 2. West Nusa Tenggara	[]
A2	District 1. Semarang City 2. Demak District 3. Purbalingga District 4. Surakarta City 5. Mataram City 6. Central Lombok District 7. Sumbawa District 8. Bima City	[]
A3	Subdistrict See code list for subdistricts	[][]
A4	Village See code list for villages	[][]
A5	Health Facility Type 1. Posyandu 2. Puskesmas/Pustu 3. Clinic 4. Independent midwife practice 5. Hospital 6. Other, specify	[]
A6	Selected respondent number	[][]
A7	Unique interview ID	[][]
A8	Substitute respondent	[]yes []no
A9	Interview date	__/__/__
A10	Interview start time	__ __
A11	Interview finish time	__ __
A12	Interviewer name	
A13	Interviewer Initials	
	CHECKING STATUS AND DATA ENTRY	
B1	Checking date by field coordinator	__/__/__
B2	Field coordinator name	
B3	Field coordinator initials	
B4	Date of data entry	__/__/__
B5	Data entry officer initials	

	Section 1: Demographics	Response code
--	-------------------------	---------------

C0	Could you tell me your name, for the purposes of this interview?	
C1	Gender of respondent <i>[interviewer to observe only]</i> 1. Female 2. Male	[]
C2	How old are you? <input type="text"/> <input type="text"/> years	[]
C3	C3. What is your highest level of education? 1. Completed high school 2. Completed a diploma (D3) 3. Completed diploma (D4) 4. Undergraduate degree 5. Profession degree 6. Masters 7. PhD	[]
C4	What is your current role? 1. Doctor 2. Nurse 3. Midwife 4. Allied health 5. Community health worker/village midwife 6. Traditional healer 7. Other, specify	[]
C5	How many years have you worked in this professional role? 1. Less than 2 years 2. Between 2 and less than 5 years 3. Between 5 and less than 10 years 4. 10 years or more	[]
C6	What health facility do you primarily work in? 1. Puskesmas (<i>continue to C7</i>) 2. Poskesdes (<i>continue to C7</i>) 3. Polindes (<i>continue to C7</i>) 4. Pustu (<i>continue to C7</i>) 5. Independent midwife practice (<i>continue to C7</i>) 6. Private clinic (<i>continue to C7 & C8</i>) 7. Other, specify (<i>continue to C7</i>)	[]
C7	Do you work in any other health facility? <i>[Can have more than one response. Mark (write the number 1 on all the answers mentioned by the respondent). The interviewer MUST NOT read out the answer options, but select the option mentioned which is closest]</i>	
	A. No	A.
	B. Yes, Puskesmas	B.
	C. Yes, Poskesdas	C.
	D. Yes, Pustu	D.

	E. Yes, independent midwife practice	E.
	F. Yes, private clinic	F.
	G. Other, specify	G.
C8	<p><i>[If respondent answered 'Yes' for private clinic in C6 OR C7, ask the question below. If the respondent does not work in a private clinic continue to C9]</i></p> <p>How would you describe the private clinic you work in. Is it: <i>[Can have more than one response. Mark (write the number 1 on all the answers mentioned by the respondent). The interviewer MUST NOT read out the answer options, but select the option mentioned which is closest]</i></p>	
	A. Not for profit - Religiously affiliated	A.
	B. Not for profit – not religiously affiliated	B.
	C. For profit	C.
	D. Other, specify	D.
	E. Don't know	E.
C9	<p>Which health services do you support? <i>[Can have more than one response. Mark (write the number 1 on all the answers mentioned by the respondent). The interviewer MUST NOT read out the answer options, but select the option mentioned which is closest]</i></p>	
	A. Routine child Immunization	A.
	B. COVID-19 Immunization	B.
	C. Antenatal care	C.
	D. Labor and birth	D.
	E. Postnatal care	E.
	F. Family planning	F.
	G. Baby and child services (weighing, vitamin A supplementation, etc)	G.
	H. Other, specify	H.
Section 2: Impact of COVID-19 on Routine Immunisation and MCH services		
<i>If the respondent did not select 'routine child immunisation' or 'COVID-19 immunisation' in Section 1, C9, proceed to D8.</i>		
D1	<p>Do you have systems in place to track people who miss out on routine vaccines or defaulters?</p> <p>1. No 2. Yes 3. Not sure</p>	[]
D2	<p>In the past year have you had a stock-out of any routine vaccines?</p> <p>1. No (<i>skip to D4</i>) 2. Yes (<i>continue to D3</i>) 3. Not sure (<i>skip to D4</i>)</p>	[]

D3	Which vaccines have been stocked out? <i>[Can have more than one response. Mark (write the number 1 on all the answers mentioned by the respondent). The interviewer MUST NOT read out the answer options, but select the option mentioned which is closest]</i>	
	A. BCG	A.
	B. HepB	B.
	C. OPV	C.
	D. IPV	D.
	E. Pentavalent (DPT-HB-Hib)	E.
	F. PCV	F.
	G. MMR/MR	G.
	H. Other, specify	H.
D4	In the past year have you had a stock out of any injection equipment? <i>[Can have more than one response. Mark (write the number 1 on all the answers mentioned by the respondent). The interviewer MUST NOT read out the answer options, but select the option mentioned which is closest]</i>	
	A. No	A.
	B. Yes, syringes	B.
	C. Yes, needles	C.
	D. Yes, diluent	D.
	E. Don't know	E.
D5	Apart from your basic professional training, have you had any formal training in immunisation? 1. No 2. Yes	[]
D6	Do you have a copy of the standard operating procedure for routine immunisation in your clinic? 1. No 2. Yes 3. Don't know	[]
D7	If required, would it bother you to give more than one injection during a vaccine encounter? 1. No 2. Yes	[]
D8	In the facility that you primarily work in (service mentioned earlier), did you experience any of the following disruptions due to COVID-19?	

	8a. Closure of the service 1. No 2. Yes 3. Don't know	[]
	8b. Lockdowns hindering access to the service for parents or caregivers 1. No 2. Yes 3. Don't know	[]
	8c. Fewer parents/caregivers presenting to service for reasons other than lockdown 1. No 2. Yes 3. Don't know	[]
	8d. Not enough staff to provide services 1. No 2. Yes 3. Don't know	[]
	8e. Staff deployed to provide COVID-19 relief 1. No 2. Yes 3. Don't know	[]
	8f. Insufficient personal protective equipment (PPE) available for health care providers to provide services 1. No 2. Yes 3. Don't know	[]
	8g. Unavailability/stock out of vaccines or injecting equipment at service 1. No 2. Yes 3. Don't know	[]
	8h. Changes in policies for who can be vaccinated 1. No 2. Yes 3. Don't know	[]
	8i. Others (please specify what are the other causes of this disruption and/or changes in service utilization):	
D9	In the facility that you primarily work in (service mentioned earlier), what approaches were used to overcome the disruptions due to COVID-19? Did you see.....	
	9a. Task shifting / role delegation 1. No 2. Yes 3. Don't know	[]
	9b. Redirection of parents or caregivers to alternative health care facilities 1. No 2. Yes	[]

	3. Don't know	
	9c. Community outreach to explain service disruptions and changes 1. No 2. Yes 3. Don't know	[]
	9d. Patients or caregivers seen outdoors from the facility 1. No 2. Yes 3. Don't know	[]
	9e. Government removal of user fees 1. No 2. Yes 3. Don't know	[]
	9f. Phone calls to patients or caregivers 1. No 2. Yes 3. Don't know	[]
	9g. Others, describe what other approaches are being used	[]
Section 3: COVID-19		
E1	Have you ever had COVID-19? 1. No (<i>skip to E6</i>) 2. Yes (<i>continue to E2</i>) 3. Not sure (<i>skip to E6</i>)	[]
E2	In your opinion, based on the symptoms you suffered/felt, was it mild, moderate or severe? 1. Mild 2. Moderate 3. Severe 4. Not Sure	[]
E3	Was it confirmed by a test? 1. Confirmed by a test (<i>continue to E4</i>) 2. Not confirmed by a test (<i>Skip to E5</i>)	[]
E4	If confirmed by tests, what tests were performed? <i>[Can have more than one response. Mark (write the number 1 on all the answers mentioned by the respondent). The interviewer MUST NOT read out the answer options, but select the option mentioned which is closest]</i>	
	A. PCR test	A.
	B. Antigen test	B.
	C. Antibody test	C.
	D. Other, specify	D.
	E. Don't know	E.
E5	Were you hospitalized? 1. No 2. Yes	[]

E6	Has anyone you know had COVID-19? 1.. No 2. Yes, family only 3. Yes, other people only (not family) 4. Yes, family and other people 5. Not sure	[]
E7	Do you know where to go to get a COVID-19 vaccine for yourself? 1. No 2. Yes	[]
E8	Have you received a COVID-19 vaccine? Would you say... 1. No (<i>skip to E11</i>) 2. Yes, you received one dose (<i>continue to E9</i>) 3. Yes, you received two doses (<i>continue to E9</i>) 4. Yes, you received three or more doses? (<i>continue to E9</i>) 5. Not sure (<i>skip to E11</i>)	[]
E9	Do you still need another dose of COVID-19 vaccine? Would you say... 1. No, you do <u>not</u> need another dose 2. Yes, you do need another dose or doses 3. Not sure	[]
E10	Do you have a vaccine card/certificate? 1. Yes, showable (peduli lindungi/card) (<i>ask to view the card</i>) 2. Yes, but can't be shown 3. No	[]
E11	<i>[If respondent is not vaccinated – see E8]</i> Do you want to get a COVID-19 vaccine? Would you say... 1. No, you do not want to 2. Yes, you do want to, or are you 3. Not sure	[]
E12	Do you think most of your close family and friends would want you to get a COVID-19 vaccine? 1. No 2. Yes	[]
E13a	How easy is it to get a COVID vaccine for yourself? Would you say... 1. Not at all easy 2. A little easy 3. Moderately easy 4. Very easy	[]
E13b	What makes it hard for you to get a COVID-19 vaccine? Would you say... <i>[Can have more than one response. Mark (write the number 1 on all the answers mentioned by the respondent). The interviewer may read out the answer options]</i>	
	A. COVID-19 vaccination is not yet available for me	A.
	B. Making an appointment is hard	B.
	C. I can't go on my own (I have a physical limitation)	C.
	D. The vaccination site is hard to get to	D.
	E. The opening times are inconvenient	E.

	F. Sometimes people are turned away without vaccination	F.
	G. The waiting time takes too long	G.
	H. Something else, specify	H.
	I. Nothing. It's not hard	I.
E14	How cheap is it to pay for COVID-19 vaccination? When you think about the cost, please consider any payments to the clinic, the cost of getting there, plus the cost of taking time away from work. Would you say... 1. Not at all cheap 2. A little easy cheap 3. Moderately cheap 4. Very cheap	[]
E15	How concerned are you about getting COVID-19? Would you say... 1. Not at all concerned 2. A little concerned 3. Moderately concerned 4. Very concerned	[]
E16	How concerned are you about your patients getting COVID-19 from you? Would you say... 1. Not at all concerned 2. A little concerned 3. Moderately concerned 4. Very concerned	[]
E17	How important do you think getting a COVID-19 vaccination will be for your health? Would you say... 1. Not at all important 2. A little important 3. Moderately important 4. Very important	[]
E18	How confident are you that you could answer patient questions about getting a COVID-19 vaccine? Would you say... 1. Not at all confident 2. A little confident 3. Moderately confident 3. Very confident	[]
E19	In the last year, have you seen or heard anything that made you worry about the COVID-19 vaccine? 1. No 2. Yes	[]
E20	Have you encountered information on COVID-19 where you found it hard to decide whether it was right or wrong? For example, information about ways to prevent the disease or to recover? 1. No (<i>skip to E22</i>) 2. Yes (<i>continue to E21</i>)	[]
E21	Where did you see or hear this information? <i>[Can have more than one response. Mark (write the number 1 on all the answers mentioned by the respondent). The interviewer MUST NOT read out the answer options, but select the option mentioned which is closest]</i>	
	A. Via an online search	A.

	B. Facebook	B.
	C. WhatsApp	C.
	D. YouTube	D.
	E. Twitter	E.
	F. TikTok	F.
	G. Instagram	G.
	H. Snapchat	H.
	I. Word of mouth	I.
	J. Other, specify	J.
E22	How confident are you that you could respond to misinformation about the COVID-19 vaccine? 1. Not at all confident 2. A little confident 3. Moderately confident 4. Very confident	[]
E23	Apart from your basic professional training, have you had any formal training on the COVID-19 vaccination? 1. No 2. Yes	[]
E24	In the past year have you had a stock-out of the COVID-19 vaccines? 1. No 2. Yes 3. Not sure 4. Do not stock the COVID-19 vaccine	[]
E25	Has delivering COVID-19 vaccination taken you away from other priorities? 1. Not at all 2. Slightly 3. Moderately 4. Very 5. I have not delivered the COVID-19 vaccination	[]
E26	Have you been treated poorly during the COVID-19 pandemic because you are a health worker? 1. No (<i>skip to E28</i>) 2. Yes (<i>continue to E27</i>) 3. Not sure (<i>skip to E28</i>)	[]
E27	What were the reasons for this? <i>[Can have more than one response. Mark (write the number 1 on all the answers mentioned by the respondent). The interviewer MUST NOT read out the answer options, but select the option mentioned which is closest]</i>	
	A. Patients angry at service closures	A.
	B. Patients angered by the vaccine brands offered	B.
	C. Patients angered by request for medical exemption	C.
	D. Patients angered by longer wait times	D.
	E. Patients believe I am a COVID-19 risk	E.
	F. Other, specify	F.

E28	Would you say you've experienced trauma or burnout related to the COVID-19 pandemic? 1. No (<i>skip to E30</i>) 2. Yes (<i>continue to E29</i>) 3. Not sure (<i>skip to E30</i>)	[]
E29	What are the main reasons you have felt trauma or burnout in your job related to COVID-19? <i>[Can have more than one response. Mark (write the number 1 on all the answers mentioned by the respondent). The interviewer MUST NOT read out the answer options, but select the option mentioned which is closest]</i>	
	A. Shifting work duties	A.
	B. Changes in service delivery	B.
	C. Long work hours	C.
	D. Patient distress	D.
	E. Use of PPE	E.
	F. Lack of access to PPE	F.
	G. Fear of illness	G.
	H. Fear of making family ill	H.
	I. Training requirements	I.
	J. Other, specify	J.
E30	Do you feel like your job has increased risk of exposure to COVID-19? 1. No 2. Yes	[]
E31	How much has your mental/emotional health been worsened by the COVID-19 pandemic? 1. Not at all 2. Slightly 3. Moderately 4. Very 5. Extremely	[]
Section 4: Self-efficacy		
<i>Now I will ask you some questions, each with four responses and would like to think about how true these are in regard to your current role. (Use Option Card/Kartu Bantu)</i>		
F1	I can always manage to solve difficult problems if I try hard enough 1. Not true at all 2. Hardly true 3. Moderately true 4. Exactly true	[]
F2	If someone opposes me, I can find the means and ways to get what I want. 1. Not true at all 2. Hardly true 3. Moderately true 4. Exactly true	[]
F3	It is easy for me to stick to my aims and accomplish my goals. 1. Not true at all	[]

	<p>2. Hardly true 3. Moderately true 4. Exactly true</p>	
F4	<p>I am confident that I could deal efficiently with unexpected events. 1. Not true at all 2. Hardly true 3. Moderately true 4. Exactly true</p>	[]
F5	<p>Thanks to my resourcefulness, I know how to handle unforeseen situations. 1. Not true at all 2. Hardly true 3. Moderately true 4. Exactly true</p>	[]
F6	<p>I can solve most problems if I invest the necessary effort. 1. Not true at all 2. Hardly true 3. Moderately true 4. Exactly true</p>	[]
F7	<p>I can remain calm when facing difficulties because I can rely on my coping abilities. 1. Not true at all 2. Hardly true 3. Moderately true 4. Exactly true</p>	[]
F8	<p>When I am confronted with a problem, I can usually find several solutions. 1. Not true at all 2. Hardly true 3. Moderately true 4. Exactly true</p>	[]
F9	<p>If I am in trouble, I can usually think of a solution. 1. Not true at all 2. Hardly true 3. Moderately true 4. Exactly true</p>	[]
F10	<p>I can usually handle whatever comes my way. 1. Not true at all 2. Hardly true 3. Moderately true 4. Exactly true</p>	[]

Supplementary material: Reflexivity statement**Manuscript titled:**

Service disruptions, trauma and burnout during the COVID-19 pandemic amongst healthcare providers delivering immunisation and maternal and child health services in Indonesia

Study conceptualisation	<p><i>How does this study address local research and policy priorities?</i></p> <p>The study was commissioned by UNICEF regional office but was developed in consultation with the country office in Indonesia and based on local epidemiological situation and what was happening in the country. In 2021, global reports had already presented the impact of the pandemic on essential health services in Indonesia and elsewhere. We engaged our research partners from Universitas Indonesia during the development of the funding proposal. Through this, we were able to also discuss the needs with our research partners based on their experience prior to proceeding. For example, the provinces in which the study was conducted was based on their knowledge and expertise of the epidemiological situation in the country, provinces, feasibility and impact/ relevance of the findings. This study was conducted with the Indonesian Ministry of Health forming part of the Technical Advisory Group to ensure that it continued to support local research and study priorities.</p>
	<p><i>How were local researchers involved in study design?</i></p> <p>Throughout the course of the study, the research team worked collaboratively and in equal partnership with the local research team from Universitas Indonesia. The collaborators from USYD (lead PI) met with the team ahead of the grant submission; and throughout design and implementation – the two lead institutions worked together to develop and implement the study. The USYD team worked proactively to ensure participation and engagement from Indonesia research team and MOH was meaningful and respectful – we used emails, regular zoom and Whatsapp. Noting that the COVID-19 pandemic was still raging when the study was designed and implemented, in-country visits by the international team weren't possible. It was also considered more appropriate for the in-country engagement to be led by our in-country partners and UNICEF country officers, both government departments as well as other local stakeholders.</p>
Research management	<p><i>How has funding been used to support the local research team(s)?</i></p> <p>At the inception stage, the Indonesian research team developed a budget to undertake the research including fieldwork activities, which were dedicated to supporting local</p>

	<p>research teams. It covered team salaries as well as costs associated with data collection processes. The in-country research team was completely responsible for governance and research management.</p>
Data acquisition and analysis	<p><i>How are research staff who conducted data collection acknowledged?</i></p> <p>The research staff who conducted data collection have also played a role as co-authors in this and other manuscripts and have been acknowledged accordingly. Individuals who supported data collection as enumerators and field coordinators have been formally acknowledged in the “Acknowledgements” section of the manuscript.</p>
	<p><i>How have members of the research partnership been provided with access to study data?</i></p> <p>The local research team lead the data collection as well as data management processes. This meant that data was shared from local research team members to international team members through a secure portal in line with the ethics process.</p>
	<p><i>How were data used to develop analytical skills within the partnership?</i></p> <p>The local research team’s leading role in data management provided opportunities for team members to develop skills in processes such as data entry and data cleaning to ensure that we had a high-quality data set to use for analysis. Data used in this manuscript was part of a broader study. In reporting of this study, local team members also helped to provide initial descriptive analyses.</p>
Data interpretation	<p><i>How have research partners collaborated in interpreting study data?</i></p> <p>Local research partners have played an important role in helping to interpret study data in context, providing insights which supported how we interpreted our findings in way that was relevant to Indonesia. The results were also presented to key stakeholders including ministry of health, UNICEF to ensure interpretation of the results was relevant to the local context.</p>
Drafting and revising for intellectual content	<p><i>How were research partners supported to develop writing skills?</i></p> <p>Our local research team have been involved as co-authors in all manuscripts either published or under review for publication which has supported opportunities to develop writing skills. The research partner team also wrote sections of the original protocol, reports and papers and feedback was provided by the study team.</p>

	<p><i>How will research products be shared to address local needs?</i></p> <p>The study findings were shared through multiple modalities – through UNICEF to the Indonesian Ministry of Health; through presentations and also CIs (JL and MS) met with multiple in-country personnel (for example, NITAG Chair and other forums) and presented findings through small discussions’ and 1-to-1 presentations. A national large meeting could not be organised due to competing priorities of colleagues in Indonesia but complete efforts have been made to share the findings and disseminate widely. A report was also prepared which shared with in-country government counterparts for early feedback, contextual input etc prior to any formal dissemination. Further, we have involved Ministry of Health team members as co-authors on all published manuscripts or those under review for publication to ensure that they can support the dissemination of these findings to help address local needs.</p>
<p>Authorship</p>	<p><i>How is the leadership, contribution and ownership of this work by LMIC researchers recognised within the authorship?</i></p> <p>All local research team members have been included as co-authors in published manuscripts or those under review for publication. The lead investigator for the local research team at Universitas Indonesia is the senior author for another paper published from the broader study. He is second author on this paper. Future papers (for example a qualitative study) will have other authors from the Indonesia team in other prominent places depending on contribution.</p>
	<p><i>How have early career researchers across the partnership been included within the authorship team?</i></p> <p>A number of the local research team members are early career researchers and have been included as co-authors on all manuscripts. The first author (MR) and author 4 (AT) for this manuscript is an early career researcher. The corresponding researcher is a mid-career researcher (MS) is an inter-sectional researcher from the global south. Future papers are also expected to be led by early career researchers across the research partnership.</p>
	<p><i>How has gender balance been addressed within the authorship?</i></p> <p>The majority of authors on this manuscript (14/17) and others developed as part of a broader study are female.</p>

Training	<i>How has the project contributed to training of LMIC researchers?</i> Training on research methods and ethics was provided by the international team (along with co-PI from the local team) ahead of fieldwork. The local research team implemented training on data collection processes to locally engaged enumerators, offering important capacity building opportunities to researchers at a district level.
Infrastructure	<i>How has the project contributed to improvements in local infrastructure?</i> Limited infrastructure was used for this study as it was a low-cost survey and did not involve any major equipment or infrastructure.
Governance	<i>What safeguarding procedures were used to protect local study participants and researchers?</i> Data collection protocols were designed by the local research team and received local ethics approval. Further government approvals were granted from the provincial and district governments where data collection took place. As data collection occurred during the COVID-19 pandemic, COVID-19 protocols were enforced to ensure the safety of participants and local researchers.