

**Table S2:** Mortality prevalence by birthweight category among the inborn neonates in the NBUs of the 16 CIN hospitals (Population B)

	<b>&lt; 1 Kg</b>	<b>1 - &lt; 1.5 Kg</b>	<b>1.5 - &lt; 2 Kg</b>	<b>2 - &lt; 2.5 Kg</b>	<b>2.5 - 4 Kg</b>	<b>&gt; 4 Kg</b>	<b>Total</b>
H1	85% (11/13)	58% (25/43)	24% (34/142)	13% (18/138)	12% (16/568)	14% (8/57)	<b>17% (165/961)</b>
H2	71% (12/17)	34% (30/87)	17% (21/123)	17% (13/75)	10% (46/468)	8% (3/37)	<b>15% (125/807)</b>
H3	95% (39/41)	47% (51/108)	19% (39/204)	12% (36/306)	6% (139/2340)	5% (5/110)	<b>10% (309/3109)</b>
H4	81% (25/31)	41% (33/80)	11% (14/132)	6% (7/119)	9% (38/440)	6% (2/35)	<b>14% (119/837)</b>
H5	87% (45/52)	50% (93/187)	18% (60/337)	8% (27/323)	12% (151/1275)	3% (9/297)	<b>16% (385/2471)</b>
H6	79% (30/38)	40% (31/78)	11% (19/178)	10% (25/249)	5% (67/1312)	3% (3/86)	<b>9% (175/1941)</b>
H7	84% (52/62)	60% (100/168)	19% (65/335)	11% (48/442)	6% (132/2329)	2% (5/232)	<b>11% (402/3568)</b>
H8	71% (15/21)	38% (19/50)	15% (16/108)	5% (6/131)	6% (35/593)	0% (0/52)	<b>10% (91/955)</b>
H9	79% (31/39)	36% (47/130)	22% (51/236)	14% (22/162)	12% (97/777)	6% (3/47)	<b>18% (251/1391)</b>
H10	53% (41/77)	32% (58/181)	7% (27/390)	5% (29/551)	3% (106/3616)	3% (7/258)	<b>5% (286/5073)</b>
H11	87% (85/98)	49% (143/293)	14% (80/566)	9% (45/512)	7% (164/2240)	1% (5/461)	<b>13% (522/4170)</b>
H12	67% (44/66)	32% (55/174)	12% (37/306)	10% (25/241)	10% (95/905)	0% (0/55)	<b>15% (256/1747)</b>
H13	80% (28/35)	38% (36/94)	12% (24/206)	5% (15/275)	5% (63/1351)	0% (1/219)	<b>8% (167/2180)</b>
H14	86% (6/7)	50% (20/40)	11% (12/109)	13% (11/84)	8% (30/380)	0% (0/21)	<b>12% (79/641)</b>
H15	86% (31/36)	44% (45/103)	16% (260/41)	10% (37/389)	4% (73/1849)	1% (2/296)	<b>8% (229/2933)</b>
H16	81% (73/90)	42% (125/297)	12% (88/730)	7% (73/1007)	4% (223/5898)	2% (7/360)	<b>7% (589/8382)</b>
<b>Total</b>	<b>79% (568/723)</b>	<b>43% (911/2113)</b>	<b>14% (628/4362)</b>	<b>9% (437/5004)</b>	<b>6% (1528/26341)</b>	<b>2% (60/2623)</b>	<b>10% (4132/41166)</b>