

Supplementary table 7. Effect of providing BCG and OPV vaccines at a home visit on growth

	MUAC enrolment Mean (SD)	MUAC follow-up* Mean (SD)	Difference in MUAC at follow-up (95% CI)	WAZ enrolment Z-score (SD)	WAZ follow-up* Z-score (SD)	Difference in WAZ at follow-up (95% CI)
Intervention	98 (9)	118 (14)	0.82 (-0.39 - 2.03)	-0.78 (1.01)	-0.50 (1.08)	0.04 (-0.05 - 0.13)
Control	97 (10)	115 (14)	Reference	-0.91 (1.04)	-0.67 (1.10)	Reference
Rural						
Intervention	97 (9)	114 (13)	1.02 (-0.43 - 2.46)	-0.83 (1.02)	-0.59 (1.11)	0.05 (-0.06 - 0.17)
Control	97 (10)	112 (13)	Reference	-0.96 (1.03)	-0.76 (1.10)	Reference
Urban						
Intervention	98 (8)	126 (10)	-0.28 (-2.48 - 1.91)	-0.67 (0.98)	-0.33 (1.01)	-0.02 (-0.19 - 0.15)
Control	99 (11)	126 (11)	Reference	-0.70 (1.02)	-0.33 (1.06)	Reference

MUAC: Mid-upper-arm circumference, SD: Standard deviation, CI: Confidence interval, WAZ: Weight-for-age Z-score

* Measured at first follow-up visit within 6 months of age.